

Colorado Department of Health Care Policy and Financing Preferred Drug List (PDL) Effective January 1, 2026

Prior Authorization Forms: Available online at <https://hcpf.colorado.gov/pharmacy-resources>

Prior Authorization (PA) Requests: Colorado Pharmacy Call Center Phone Number: 800-424-5725 | Fax Number: 800-424-5881

Electronic Prior Authorization (ePA): Electronic Prior Authorization Requests are supported by CoverMyMeds and may be submitted via Electronic Health Record (EHR) systems or through the CoverMyMeds provider portal.

The PDL applies to Medicaid fee-for-service members. It does not apply to members enrolled in Rocky Mountain Health HMO or Denver Health Medicaid Choice.

Initiation of pharmaceutical product subject to Prior Authorization: Please note that starting the requested drug, including a non-preferred drug, prior to a PA request being reviewed and approved, through either inpatient use, by using office “samples,” or by any other means, does not necessitate Medicaid approval of the PA request.

Health First Colorado, at section 25.5-5-501, C.R.S., requires the generic of a brand name drug be prescribed if the generic is therapeutically equivalent to the brand name drug. Exceptions to this rule are: 1) If the brand name drug is more cost effective than the generic as determined by the Department, 2) If the patient has been stabilized on a brand name drug and the prescriber believes that transition to a generic would disrupt care, and 3) If the drug is being used for treatment of mental illness, cancer, epilepsy, or human immunodeficiency virus and acquired immune deficiency syndrome.

Please see the [Brand Favored Product List](#) for a list of medications where the brand name drug is more cost effective than the generic drug.

A provider may request a step therapy exception for the treatment of a serious or complex medical condition pursuant to section 25.5-4-428, C.R.S. Serious or complex medical condition means the following medical conditions: serious mental illness, cancer, epilepsy, multiple sclerosis, or human immunodeficiency virus (HIV)/ acquired immune deficiency syndrome (AIDS), or a condition requiring medical treatment to avoid death, hospitalization, or a worsening or advancing of disease progression resulting in significant harm or disability. The step therapy exception request form is available by visiting <https://hcpf.colorado.gov/pharmacy-resources>

Brand Name Required = BNR, Prior Authorization = PA, AutoPA = authorization can be automated at the point-of-sale transaction if criteria are met
Preferred drug list applies only to prescription (RX) products, unless specified.

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-preferred products will be approved for one year unless otherwise stated.)
<div>I. Analgesics</div>		
Therapeutic Drug Class: NON-OPIOID ANALGESIA AGENTS - Oral – <i>Effective 4/1/2025</i>		
<div>No PA Required</div> <div>Duloxetine 20 mg, 30 mg, 60 mg capsule</div> <div>Gabapentin capsule, tablet, solution</div> <div>Pregabalin capsule</div> <div>SAVELLA (milnacipran) tablet, titration pack</div>	<div>PA Required</div> <div>CYMBALTA (duloxetine) capsule</div> <div>DRIZALMA (duloxetine DR) sprinkle capsules</div> <div>Duloxetine 40 mg capsule</div> <div>GRALISE (gabapentin ER) tablet</div> <div>Gabapentin ER tablet</div> <div>HORIZANT (gabapentin ER) tablet</div> <div>JOURNAVX (suzetrigine) tablet</div> <div>LYRICA (pregabalin) capsule, solution, CR tablet</div> <div>NEURONTIN (gabapentin) capsule, solution, tablet</div> <div>Pregabalin solution, ER tablet</div>	<div>JOURNAVX (suzetrigine) may be approved if the following criteria are met:</div> <div><ul style="list-style-type: none">Member is ≥ 18 years of age ANDMember is being prescribed suzetrigine for up to 14 days of treatment for moderate-to- severe acute pain ANDPrescriber attests that the member’s pain is unable to be managed with an NSAID, acetaminophen, or other non-opioid analgesic ANDJournavx (suzetrigine) is not being prescribed to treat chronic pain ANDThe medication is not being prescribed to treat pain associated with migraine ANDMember does not have severe hepatic impairment (Child-Pugh Class C) ANDMember has been counseled to avoid food or drink containing grapefruit during treatment with Journavx (suzetrigine) ANDMember is not concurrently taking a strong CYP3A inhibitor (such as ketoconazole, itraconazole, posaconazole, ritonavir, indinavir, saquinavir, clarithromycin, fluvoxamine) ANDMember is not concurrently taking a strong or moderate CYP3A inducer (such as carbamazepine, phenytoin, rifampin, efavirenz, rifabutin, St. John’s Wort) · Members using hormonal contraceptives containing progestins other than levonorgestrel and norethindrone have been counseled regarding alternative or additional contraception, if appropriate, per product labeling.</div> <div><div>Duration of Approval: 3 months</div><div>Dosing Limit: One 14-day course per approval on file</div><div>Quantity limit: 29 tablets/14 days</div></div> <div>All other non-preferred oral non-opioid analgesic agents may be approved if member meets all of the following criteria:</div> <div><ul style="list-style-type: none">Member has trialed and failed duloxetine (20mg, 30mg, or 60mg) AND has trialed and failed gabapentin OR pregabalin capsule (Failure is defined as lack of efficacy with 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction)</div> <div>Prior authorization will be required for Lyrica (pregabalin) capsule dosages > 600mg per day (maximum of 3 capsules daily) and gabapentin dosages > 3600mg per day.</div>
Therapeutic Drug Class: NON-OPIOID ANALGESIA AGENTS - Topical – <i>Effective 4/1/2025</i>		
<div>No PA Required</div> <div>Lidocaine patch</div> <div>LIDODERM (lidocaine) patch</div>	<div>PA Required</div> <div>Lidocaine patch (Puretek)</div> <div>ZTLIDO (lidocaine) topical system</div>	<div>Non-preferred topical products require a trial/failure with an adequate 8-week trial of gabapentin AND pregabalin AND duloxetine AND a preferred lidocaine 5% patch. Failure is defined as lack of efficacy with an 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction.</div> <div><div>Lidocaine 5% patch (<i>Puretek manufacturer only</i>) may be approved if the following criteria are met:</div><div><ul style="list-style-type: none">Member is ≥ 18 years of age AND</div></div>

		<ul style="list-style-type: none">Member has had an adequate 8-week trial and failure of: gabapentin AND pregabalin AND duloxetine AND a preferred lidocaine 5% patch. Failure is defined as lack of efficacy with an 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction ANDPrescriber has provided a justification of clinical necessity indicating that an alternative generic lidocaine 5% patch formulation cannot be used.
Therapeutic Drug Class: NON-STEROIDAL ANTI-INFLAMMATORIES (NSAIDs) - Oral – <i>Effective 4/1/2025</i>		
No PA Required	PA Required	
Celecoxib capsule	ARTHROTEC (diclofenac sodium/ misoprostol) tablet	<p>DUEXIS (ibuprofen/famotidine) or VIMOVO (naproxen/esomeprazole) may be approved if the member meets the following criteria:</p> <ul style="list-style-type: none">Trial and failure[‡] of all preferred NSAIDs at maximally tolerated doses ANDTrial and failure[‡] of three preferred proton pump inhibitors in combination with NSAID within the last 6 months ANDHas a documented history of gastrointestinal bleeding <p>Diclofenac potassium 25 mg immediate-release tablets may be approved if the following criteria are met:</p> <ul style="list-style-type: none">Member is ≥ 18 years of age ANDMember does not have any of the following medical conditions:<ul style="list-style-type: none">History of recent coronary artery bypass graft (CABG) surgeryHistory of myocardial infarctionSevere heart failureAdvanced renal diseaseHistory of gastrointestinal bleeding <p>AND</p> <ul style="list-style-type: none">Member has trial and failure[‡] of four preferred oral NSAIDs at maximally tolerated doses <p>ELYXYB (celecoxib) oral solution may be approved if the following criteria are met:</p> <ul style="list-style-type: none">Member is ≥ 18 years of age ANDRequested medication is being prescribed for acute treatment of migraine (with or without aura) ANDMember does <u>not</u> have any of the following medical conditions:<ul style="list-style-type: none">History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDsHistory of recent coronary artery bypass graft (CABG) surgeryHistory of allergic-type reactions to sulfonamidesSevere heart failureHistory of myocardial infarctionHistory of gastrointestinal bleedingAdvanced renal diseasePregnancy past 30 weeks gestation <p>AND</p> <ul style="list-style-type: none">Member is unable to take an alternative NSAID in a solid oral dosage form ANDMember has tried and failed[†] one preferred NSAID oral liquid ANDMember is unable to use celecoxib capsules, opened and sprinkled into applesauce or other soft food
Diclofenac potassium 50 mg tablet	CELEBREX (celecoxib) capsule	
Diclofenac sodium EC/DR tablet	COMBOGESIC (Ibuprofen/Acetaminophen) tablet	
Ibuprofen suspension, tablet (RX)	DAYPRO (oxaprozin) caplet	
Indomethacin capsule, ER capsule	Diclofenac potassium capsule, powder pack	
Ketorolac tablet*	Diclofenac potassium 25 mg tablet	
Meloxicam tablet	Diclofenac sodium ER/SR tablet	
Nabumetone tablet	Diclofenac sodium/misoprostol tablet	
Naproxen DR/ER, tablet (RX)	Diflunisal tablet	
Naproxen suspension	DUEXIS (ibuprofen/famotidine) tablet	
Sulindac tablet	ELYXYB (celecoxib) solution	
	Etodolac capsule; IR, ER tablet	
	FELDENE (piroxicam) capsule	
	Fenoprofen capsule, tablet	
	Flurbiprofen tablet	
	Ibuprofen/famotidine tablet	

	<div>Ibuprofen 300 mg tablet</div> <div>Ketoprofen IR, ER capsule</div> <div>LOFENA (diclofenac) tablet</div> <div>Meclofenamate capsule</div> <div>Mefenamic acid capsule</div> <div>Meloxicam submicronized capsule, suspension</div> <div>NALFON (fenoprofen) capsule, tablet</div> <div>NAPRELAN (naproxen CR) tablet</div> <div>Naproxen sodium CR, ER, IR tablet</div> <div>Naproxen/esomeprazole DR tablet</div> <div>Oxaprozin tablet</div> <div>Piroxicam capsule</div> <div>RELAFEN DS (nabumetone) tablet</div> <div>Tolmetin tablet</div> <div>VIMOVO (naproxen/esomeprazole) DR tablet</div>	<div>[†]Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions.</div> <div><u>Maximum dose:</u> 120 mg/day</div> <div>All other non-preferred oral agents may be approved following trial and failure[‡] of four preferred agents. [‡]Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions.</div> <div>*Ketorolac tablets quantity limit: 5-day supply per 30 days and 20 tablets per 30 days</div>
Therapeutic Drug Class: NON-STEROIDAL ANTI-INFLAMMATORIES (NSAIDS) - Non-Oral – <i>Effective 4/1/2025</i>		
<div>No PA Required</div> <div>Diclofenac 1.5% topical solution</div> <div>Diclofenac sodium 1% gel (OTC/Rx)</div>	<div>PA Required</div> <div>Diclofenac 1.3% topical patch, 2% pump</div> <div>FLECTOR (diclofenac) 1.3% topical patch</div> <div>Ketorolac nasal spray</div>	<div>SPRIX (ketorolac) may be approved if meeting the following criteria:</div> <div><ul style="list-style-type: none">Member is unable to tolerate, swallow or absorb oral NSAID formulations ORMember has trialed and failed three preferred oral or topical NSAID agents (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions)Quantity limit: 5-single day nasal spray bottles per 30 days</div> <div>All other non-preferred topical agents may be approved for members who have trialed and failed one preferred agent. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction.</div> <div>Diclofenac topical patch quantity limit: 2 patches per day</div>

	LICART (diclofenac) 1.3% topical patch PENNSAID (diclofenac solution) 2% pump, 2% solution packet	Diclofenac 3% gel (generic Solaraze) prior authorization criteria can be found in the Antineoplastic agents, topical, section of the PDL.
<p>Opioid Utilization Policy (long-acting and short-acting opioids):</p> <p>It is highly encouraged that the healthcare team utilize the Prescription Drug Monitoring Program (PDMP) to aid in ensuring safe and efficacious therapy for members using controlled substances.</p> <p><u>Total Morphine Milligram Equivalent Policy Effective 10/1/17:</u></p> <ul style="list-style-type: none">• The maximum allowable morphine milligram equivalent (MME) is 200 MME. Prescriptions for short-acting (SA) and long-acting (LA) opioids are cumulatively included in this calculation. The prescription that exceeds the cumulative MME limit of 200 MME for a member will require prior authorization and may require a provider-to-provider telephone consultation with the pain management physician (free of charge and provided by Health First Colorado).• Prior authorization will be granted to allow for tapering• Prior authorization for 1 year will be granted for diagnosis of sickle cell anemia• Prior authorization for 1 year will be granted for admission to or diagnosis of hospice or end of life care• Prior authorization for 1 year will be granted for pain associated with cancer <p>MME calculation is conducted using conversion factors from the following link: https://pharmacypmp.az.gov/resources/mme-calculator</p> <p>Only one long-acting opioid agent (including different strengths) and one short-acting opioid agent (including different strengths) will be considered for a prior authorization.</p> <p>Medicaid provides guidance on the treatment of pain, including tapering, on our webpage under the heading Pain Management Resources and Opioid Use at: https://www.colorado.gov/pacific/hcpf/pain-management-resources-and-opioid-use</p> <p><u>Opioid Naïve Policy Effective 8/1/17 (<i>Update effective 04/01/23 in Italics</i>):</u></p> <p>Members who have not filled a prescription for an opioid within the past 180 days will be identified as “opioid treatment naïve” and have the following limitations placed on the initial prescription(s):</p> <ul style="list-style-type: none">• The prescription is limited to short-acting opioid agents <i>or Butrans (buprenorphine)</i>. Use of other long-acting opioid agents will require prior authorization approval for members identified as opioid treatment naïve.• The days’ supply of the first, second, and third prescription for an opioid will be limited to 7 days, the quantity will be limited to 8 dosage forms per day (tablets, capsules), maximum #56 tablets/capsules for a 7-day supply• The fourth prescription for an opioid will require prior authorization, filling further opioid prescriptions may require a clinical pharmacist review or provider to provider telephone consultation with a pain management physician (free of charge and provided by Health First Colorado).• If a member has had an opioid prescription filled within the past 180 days, then this policy would not apply to that member and other opioid policies would apply as applicable. <p><u>Dental Prescriptions Opioid Policy Effective 11/15/18 (implemented in the claims system 01/07/19):</u></p> <p>Members who receive an opioid prescribed by a dental provider will be subject to day supply limits and quantity per day limits for short acting opioids.</p> <ul style="list-style-type: none">• The prescription is limited to short-acting opioid agents only. Use of long-acting opioid agents and short acting fentanyl agents will require prior authorization approval for members’ prescriptions written by a dental provider.• The days’ supply of the first, second, and third prescription for an opioid will be limited to 4 days, the quantity will be limited to 6 dosage forms per day (tablets, capsules), maximum #24 tablets/capsules for a 4-day supply• The fourth prescription for an opioid will require prior authorization. A prior authorization for the fourth fill may be approved for up to a 7-day supply and the quantity will be limited to 8 dosage forms per day (#56 tablets/capsules) for members with any of the following diagnoses/undergoing any of the following procedures:<ul style="list-style-type: none">○ Traumatic oro-facial tissue injury with major mandibular/maxillary surgical procedures○ Severe cellulitis of facial planes○ Severely impacted teeth with facial space infection necessitating surgical management		

- Other potential exemptions that exceed the first 3 fill limits (day supply and quantity) may be evaluated with a provider-to-provider telephone consult with a pain management specialist (free of charge and provided by Health First Colorado)

If a member has had an opioid prescription prescribed by a non-dental provider, then this policy would not apply to that member and other opioid policies would apply as applicable. Dental prescriptions do not impact the opioid treatment naïve policy, but the prescriptions will be counted towards the Morphine Milligram Equivalent (MME) daily dose.

Opioid and Benzodiazepine Combination Effective 9/15/19:

Prior authorization will be required for members receiving long-term therapy with an opioid medication who are newly started on a benzodiazepine medication OR for members receiving long-term therapy with a benzodiazepine medication who are newly started on an opioid medication. Prior authorization may be approved if meeting the following:

- The member discontinued or is no longer taking either the opioid or benzodiazepine medication and will not be using these in combination **OR**
- The member will not be taking the prescribed opioid and benzodiazepine medications at the same time based on prescribed dosing interval (such as prn administration) for the regimen AND the prescriber attests that the member has received appropriate counseling* regarding the risks associated with combining opioid and benzodiazepine medications including increased risk for sedation, respiratory depression, overdose, and overdose-related death and counseling regarding the FDA Boxed Warning for combining these medications **OR**
- The prescriber has evaluated the regimen and attests that it is appropriate for the member to continue use of the concomitant opioid and benzodiazepine medication regimen as prescribed AND the prescriber attests that the member has received appropriate counseling* regarding the risks associated with combining opioid and benzodiazepine medications including increased risk for sedation, respiratory depression, overdose, and overdose-related death and counseling regarding the FDA Boxed Warning for combining these medications **OR**
- Prior authorization may be approved for members receiving palliative or hospice care **OR**
- For benzodiazepine prior authorizations, approval may be granted if the benzodiazepine is being prescribed for seizure disorder or convulsions.

**If counseling has not been provided, the prescriber attests that a reasonable effort will be made to contact the member or the member’s pharmacy to ensure that counseling is provided.*

Opioid and Quetiapine Combination Effective 9/15/19:

Pharmacy claims for members receiving opioid and quetiapine medications in combination will require entry of point-of-sale DUR service codes (Reason for Service, Professional Service, Result of Service) for override of drug-drug interaction (DD) related to risk of increased sedation from concomitant use of this drug combination.

Opioid and Buprenorphine-Containing substance use disorder (SUD) Product Combination Effective 06/01/21:

Opioid claims submitted for members currently receiving buprenorphine-containing SUD medications will require entry of point-of-sale DUR service codes (Reason for Service, Professional Service, Result of Service) for override of drug-drug interaction (DD) with use of this drug combination.

Therapeutic Drug Class: OPIOIDS, Short Acting – <i>Effective 4/1/2025</i>		
Preferred No PA Required* (If criteria and quantity limit are met)	Non-Preferred PA Required	<p>*Preferred codeine and tramadol products do not require prior authorization for adult members (18 years of age or greater) if meeting all other opioid policy criteria.</p> <p>Preferred codeine or tramadol products prescribed for members < 18 years of age must meet the following criteria:</p> <ul style="list-style-type: none">• Preferred tramadol and tramadol-containing products may be approved for members < 18 years of age if meeting the following:<ul style="list-style-type: none">○ Member is 12 years to 17 years of age AND○ Tramadol is NOT being prescribed for post-surgical pain following tonsil or adenoid procedure AND○ Member’s BMI-for-age is not > 95th percentile per CDC guidelines AND○ Member does not have obstructive sleep apnea or severe lung disease OR○ For members < 12 years of age with complex conditions or life-limiting illness who are receiving care under a pediatric specialist, tramadol and tramadol-containing products may be approved on a case-by-case basis• Preferred Codeine and codeine-containing products will receive prior authorization approval for members meeting the following criteria may be approved for members < 18 years of age if meeting the following:<ul style="list-style-type: none">○ Member is 12 years to 17 years of age AND
*Acetaminophen/codeine tablets	Acetaminophen / codeine elixir	
Hydrocodone/acetaminophen solution, tablet	ASCOMP WITH CODEINE	
Hydromorphone tablet	(codeine/butalbital/aspirin/caffeine)	
Morphine IR solution, tablet	*Butalbital/caffeine/acetaminophen/codeine capsule	
Oxycodone solution, tablet		
Oxycodone/acetaminophen tablet		

<div>*Tramadol 25mg, 50mg</div> <div>*Tramadol/acetaminophen tablet</div>	Butalbital/caffeine/aspirin/codeine capsule	<div><ul style="list-style-type: none">○ Codeine is NOT being prescribed for post-surgical pain following tonsil or adenoid procedure AND○ Member’s BMI-for-age is not > 95th percentile per CDC guidelines AND○ Member does not have obstructive sleep apnea or severe lung disease AND○ Member is not pregnant, or breastfeeding AND○ Renal function is not impaired (GFR > 50 ml/min) AND○ Member is not receiving strong inhibitors of CYP3A4 (such as erythromycin, clarithromycin, itraconazole, ketoconazole, posaconazole, fluconazole [≥200mg daily], voriconazole, delavirdine, and milk thistle) AND○ Member meets <u>one</u> of the following:<ul style="list-style-type: none">• Member has trialed codeine or codeine-containing products in the past with no history of allergy or adverse drug reaction to codeine• Member has not trialed codeine or codeine-containing products in the past and the prescriber acknowledges reading the following statement: “Approximately 1-2% of the population metabolizes codeine in a manner that exposes them to a much higher potential for toxicity. Another notable proportion of the population may not clinically respond to codeine. We ask that you please have close follow-up with members newly starting codeine and codeine-containing products to monitor for safety and efficacy.”</div> <div>Non-preferred tramadol products may be approved following trial and failure of generic tramadol 50mg tablet AND generic tramadol/acetaminophen tablet.</div> <div>All other non-preferred short-acting opioid products may be approved following trial and failure of three preferred products. Failure is defined as allergy‡, lack of efficacy, intolerable side effects, or significant drug-drug interaction.</div> <div>‡Allergy: hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, and angioedema</div> <div><u>Quantity Limits:</u> Short-acting opioids will be limited to a total of 120 tablets per 30 days (4/day) per member for members who are not included in the opioid treatment naive policy.<ul style="list-style-type: none">• Exceptions will be made for members with a diagnosis of a terminal illness (hospice or palliative care) or sickle cell anemia.• For members who are receiving more than 120 tablets currently and who do not have a qualifying exemption diagnosis, a 6-month prior authorization can be granted via the prior authorization process for providers to taper members.• Please note that if more than one agent is used, the combined total utilization may not exceed 120 units in 30 days. There may be allowed certain exceptions to this limit for acute situations (for example: post-operative surgery, fractures, shingles, car accident).</div> <div><u>Maximum Doses:</u> Tramadol: 400mg/day Codeine: 360mg/day Butorphanol intranasal: 10ml per 30 days (four 2.5ml 10mg/ml package units per 30 days)</div>
	Butalbital compound/codeine	
	Butorphanol tartrate (nasal) spray	
	Carisoprodol/aspirin/codeine	
	Codeine tablet	
	Dihydrocodeine/acetaminophen/caffeine tablet	
	DILAUDID (hydromorphone) solution, tablet	
	FIORICET/CODEINE (codeine/butalbital/acetaminophen/caffeine) capsule	
	Hydrocodone/ibuprofen tablet	
	Hydromorphone solution	
	Levorphanol tablet	
	Meperidine solution, tablet	
	Morphine concentrated solution, oral syringe	
	NALOCET (oxycodone/acetaminophen) tablet	

	<div>Pentazocine/naloxone tablet</div> <div>PERCOCET (oxycodone/ acetaminophen) tablet</div> <div>ROXICODONE (oxycodone) tablet</div> <div>ROXYBOND (oxycodone) tablet</div> <div>SEGLENTIS (tramadol/celecoxib) tablet</div> <div>Tramadol 100mg tablet</div> <div>Tramadol solution</div>	
Therapeutic Drug Class: FENTANYL PREPARATIONS (buccal, transmucosal, sublingual) – <i>Effective 4/1/2025</i>		
	<div>PA Required</div> <div>ACTIQ (fentanyl citrate) lozenge</div> <div>Fentanyl citrate lozenge, buccal tablet</div> <div>FENTORA (fentanyl citrate) buccal tablet</div>	<div>Fentanyl buccal, intranasal, transmucosal, and sublingual products:</div> <div>Prior authorization approval may be granted for members experiencing breakthrough cancer pain and those that have already received and are tolerant to opioid drugs for the cancer pain AND are currently being treated with a long-acting opioid drug. The prior authorization may be granted for up to 4 doses per day. For patients in hospice or palliative care, prior authorization will be automatically granted regardless of the number of doses prescribed.</div>
Therapeutic Drug Class: OPIOIDS, Long Acting – <i>Effective 4/1/2025</i>		
<div>Preferred No PA Required (unless indicated by * criteria)</div> <div>BELBUCA (buprenorphine) buccal film</div> <div>BUTRANS^{BNR} (buprenorphine) transdermal patch</div>	<div>Non-Preferred PA Required</div> <div>**OXYCONTIN (oxycodone ER) tablet</div> <div>Buprenorphine transdermal patch</div> <div>CONZIP (tramadol ER) capsule</div>	<div>*Belbuca (buprenorphine) buccal film may be approved for members who have trialed and failed‡ treatment with Butrans (buprenorphine) patch at a dose of 20 mcg/hr OR with prescriber confirmation that the maximum dose of Butrans 20 mcg/hr will not provide adequate analgesia. <u>Quantity limit</u>: 60 films/30 days.</div> <div>Oxycontin (oxycodone ER) may be approved for members who have trialed and failed‡ treatment with TWO preferred agents.</div>

<div>*Fentanyl 12mcg, 25mcg, 50mcg, 75mcg, 100mcg transdermal patch</div> <div>Morphine ER (generic MS Contin) tablet</div> <div>Tramadol ER (generic Ultram ER) tablet</div>	<div>Fentanyl 37mcg, 62mcg, 87mcg transdermal patch</div> <div>Hydrocodone ER capsule, tablet</div> <div>Hydromorphone ER tablet</div> <div>HYSINGLA (hydrocodone ER) tablet</div> <div>Methadone (all forms)</div> <div>Morphine ER capsule</div> <div>MS CONTIN (morphine ER) tablet</div> <div>Oxycodone ER tablet</div> <div>Oxymorphone ER tablet</div> <div>Tramadol ER capsule</div>	<div>All other non-preferred products may be approved for members who have trialed and failed‡ three preferred products.</div> <div>‡Failure is defined as lack of efficacy with 14-day trial, allergy (hives, maculopapular rash, erythema multiforme, pustular rash, intolerable application site skin reactions, severe hypotension, bronchospasm, and angioedema), intolerable side effects, or significant drug-drug interaction.</div> <div>Methadone: Members may receive 30-day approval when prescribed for neonatal abstinence syndrome without requiring trial and failure of preferred agents or opioid prescriber consultation.</div> <div>Methadone Continuation: Members who have been receiving methadone for pain indications do not have to meet non-preferred criteria. All new starts for methadone will require prior authorization under the non-preferred criteria listed above.</div> <div>If a prescriber would like to discuss strategies for tapering off methadone or transitioning to other pain management therapies for a Health First Colorado member, consultation with the Health First Colorado pain management physician is available free of charge by contacting the pharmacy call center helpdesk and requesting an opioid prescriber consult.</div> <div>Reauthorization: Reauthorization for a non-preferred agent may be approved if the following criteria are met:<ul style="list-style-type: none">Provider attests to continued benefit outweighing risk of opioid medication use ANDMember met original prior authorization criteria for this drug class at time of original authorization</div> <div>Quantity/Dosing Limits:<ul style="list-style-type: none">Oxycontin and Hydrocodone ER (generic Zohydro ER) will only be approved for twice daily dosing.Hysingla will only be approved for once daily dosing.Fentanyl patches will require a PA for doses of more than 15 patches/30 days (if using one strength) or 30 patches for 30 days (if using two strengths). For fentanyl patch strengths of 37mcg/hr, 62mcg/hr, and 87mcg/hr, member must trial and fail two preferred strengths of separate patches that will provide the desired dose (such as 12mcg/hr + 50mcg/hr = 62mcg/hr).</div>
Therapeutic Drug Class: BUPRENORPHINE, Injectable – <i>Effective 7/1/2025</i>		
<div>Preferred No PA Required (*Must meet eligibility criteria)</div> <div>Brixadi Weekly/Monthly (buprenorphine) syringe</div> <div>Sublocade (buprenorphine) syringe</div>	<div>Non-Preferred PA Required</div>	<div>Preferred agents may be approved if the following criteria are met:<ul style="list-style-type: none">The requested medication is being dispensed directly to the healthcare professional (medication should not be dispensed directly to the member) ANDProvider attests to member’s enrollment in a complete treatment program, including counseling and psychosocial support ANDMember has a documented diagnosis of moderate to severe opioid use disorder ANDFor members newly started on therapy who are not currently using a transmucosal buprenorphine-containing product, prescriber attests that transmucosal buprenorphine induction therapy will be initiated in accordance with product labeling.</div> <div>Maximum dose:<ul style="list-style-type: none">Brixadi (buprenorphine) injection: 128 mg/month</div>

		<ul style="list-style-type: none">Sublocade (buprenorphine) injection: 600 mg/month during 1st month of induction therapy; 300 mg/month maintenance dose thereafter																				
<h2>II. Anti-Infectives</h2>																						
Therapeutic Drug Class: ANTIBIOTICS, Inhaled – <i>Effective 1/1/2026</i>																						
<p>Preferred No PA Required (*Must meet eligibility criteria)</p> <p>Tobramycin inhalation solution (generic TOBI)</p> <p>*CAYSTON (aztreonam) inhalation solution</p>	<p>Non-Preferred PA Required</p> <p>ARIKAYCE (amikacin liposomal) inhalation vial</p> <p>BETHKIS (tobramycin) inhalation ampule</p> <p>KITABIS (tobramycin) nebulizer pak</p> <p>TOBI (tobramycin) inhalation solution</p> <p>TOBI PODHALER (tobramycin) inhalation capsule</p> <p>Tobramycin inhalation ampule (generic Bethkis)</p> <p>Tobramycin nebulizer pak (generic Kitabis)</p>	<p>*CAYSTON (aztreonam) inhalation solution may be approved if the following criteria are met:</p> <ul style="list-style-type: none">Member has a history of trial and failure of preferred tobramycin solution for inhalation (failure is defined as lack of efficacy with a 4-week trial, intolerable side effects, or significant drug-drug interactions) OR provider attests that member cannot use preferred tobramycin solution for inhalation due to documented allergy or contraindication to therapy ANDThe member has known colonization of <i>Pseudomonas aeruginosa</i> in the lungs ANDThe member has been prescribed an inhaled beta agonist to use prior to nebulization of Cayston (aztreonam). <p>ARIKAYCE (amikacin) may be approved if the following criteria are met:</p> <ul style="list-style-type: none">Member has refractory mycobacterium avium complex (MAC) lung disease with limited or no alternative treatment options available ANDMember has trialed and failed 6 months of therapy with a 3-drug regimen that includes a macrolide (failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions). <p>All other non-preferred inhaled antibiotic agents may be approved if the following criteria are met:</p> <ul style="list-style-type: none">The member has a diagnosis of cystic fibrosis with known colonization of <i>Pseudomonas aeruginosa</i> in the lungs ANDMember has history of trial and failure of preferred tobramycin solution for inhalation (failure is defined as lack of efficacy with a 4-week trial, contraindication to therapy, allergy, intolerable side effects or significant drug-drug interactions). <table><tr><th colspan="4">Table 1: Minimum Age, Maximum Dose, and Quantity Limitations</th></tr><tr><th>Drug Name</th><th>Minimum Age</th><th>Maximum Dose</th><th>Quantity Limit (Based on day supply limitation for pack size dispensed)</th></tr><tr><td>ARIKAYCE (amikacin)</td><td>≥ 18 years</td><td>590 mg once daily</td><td>Not applicable</td></tr><tr><td>BETHKIS (tobramycin)</td><td>Age ≥ 6 years</td><td>300 mg twice daily</td><td>28-day supply per 56-day period</td></tr><tr><td>CAYSTON (aztreonam)</td><td>≥ 7 years</td><td>75 mg three times daily</td><td>28-day supply per 56-day period</td></tr></table>	Table 1: Minimum Age, Maximum Dose, and Quantity Limitations				Drug Name	Minimum Age	Maximum Dose	Quantity Limit (Based on day supply limitation for pack size dispensed)	ARIKAYCE (amikacin)	≥ 18 years	590 mg once daily	Not applicable	BETHKIS (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period	CAYSTON (aztreonam)	≥ 7 years	75 mg three times daily	28-day supply per 56-day period
Table 1: Minimum Age, Maximum Dose, and Quantity Limitations																						
Drug Name	Minimum Age	Maximum Dose	Quantity Limit (Based on day supply limitation for pack size dispensed)																			
ARIKAYCE (amikacin)	≥ 18 years	590 mg once daily	Not applicable																			
BETHKIS (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period																			
CAYSTON (aztreonam)	≥ 7 years	75 mg three times daily	28-day supply per 56-day period																			

		<table><tr><td>KITABIS PAK (tobramycin)</td><td>Age ≥ 6 years</td><td>300 mg twice daily</td><td>28-day supply per 56-day period</td></tr><tr><td>TOBI[†] (tobramycin)</td><td>Age ≥ 6 years</td><td>300 mg twice daily</td><td>28-day supply per 56-day period</td></tr><tr><td>TOBI PODHALER (tobramycin)</td><td>Age ≥ 6 years</td><td>112 mg twice daily</td><td>28-day supply per 56-day period</td></tr><tr><td colspan="4">[†] Limitations apply to brand product formulation only</td></tr></table>	KITABIS PAK (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period	TOBI [†] (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period	TOBI PODHALER (tobramycin)	Age ≥ 6 years	112 mg twice daily	28-day supply per 56-day period	[†] Limitations apply to brand product formulation only			
KITABIS PAK (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period															
TOBI [†] (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period															
TOBI PODHALER (tobramycin)	Age ≥ 6 years	112 mg twice daily	28-day supply per 56-day period															
[†] Limitations apply to brand product formulation only																		
		Members currently stabilized on any inhaled antibiotic agent in this class may receive approval to continue that agent.																
Therapeutic Drug Class: ANTI-HERPETIC AGENTS - Oral – <i>Effective 1/1/2026</i>																		
No PA Required Acyclovir tablet, capsule *Acyclovir suspension (<i>members under 18 years or cannot swallow a solid dosage form</i>) Famciclovir tablet Valacyclovir tablet	PA Required Acyclovir suspension (<i>all other members</i>) VALTREX (valacyclovir) tablet	<p>Non-preferred products may be approved for members who have failed an adequate trial with two preferred products with different active ingredients. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Sitavig (acyclovir) buccal tablet may be approved for diagnosis of recurrent herpes labialis (cold sores) if member meets non-preferred criteria listed above AND has failed trial with oral acyclovir suspension. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>*Acyclovir suspension does not require prior authorization for members < 18 years of age and may be approved for members ≥ 18 years of age who cannot swallow an oral dosage form.</p> <table><tr><th colspan="3">Maximum Dose Table</th></tr><tr><th></th><th>Adult</th><th>Pediatric</th></tr><tr><td>Acyclovir</td><td>4,000 mg/day</td><td>3,200 mg/day</td></tr><tr><td>Famciclovir</td><td>2,000 mg/day</td><td></td></tr><tr><td>Valacyclovir</td><td>4,000 mg/day</td><td>Age 2-11 years: 3,000 mg/day Age ≥ 12 years: 4,000 mg/day</td></tr></table>		Maximum Dose Table				Adult	Pediatric	Acyclovir	4,000 mg/day	3,200 mg/day	Famciclovir	2,000 mg/day		Valacyclovir	4,000 mg/day	Age 2-11 years: 3,000 mg/day Age ≥ 12 years: 4,000 mg/day
Maximum Dose Table																		
	Adult	Pediatric																
Acyclovir	4,000 mg/day	3,200 mg/day																
Famciclovir	2,000 mg/day																	
Valacyclovir	4,000 mg/day	Age 2-11 years: 3,000 mg/day Age ≥ 12 years: 4,000 mg/day																
Therapeutic Drug Class: ANTI-HERPETIC AGENTS - Topical – <i>Effective 1/1/2026</i>																		
No PA Required Acyclovir cream (<i>Teva only</i>) Acyclovir ointment DENA VIR ^{BNR} (penciclovir) cream	PA Required Acyclovir cream (<i>all other manufacturers</i>) Penciclovir cream XERESE (acyclovir/ hydrocortisone) cream ZOVIRAX (acyclovir) cream, ointment	<p>Non-Preferred Zovirax and acyclovir ointment/cream formulations may be approved for members who have failed an adequate trial with the preferred topical acyclovir ointment/cream product (diagnosis, dose and duration) as deemed by approved compendium. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction)</p> <p>Xerese (acyclovir/hydrocortisone) prior authorization may be approved for members that meet the following criteria:</p> <ul style="list-style-type: none">• Documented diagnosis of recurrent herpes labialis AND• Member is immunocompetent AND• Member has failed treatment of at least 10 days with acyclovir (Failure is defined as significant drug-drug interaction, lack of efficacy, contraindication to or intolerable side effects) AND																

		<ul style="list-style-type: none">Member has failed treatment of at least one day with famciclovir 1500 mg OR valacyclovir 2 grams twice daily (Failure is defined as significant drug-drug interaction, lack of efficacy, contraindication to or intolerable side effects)
Therapeutic Drug Class: FLUOROQUINOLONES – Oral – <i>Effective 1/1/2026</i>		
<p>Preferred No PA Required (*if meeting eligibility criteria)</p> <p>*CIPRO (ciprofloxacin) oral suspension^{BNR}</p> <p>Ciprofloxacin tablet</p> <p>Levofloxacin tablet</p> <p>Moxifloxacin tablet</p>	<p>Non-Preferred PA Required</p> <p>BAXDELA (delafloxacin) tablet</p> <p>CIPRO (ciprofloxacin) tablet</p> <p>Ciprofloxacin oral suspension</p> <p>Levofloxacin oral solution</p> <p>Ofloxacin tablet</p>	<p>*CIPRO suspension does not require prior authorization for members < 18 years of age and may be approved for members ≥ 18 years of age</p> <p>Non-preferred products may be approved for members who have failed an adequate trial (7 days) with at least one preferred product. (Failure is defined as: lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>Levofloxacin solution may be approved for members with prescriber attestation that member:</p> <ul style="list-style-type: none">is unable to take Cipro (ciprofloxacin) crushed tablet or suspension ORis < 5 years of age and being treated for pneumonia ORis ≥ 6 months old and being treated for fever in the setting of chemotherapy-induced neutropenia ORhas failed† an adequate trial (7 days) of ciprofloxacin suspension <p>†Failure is defined as lack of efficacy, allergy, intolerable side effects, significant drug-drug interaction, or contraindication to therapy.</p>
Therapeutic Drug Class: HEPATITIS C VIRUS TREATMENTS – <i>Effective 1/1/2026</i>		
Direct Acting Antivirals (DAAs)		
<p>Preferred No PA Required for initial treatment (*must meet eligibility criteria)</p> <p>EPCLUSA (sofosbuvir/velpatasvir) 200 mg -50 mg, 150 mg-37.5 mg tablet, pellet pack</p> <p>HARVONI (ledipasvir/sofosbuvir) 45mg-200mg tablet, pellet pack</p> <p>Ledipasvir/Sofosbuvir 90 mg-400 mg tablet (<i>Asegua only</i>)</p> <p>MAVYRET (glecaprevir/pibrentasvir) tablet, pellet pack</p> <p>Sofosbuvir/Velpatasvir 400mg-100mg (<i>Asegua only</i>)</p> <p>*VOSEVI tablet (sofosbuvir/velpatasvir/voxilaprevir)</p>	<p>Non-Preferred PA Required</p> <p>EPCLUSA 400 mg-100 mg (sofosbuvir/velpatasvir) tablet</p> <p>HARVONI 90 mg-400 mg (ledipasvir/sofosbuvir) tablet</p> <p>SOVALDI (sofosbuvir) tablet, pellet packet</p> <p>ZEPATIER (elbasvir/grazoprevir) tablet</p>	<p>Pharmacy claims for preferred products prescribed for initial treatment will be eligible for up to a 90-day supply fill allowing for the appropriate days’ duration for completing the initial treatment regimen (with no PA required). Subsequent fills will require prior authorization meeting re-treatment criteria below.</p> <p>*Second line preferred agents (Vosevi) may be approved for members 18 years of age or older with chronic HCV infection who are non-cirrhotic or have compensated cirrhosis (Child-Pugh A) AND meet the following criteria:</p> <ul style="list-style-type: none">GT 1-6 and has previously failed treatment with a regimen containing an NS5A inhibitor (such as ledipasvir, daclatasvir, or ombitasvir) ORGT 1a or 3 and has previously failed treatment with a regimen containing sofosbuvir without an NS5A inhibitor ANDRequest meets the applicable criteria below for re-treatment. <p>Re-treatment: All requests for HCV re-treatment for members who have failed therapy with a DAA will be reviewed on a case-by-case basis. Additional information may be requested for re-treatment requests including:</p> <ul style="list-style-type: none">Assessment of member readiness for re-treatmentPrevious regimen medications and dates treatedGenotype of previous HCV infectionAny information regarding adherence to previously trialed regimen(s) and current chronic medicationsAdverse effects experienced from previous treatment regimenConcomitant therapies during previous treatment regimen

		<ul style="list-style-type: none">Vosevi regimens will require verification that member has been tested for evidence of active hepatitis B virus (HBV) infection and for evidence of prior HBV infection prior to initiating treatment. <p>Non-preferred agents may be approved if documentation is provided indicating an acceptable rationale for not prescribing a preferred treatment regimen (acceptable rationale may include patient-specific medical contraindications to a preferred treatment or cases where a member has initiated treatment on a non-preferred drug and needs to complete therapy).</p> <p>Members currently receiving treatment with a non-preferred agent will receive approval to finish their treatment regimen, provided required documentation is sent via normal prior authorization request process.</p>
Ribavirin Products		
No PA Required Ribavirin capsule Ribavirin tablet		Preferred products are eligible for up to a 90-day supply fill. Non-preferred ribavirin products require prior authorizations which will be evaluated on a case-by-case basis.
Therapeutic Drug Class: HUMAN IMMUNODEFICIENCY VIRUS (HIV) TREATMENTS, ORAL – <i>Effective 1/1/2026</i> Oral products indicated for HIV pre-exposure prophylaxis (PrEP) or post-exposure prophylaxis (PEP) are eligible for coverage with a written prescription by an enrolled pharmacist. Additional information regarding pharmacist enrollment can be found at https://hcpf.colorado.gov/pharm-serv.		
Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs)		
No PA Required EDURANT (rilpivirine) tablet Efavirenz capsule, tablet Etravirine tablet INTELENCE (etravirine) tablet Nevirapine suspension, IR tablet, ER tablet PIFELTRO (doravirine) tablet		All products are preferred and do not require prior authorization.
Nucleoside/Nucleotide Reverse Transcriptase Inhibitors (NRTIs)		
No PA Required Abacavir solution, tablet		All products are preferred and do not require prior authorization.

Didanosine DR capsule		
Emtricitabine capsule		
EMTRIVA (emtricitabine) capsule, solution		
EPIVIR (lamivudine) solution, tablet		
Lamivudine solution, tablet		
RETROVIR (zidovudine) capsule, syrup		
Stavudine capsule		
Tenofovir disoproxil fumarate (TDF) tablet		
VIREAD (TDF) oral powder, tablet		
ZIAGEN (abacavir) solution, tablet		
Zidovudine capsule, syrup, tablet		
Protease Inhibitors (PIs)		
No PA Required		All products are preferred and do not require prior authorization.
APTIVUS (tipranavir) capsule		
Atazanavir capsule		
Darunavir tablet		
Fosamprenavir tablet		
LEXIVA (fosamprenavir) suspension, tablet		
NORVIR (ritonavir) powder packet, tablet		
PREZISTA (darunavir) suspension, tablet		
REYATAZ (atazanavir) capsule, powder pack		
Ritonavir tablet		
VIRACEPT (nelfinavir) tablet		
Other Agents		

<div>No PA Required</div> <div>ISENTRESS (raltegravir) chewable, powder pack, tablet</div> <div>ISENTRESS HD (raltegravir) tablet</div> <div>Maraviroc tablet</div> <div>RUKOBIA (fostemsavir tromethamine ER) tablet</div> <div>SELZENTRY (maraviroc) solution, tablet</div> <div>SUNLENCA (lenacapavir) tablet</div> <div>TIVICAY (dolutegravir) tablet</div> <div>TIVICAY PD (dolutegravir) tablet for suspension</div> <div>TYBOST (cobicistat) tablet</div> <div>VOCABRIA (cabotegravir) tablet</div> <div>YEZTUGO (lenacapavir) tablet</div>		<div>All products are preferred and do not require prior authorization.</div>
Combination Agents		
<div>No PA Required</div> <div>Abacavir/Lamivudine tablet</div> <div>ATRIPLA (efavirenz/Emtricitabine/TDF) tablet</div> <div>BIKTARVY (bictegravir/emtricitabine/TAF) tablet</div> <div>CIMDUO (lamivudine/TDF) tablet</div> <div>COMBIVIR (lamivudine/zidovudine) tablet</div> <div>COMPLERA (emtricitabine/rilpivirine/TDF) tablet</div> <div>DELSTRIGO (doravirine/lamivudine/TDF) tablet</div> <div>DESCOVY (emtricitabine/TAF) tablet</div> <div>DOVATO (dolutegravir/lamivudine) tablet</div> <div>Efavirenz/Emtricitabine/TDF tablet</div> <div>Efavirenz/Lamivudine/TDF tablet</div>		<div>All products are preferred and do not require prior authorization.</div>

Emtricitabine/rilpivirine/TDF tablet		
Emtricitabine/TDF tablet		
EPZICOM (abacavir/lamivudine) tablet		
EVOTAZ (atazanavir/cobicistat) tablet		
GENVOYA (elvitegravir/cobicistat/ emtricitabine/TAF) tablet		
JULUCA (dolutegravir/rilpivirine) tablet		
KALETRA (lopinavir/ritonavir) solution, tablet		
Lamivudine/Zidovudine tablet		
Lopinavir/Ritonavir solution, tablet		
ODEFSEY (emtricitabine/rilpivirine/TAF) tablet		
PREZCOBIX (darunavir/cobicistat) tablet		
STRIBILD (elvitegravir/cobicistat/ emtricitabine/TDF) tablet		
SYMFI/SYMFI LO (efavirenz/lamivudine/TDF) tablet		
SYMTUZA (darunavir/cobicistat/ emtricitabine/TAF) tablet		
TRIUMEQ (abacavir/dolutegravir/ lamivudine) tablet		
TRIUMEQ PD (abacavir/dolutegravir) tablet for suspension		
TRIZIVIR (abacavir/lamivudine/zidovudine) tablet		
TRUVADA (emtricitabine/TDF) tablet		
Therapeutic Drug Class: TETRACYCLINES – <i>Effective 7/1/2025</i>		
No PA Required	PA Required	Prior authorization for non-preferred tetracycline agents may be approved if member has trialed/failed a preferred doxycycline product AND preferred minocycline. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. Prior authorization for liquid oral tetracycline formulations may be approved if member is unable to take a solid oral dosage form.
Doxycycline hyclate capsules	Demeclocycline tablet	
Doxycycline hyclate tablets	DORYX (doxycycline DR) tablet	
Doxycycline monohydrate 50mg, 100mg capsule	Doxycycline hyclate DR tablet	
Doxycycline monohydrate tablets		

Minocycline capsules	Doxycycline monohydrate 75mg, 150mg capsule Doxycycline monohydrate suspension Minocycline IR, ER tablet MINOLIRA (minocycline ER) tablet MORGIDOX (doxycycline/skin cleanser) kit NUZYRA (omadacycline) tablet SOLODYN ER (minocycline ER) tablet Tetracycline capsule XIMINO (minocycline ER) capsule	Nuzyra (omadacycline) prior authorization may be approved if member meets all of the following criteria: the above “non-preferred” prior authorization criteria and the following: <ul style="list-style-type: none">Member has trialed and failed[†] therapy with a preferred doxycycline product and preferred minocycline OR clinical rationale is provided describing why these medications cannot be trialed (including resistance and sensitivity) ANDMember has diagnosis of either Community Acquired Bacterial Pneumonia (CABP) or Acute Bacterial Skin and Skin Structure Infection (ABSSSI) or clinical rationale and supporting literature describing/supporting intended use AND one of the following:<ul style="list-style-type: none">If member diagnosis is ABSSSI, member must have trial and failure[†] of sulfamethoxazole/trimethoprim product in addition to preferred tetracyclines ORIf member diagnosis is CABP, member must have trial and failure[†] of either a beta-lactam antibiotic (amoxicillin/clavulanic acid) or a macrolide (azithromycin) AND <ul style="list-style-type: none">Maximum duration of use is 14 days <p>[†]Failure is defined as lack of efficacy with 7-day trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.</p>
----------------------	---	---

III. Cardiovascular

Therapeutic Drug Class: ALPHA-BLOCKERS – Effective 7/1/2025		
No PA Required	PA Required	Non-preferred products may be approved following trial and failure of one preferred product (failure is defined as lack of efficacy with 4-week trial, allergy or intolerable side effects).
Prazosin capsule	MINIPRESS (prazosin) capsule	
Therapeutic Drug Class: BETA-BLOCKERS – Effective 7/1/2025		
Beta-Blockers, Single Agent		
No PA Required (*Must meet eligibility criteria)	PA Required	*HEMANGEOL (propranolol) oral solution may be approved for members between 5 weeks and 1 year of age with proliferating infantile hemangioma requiring systemic therapy. Maximum dose: 1.7 mg/kg twice daily
Acebutolol capsule	Betaxolol tablet	Non-preferred products may be approved following trial and failure with two preferred products (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Atenolol tablet	BYSTOLIC (nebivolol) tablet	
Bisoprolol tablet	COREG (carvedilol) tablet	
Carvedilol IR tablet	COREG CR (carvedilol ER) capsule	
	Carvedilol ER capsule	INNOPRAN XL (propranolol ER) capsule brand product formulation may be approved if meeting the following: <ul style="list-style-type: none">Request meets non-preferred criteria listed above AND

*HEMANGEOL (propranolol) solution	INDERAL LA/XL (propranolol ER) capsule	<ul style="list-style-type: none">Member has trialed and failed therapy with a generic propranolol ER capsule formulation OR prescriber provides clinical rationale supporting why generic propranolol ER capsule product formulations cannot be trialed. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions. <p>KAPSPARGO SPRINKLE (metoprolol succinate) extended-release capsule may be approved for members ≥ 6 years of age who are unable to take a solid oral dosage form. Maximum dose: 200mg/day (adult); 50mg/day (pediatric)</p> <p>Members currently stabilized on timolol oral tablet non-preferred products may receive approval to continue on that product.</p> <p>Members currently stabilized on the non-preferred Bystolic (nebivolol) tablets may receive approval to continue on that product.</p> <p>Members currently stabilized on the non-preferred carvedilol ER capsules may receive approval to continue on that product.</p> <table><tr><th colspan="5">Table 1: Receptor Selectivity and Other Properties of Preferred Beta Blockers</th></tr><tr><th></th><th>β₁</th><th>β₂</th><th>Alpha-1 receptor antagonist</th><th>Intrinsic sympathomimetic activity (ISA)</th></tr><tr><td>Acebutolol</td><td>X</td><td></td><td></td><td>X</td></tr><tr><td>Atenolol</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Betaxolol</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Bisoprolol</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Carvedilol</td><td>X</td><td>X</td><td>X</td><td></td></tr><tr><td>Labetalol</td><td>X</td><td>X</td><td>X</td><td></td></tr><tr><td>Metoprolol succinate</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Metoprolol tartrate</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Nadolol</td><td>X</td><td>X</td><td></td><td></td></tr><tr><td>Nebivolol</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Pindolol</td><td>X</td><td>X</td><td></td><td>X</td></tr><tr><td>Propranolol</td><td>X</td><td>X</td><td></td><td></td></tr></table>	Table 1: Receptor Selectivity and Other Properties of Preferred Beta Blockers						β ₁	β ₂	Alpha-1 receptor antagonist	Intrinsic sympathomimetic activity (ISA)	Acebutolol	X			X	Atenolol	X				Betaxolol	X				Bisoprolol	X				Carvedilol	X	X	X		Labetalol	X	X	X		Metoprolol succinate	X				Metoprolol tartrate	X				Nadolol	X	X			Nebivolol	X				Pindolol	X	X		X	Propranolol	X	X		
Table 1: Receptor Selectivity and Other Properties of Preferred Beta Blockers																																																																								
	β ₁		β ₂	Alpha-1 receptor antagonist	Intrinsic sympathomimetic activity (ISA)																																																																			
Acebutolol	X				X																																																																			
Atenolol	X																																																																							
Betaxolol	X																																																																							
Bisoprolol	X																																																																							
Carvedilol	X		X	X																																																																				
Labetalol	X	X	X																																																																					
Metoprolol succinate	X																																																																							
Metoprolol tartrate	X																																																																							
Nadolol	X	X																																																																						
Nebivolol	X																																																																							
Pindolol	X	X		X																																																																				
Propranolol	X	X																																																																						
Labetalol tablet	INNOPRAN XL (propranolol ER) capsule																																																																							
Metoprolol tartrate tablet	KASPARGO (metoprolol succinate) sprinkle capsule																																																																							
Metoprolol succinate ER tablet	LOPRESSOR (metoprolol tartrate) tablet																																																																							
Nadolol tablet	Pindolol tablet																																																																							
Nebivolol tablet	TENORMIN (atenolol) tablet																																																																							
Propranolol IR tablet, solution	Timolol tablet																																																																							
Propranolol ER capsule	TOPROL XL (metoprolol succinate) tablet																																																																							

Beta-Blockers, Anti-Arrhythmics		
No PA Required Sotalol tablet	PA Required BETAPACE/AF (sotalol) tablet SOTYLIZE (sotalol) solution	SOTYLIZE (sotalol) oral solution may be approved for members 3 days to < 5 years of age. For members ≥ 5 years of age, SOTYLIZE (sotalol) oral solution may be approved for members who are unable to take a solid oral dosage form OR members that have trialed and failed therapy with one preferred product. (Failure is defined as allergy or intolerable side effects.) Maximum dose: 320 mg/day
Beta-Blockers, Combinations		
No PA Required Atenolol/Chlorthalidone tablet Bisoprolol/HCTZ tablet Metoprolol/HCTZ tablet	PA Required TENORETIC (atenolol/chlorthalidone) tablet ZIAC (bisoprolol/HCTZ) tablet	Non-preferred products may be approved following trial and failure with two preferred products (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Therapeutic Drug Class: CALCIUM CHANNEL-BLOCKERS – <i>Effective 7/1/2025</i>		
Dihydropyridines (DHPs)		
No PA Required Amlodipine tablet Felodipine ER tablet Nifedipine ER tablet Nifedipine IR capsule	PA Required ADALAT CC (nifedipine ER) tablet NORLIQVA (amlodipine) suspension KATERZIA (amlodipine) suspension Isradipine capsule Levamlodipine tablet Nicardipine capsule Nimodipine capsule Nisoldipine ER tablet NORVASC (amlodipine) tablet	Non-preferred products may be approved following trial and failure of two preferred agents. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions. Nimodipine oral capsule may be approved for adult members (≥ 18 years of age) with subarachnoid hemorrhage NYMALIZE (nimodipine) oral syringe may be approved for adult members (≥ 18 years of age) with subarachnoid hemorrhage who also have a feeding tube or have difficulty swallowing solid dosage forms. Maximum dose: 360 mg/day for 21 days (6 syringes/day or 126 syringes/21 days) KATERZIA (amlodipine) suspension may be approved if meeting the following: <ul style="list-style-type: none">• The member has a feeding tube or confirmed difficulty swallowing solid oral dosage forms OR cannot obtain the required dose through crushed amlodipine tablets AND• For members < 6 years of age, the prescriber confirms that the member has already been receiving the medication following initiation in a hospital or other clinical setting

	NYMALIZE (nimodipine) solution, oral syringe PROCARDIA XL (nifedipine ER) tablet SULAR (nisoldipine ER) tablet	
Non-Dihydropyridines (Non-DHPs)		
No PA Required Diltiazem IR tablet Diltiazem CD/ER capsule Verapamil IR, ER tablet Verapamil ER 120 mg, 180 mg, 240 mg capsule	PA Required CARDIZEM (diltiazem) tablet CARDIZEM CD/LA (diltiazem CD/ER) capsule, tablet Diltiazem ER/LA tablet TIAZAC ER (diltiazem ER) capsule Verapamil ER 360 mg capsule Verapamil PM ER 100 mg, 200 mg, 300 mg capsule VERELAN/PM (verapamil ER) pellet capsule	Non-preferred products may be approved following trial and failure of three preferred agents. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions.
Therapeutic Drug Class: ANGIOTENSIN MODIFIERS – <i>Effective 7/1/2025</i>		
Angiotensin-converting enzyme inhibitors (ACE Inh)		
No PA Required Benazepril tablet Enalapril tablet Fosinopril tablet Lisinopril tablet Quinapril tablet Ramipril tablet	PA Required ACCUPRIL (quinapril) tablet ALTACE (ramipril) capsule Captopril tablet Enalapril solution EPANED (enalapril) solution LOTENSIN (benazepril) tablet Moexipril tablet Perindopril tablet	Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products may be approved for members who have trialed and failed treatment with three preferred products (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction). Enalapril solution may be approved without trial and failure of three preferred agents for members who are unable to take a solid oral dosage form. QBRELIS (lisinopril) solution may be approved for members 6 years of age or older who are unable to take a solid oral dosage form and have trialed and failed Epaned (enalapril) solution. Failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction.

	<div>PRINIVIL (lisinopril) tablet</div> <div>QBRELIS (lisinopril) solution</div> <div>Trandolapril tablet</div> <div>VASOTEC (enalapril) tablet</div> <div>ZESTRIL (lisinopril) tablet</div>	
ACE Inhibitor Combinations		
<div>No PA Required</div> <div>Amlodipine/Benazepril capsule</div> <div>Benazepril/HCTZ tablet</div> <div>Enalapril/HCTZ tablet</div> <div>Lisinopril/HCTZ tablet</div> <div>Quinapril/HCTZ tablet</div>	<div>PA Required</div> <div>ACCURETIC (quinapril/HCTZ) tablet</div> <div>Captopril/HCTZ tablet</div> <div>Fosinopril/HCTZ tablet</div> <div>LOTENSIN HCT (benazepril/HCTZ) tablet</div> <div>LOTREL (amlodipine/benazepril) capsule</div> <div>VASERETIC (enalapril/HCTZ) tablet</div> <div>ZESTORETIC (lisinopril/HCTZ) tablet</div>	<div>Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products may be approved for members who have trialed and failed treatment with three preferred products (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction).</div>
Angiotensin II receptor blockers (ARBs)		
<div>No PA Required</div> <div>Irbesartan tablet</div> <div>Losartan tablet</div> <div>Olmesartan tablet</div> <div>Telmisartan tablet</div> <div>Valsartan tablet</div>	<div>PA Required</div> <div>ARBLI (losartan) oral suspension</div> <div>ATACAND (candesartan) tablet</div> <div>AVAPRO (irbesartan) tablet</div> <div>BENICAR (olmesartan) tablet</div> <div>Candesartan tablet</div> <div>COZAAR (losartan) tablet</div>	<div>Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products may be approved for members who have trialed and failed treatment with three preferred products (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction).</div>

	DIOVAN (valsartan) tablet EDARBI (azilsartan) tablet Eprosartan tablet MICARDIS (telmisartan) tablet Valsartan solution	
ARB Combinations		
<p>Preferred No PA Required (Unless indicated*)</p> <p>*ENTRESTO (sacubitril/valsartan) tablet^{BNR}</p> <p>Irbesartan/HCTZ tablet</p> <p>Losartan/HCTZ tablet</p> <p>Olmesartan/Amlodipine tablet</p> <p>Olmesartan/HCTZ tablet</p> <p>Telmisartan/HCTZ tablet</p> <p>Valsartan/Amlodipine tablet</p> <p>Valsartan/HCTZ tablet</p>	<p>Non-Preferred PA Required</p> <p>ATACAND HCT (candesartan/HCTZ) tablet</p> <p>AVALIDE (irbesartan/HCTZ) tablet</p> <p>AZOR (olmesartan/amlodipine) tablet</p> <p>BENICAR HCT (olmesartan/HCTZ) tablet</p> <p>Candesartan/HCTZ tablet</p> <p>DIOVAN HCT (valsartan/HCTZ) tablet</p> <p>EDARBYCLOR (azilsartan/chlorthalidone) tablet</p> <p>ENTRESTO (sacubitril/valsartan) sprinkles</p> <p>EXFORGE (valsartan/amlodipine) tablet</p> <p>EXFORGE HCT (valsartan/amlodipine/HCTZ) tablet</p> <p>HYZAAR (losartan/HCTZ) tablet</p>	<p>Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products may be approved for members who have trialed and failed treatment with three preferred products (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>*ENTRESTO (sacubitril/valsartan) may be approved for members if the following criteria are met:</p> <ul style="list-style-type: none">• Member is 1 to 17 years of age and has a diagnosis of symptomatic heart failure with systemic left ventricular systolic dysfunction (LVSD) and/or has chronic heart failure with a below-normal left ventricular ejection fraction (LVEF) OR• Member is ≥ 18 years of age and has a diagnosis of chronic heart failure.• Diagnosis will be verified through automated verification (AutoPA) of the appropriate corresponding ICD-10 diagnosis codes related to the indicated use of the medication.

	MICARDIS HCT (telmisartan/HCTZ) tablet Olmesartan/amlodipine/HCTZ tablet Sacubitril/valsartan tablet Telmisartan/amlodipine tablet TRIBENZOR (olmesartan/amlodipine/HCTZ) tablet Valsartan/Amlodipine/HCTZ tablet	
Renin Inhibitors & Renin Inhibitor Combinations		
	PA Required Aliskiren tablet TEKTURNA (aliskiren) tablet TEKTURNA HCT (aliskiren/HCTZ) tablet	Non-preferred renin inhibitors and renin inhibitor combination products may be approved for members who have failed treatment with three preferred products from the angiotensin modifier class (failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction). Renin inhibitors and combinations will not be approved in patients with diabetes. Renin inhibitors are contraindicated when used in combination with an ACE inhibitor, ACE inhibitor combination, ARB, or ARB combination.
Therapeutic Drug Class: PULMONARY ARTERIAL HYPERTENSION THERAPIES – <i>Effective 7/1/2025</i>		
Phosphodiesterase Inhibitors		
Preferred *Must meet eligibility criteria *Sildenafil oral suspension, tablet *Tadalafil 20mg tablet	Non-Preferred PA Required ADCIRCA (tadalafil) tablet ALYQ (tadalafil) tablet LIQREV (sildenafil) suspension REVATIO (sildenafil) suspension, tablet	*Eligibility criteria for preferred products: Preferred sildenafil and tadalafil tablet formulations may be approved for a diagnosis of pulmonary hypertension or right-sided heart failure. Sildenafil suspension may be approved for a diagnosis of pulmonary hypertension for members < 5 years of age who cannot take a solid oral dosage form. Non-preferred oral tablet products may be approved if meeting the following: <ul style="list-style-type: none">Member has a diagnosis of pulmonary hypertension ANDMember has trialed and failed treatment with preferred sildenafil tablet AND preferred tadalafil tablet. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction. Members who have been previously stabilized on a non-preferred product may receive approval to continue on the medication. Non-preferred oral liquid products may be approved if meeting the following:

	TADLIQ suspension	<ul style="list-style-type: none">Member has a diagnosis of pulmonary hypertension ANDRequest meets one of the following:<ul style="list-style-type: none">Member has trialed and failed treatment with one preferred oral liquid formulation (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction) ORPrescriber verifies that the member is unable to take a solid oral dosage form and that there is clinical necessity for use of a regimen with a less frequent dosing interval.
Endothelin Receptor Antagonists		
<p>Preferred *Must meet eligibility criteria</p> <p>*Ambrisentan tablet</p> <p>*Bosentan 62.5mg, 125mg tablet</p>	<p>Non-Preferred PA Required *Must meet eligibility criteria</p> <p>Bosentan tablet for suspension</p> <p>LETAIRIS (ambrisentan) tablet</p> <p>OPSUMIT (macitentan) tablet</p> <p>OPSYNVI (macitentan/tadalafil) tablet</p> <p>TRACLEER (bosentan) 32mg tablet for suspension</p> <p>TRACLEER (bosentan) 62.5mg, 125mg tablet</p>	<p>*Eligibility Criteria for all agents in the class Approval may be granted for a diagnosis of pulmonary hypertension. Member and prescriber should be enrolled in applicable REMS program for prescribed medication.</p> <p>Non-preferred agents may be approved for members who have trialed and failed two preferred agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>TRACLEER (bosentan) tablet for suspension may be approved if meeting one of the following:</p> <ul style="list-style-type: none">The member cannot swallow a solid oral dosage form ORThe request meets eligibility criteria and non-preferred criteria listed above. <p>Members who have been previously stabilized on a non-preferred product may receive approval to continue the medication.</p>
Prostacyclin Analogues and Receptor Agonists		
<p>Preferred (*Must meet eligibility criteria)</p> <p>*FLOLAN (epoprostenol) vial</p> <p>*ORENITRAM (treprostinil ER) tablet, titration kit</p> <p>*REMODULIN (treprostinil) vial</p> <p>*VENTAVIS (iloprost) inhalation solution</p>	<p>Non-Preferred PA Required</p> <p>Epoprostenol vial</p> <p>Treprostinil vial</p> <p>TYVASO (treprostinil) inhaler, inhalation solution</p>	<p>*Eligibility Criteria for all agents in the class Approval will be granted for a diagnosis of pulmonary hypertension.</p> <p>Non-preferred products may be approved for members who have failed treatment with a Preferred Product. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, contraindication to IV therapy or significant drug-drug interaction).</p> <p>Members who have been previously stabilized on a non-preferred product may receive approval to continue on the medication.</p>

	UPTRAVI (selexipag) tablet, dose pack, vial VELETRI (epoprostenol) vial YUTREPIA (treprostinil) capsule for inhalation	
Guanylate Cyclase (sGC) Stimulator		
	Non-Preferred PA Required ADEMPAS (riociguat) tablet	ADEMPAS (riociguat) may be approved for members who meet the following criteria: <ul style="list-style-type: none">For members of childbearing potential:<ul style="list-style-type: none">Member is not pregnant and is able to receive monthly pregnancy tests while taking ADEMPAS and one month after stopping therapy ANDMember and their partners are utilizing one of the following contraceptive methods during treatment and for one month after stopping treatment (IUD, contraceptive implants, tubal sterilization, a hormone method with a barrier method, two barrier methods, vasectomy with a hormone method, or vasectomy with a barrier method) AND <ul style="list-style-type: none">Member has a CrCl \geq 15 mL/min and is not on dialysis ANDMember does not have severe liver impairment (Child Pugh C) ANDMember has a diagnosis of persistent/recurrent chronic thromboembolic pulmonary hypertension (CTEPH) (WHO Group 4) after surgical treatment or has inoperable CTEPH ORMember has a diagnosis of pulmonary hypertension and has failed treatment with a preferred product for pulmonary hypertension. (Failure is defined as a lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction).
Therapeutic Drug Class: LIPOTROPICS – <i>Effective 7/1/2025</i>		
Bile Acid Sequestrants		
No PA Required Colesevelam tablet Colestipol tablet Cholestyramine packet, light packet, powder	PA Required Colesevelam packet COLESTID (colestipol) tablet, granules Colestipol granules QUESTRAN (cholestyramine/sugar) packet, powder QUESTRAN LIGHT (cholestyramine/ aspartame) packet, powder WELCHOL (colesevelam) packet, tablet	Non-preferred bile acid sequestrants may be approved if the member has failed treatment with 2 preferred products in the last 12 months (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions). Non-preferred lipotropic agents with a preferred product with same strength, dosage form, and active ingredient may be approved with adequate trial and/or failure of the preferred product with the same ingredient (such as preferred ezetimibe and Zetia) and 2 additional agents. (Failure is defined as: lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).

Fibrates					
No PA Required Fenofibric acid DR (generic Trilipix) capsule Fenofibrate capsule, tablet (generic Lofibra/Tricor) Gemfibrozil tablet	PA Required ANTARA (fenofibrate) capsule Fenofibric acid tablet Fenofibrate capsule (generic Antara/Fenoglide/Lipofen) FENOGLIDE (fenofibrate) tablet LIPOFEN (fenofibrate) capsule LOPID (gemfibrozil) tablet TRICOR (fenofibrate nano) tablet TRILIPIX (fenofibric acid) capsule	Non-preferred fibrates may be approved if the member has failed treatment with generic gemfibrozil or generic fenofibrate and niacin ER in the last 12 months (failure is defined as lack of efficacy with 4-week trial of each drug, allergy, intolerable side effects or significant drug-drug interactions). Non-preferred lipotropic agents with a preferred product with same strength, dosage form, and active ingredient may be approved with adequate trial and/or failure of the preferred product with the same ingredient (such as preferred ezetimibe and Zetia) and 2 additional agents. (Failure is defined as: lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).			
Other Lipotropics					
No PA Required (*Must meet eligibility criteria) Ezetimibe tablet Niacin ER tablet *Omega-3 ethyl esters capsule (generic Lovaza)	PA Required Icosapent ethyl capsule LOVAZA (omega-3 ethyl esters) capsule NEXLETOL (bempedoic acid) tablet NEXLIZET (bempedoic acid/ezetimibe) tablet ZETIA (ezetimibe) tablet	Non-preferred lipotropic agents with a preferred product with same strength, dosage form, and active ingredient may be approved with adequate trial and/or failure of the preferred product with the same ingredient and 2 additional agents. (Failure is defined as: lack of efficacy with 4-week trial, contraindication, allergy, intolerable side effects or significant drug-drug interactions). *Omega-3 ethyl esters (generic Lovaza) may be approved for members who have a baseline triglyceride level ≥ 500 mg/dL Lovaza (brand name) may be approved if meeting the following: <ul style="list-style-type: none">• Member has a baseline triglyceride level ≥ 500 mg/dl AND• Member has failed an adequate trial of omega-3 Ethyl Esters AND an adequate trial of gemfibrozil or fenofibrate (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions) Nexletol (bempedoic acid) or Nexlizet (bempedoic acid/ezetimibe) may be approved if meeting the following criteria: <ul style="list-style-type: none">• Member is ≥ 18 years of age AND• Member is not pregnant AND• Member is not receiving concurrent simvastatin > 20 mg daily or pravastatin > 40 mg daily AND• Member has a diagnosis of either heterozygous familial hypercholesterolemia or established atherosclerotic cardiovascular disease (see definition below), AND <table><tr><td>Conditions Which Define Clinical Atherosclerotic Cardiovascular Disease</td></tr><tr><td>• Acute Coronary Syndrome</td></tr><tr><td>• History of Myocardial Infarction</td></tr></table>	Conditions Which Define Clinical Atherosclerotic Cardiovascular Disease	• Acute Coronary Syndrome	• History of Myocardial Infarction
Conditions Which Define Clinical Atherosclerotic Cardiovascular Disease					
• Acute Coronary Syndrome					
• History of Myocardial Infarction					

		<div><ul style="list-style-type: none">• Stable or Unstable Angina• Coronary or other Arterial Revascularization• Stroke• Transient Ischemic Attack• Peripheral Arterial Disease of Atherosclerotic Origin</div> <div><ul style="list-style-type: none">• Member is concurrently adherent (> 80% of the past 180 days) on a maximally tolerated dose of a high intensity statin therapy (atorvastatin ≥ 40 mg daily OR rosuvastatin ≥ 20 mg daily [as a single-entity or as a combination product]) AND ezetimibe (as a single-entity or as a combination product) concomitantly for ≥ 8 continuous weeks), AND• If intolerant to a statin due to side effects, member must have a one month documented trial with at least two other maximally dosed statins in addition to ezetimibe. For members with a past or current incidence of rhabdomyolysis, a one-month trial and failure of a statin is not required, AND• Member has a treated LDL > 70 mg/dL for a clinical history of ASCVD OR LDL > 100 mg/dL if familial hypercholesterolemia</div> <div><u>Initial Approval:</u> 1 year</div> <div><u>Reauthorization:</u> Reauthorization may be approved for 1 year with provider attestation of medication safety and efficacy during the initial treatment period</div>
Therapeutic Drug Class: STATINS – <i>Effective 7/1/2025</i>		
No PA Required <div>Atorvastatin tablet</div> <div>Lovastatin tablet</div> <div>Pravastatin tablet</div> <div>Rosuvastatin tablet</div> <div>Simvastatin tablet</div>	PA Required <div>ALTOPREV (lovastatin ER) tablet</div> <div>ATORVALIQ (atorvastatin) suspension</div> <div>CRESTOR (rosuvastatin) tablet</div> <div>EZALLOR (rosuvastatin) sprinkle capsule</div> <div>FLOLIPID (simvastatin) suspension</div> <div>Fluvastatin capsule, ER tablet</div> <div>LESCOL XL (fluvastatin ER) tablet</div> <div>LIPITOR (atorvastatin) tablet</div> <div>LIVALO (pitavastatin) tablet</div> <div>Pitavastatin tablet</div> <div>ZOCOR (simvastatin) tablet</div>	<div>Non-preferred products may be approved following trial and failure of treatment with two preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions). For members who are unable to take a solid oral dosage form, non-preferred liquid product formulations may be approved without requiring trial and failure of preferred products.</div> <div>Age Limitations: Altoprev (lovastatin ER) will not be approved for members < 10 years of age. Fluvastatin will not be approved for members < 10 years of age. Livalo (pitavastatin) will not be approved for members < 8 years of age.</div>

	ZYPITAMAG (pitavastatin) tablet	
Therapeutic Drug Class: STATIN COMBINATIONS – <i>Effective 7/1/2025</i>		
No PA Required Simvastatin/Ezetimibe tablet	PA Required Atorvastatin/Amlodipine tablet CADUET (atorvastatin/amlodipine) tablet VYTORIN (simvastatin/ezetimibe) tablet	Non-preferred Statin combinations may be approved following trial and failure of treatment with two preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions). <u>Age Limitations:</u> Vytorin and generic ezetimibe/simvastatin will not be approved for members < 18 years of age. Caduet and generic amlodipine/atorvastatin will not be approved for members < 10 years of age.
Therapeutic Drug Class: Movement Disorders – <i>Effective 7/1/2025</i>		
No PA Required (*Must meet eligibility criteria) *Austedo (deutetrabenazine) tablet *Austedo (deutetrabenazine) XR tablet, titration pack *Ingrezza (valbenazine) capsule, initiation pack * Tetrabenazine tablet	PA Required Xenazine (tetrabenazine) tablet	*Eligibility Criteria for all agents in the class <ul style="list-style-type: none">• Member is ≥18 years of age AND• Member has been diagnosed with tardive dyskinesia or chorea associated with Huntington’s disease AND• If the member has hepatic impairment, FDA labeling for use has been evaluated AND• <u>For chorea associated with Huntington’s disease:</u><ul style="list-style-type: none">○ Member has been evaluated for untreated or inadequately treated depression and member has been counseled regarding the risks of depression and suicidality associated with agents in this therapeutic class.AND• <u>For tardive dyskinesia:</u><ul style="list-style-type: none">○ If applicable, the need for ongoing treatment with 1st and 2nd generation antipsychotics, metoclopramide, or prochlorperazine has been evaluated AND○ A baseline Abnormal Involuntary Movement Scale (AIMS) has been performed. Xenazine (tetrabenazine) Maximum dose 50 mg/day (PA available for extensive metabolizers of CYP2D6) Ingrezza (valbenazine) Quantity limits: <ul style="list-style-type: none">• 40 mg: 1.767 capsules/day• 60 mg: 1 capsule/day• 80 mg: 1 capsule/day Austedo (deutetrabenazine) Maximum dose: 48 mg/day

		Non-preferred Movement Disorder Agents may be approved for members ≥18 years of age after trial and failure of two preferred products. Failure is defined as lack of efficacy, contraindication, allergy, intolerable side effects or significant drug-drug interaction.
<h2>IV. Central Nervous System</h2>		
Therapeutic Drug Class: ANTICONVULSANTS -Oral – <i>Effective 4/1/2025</i>		
No PA Required	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i>	Members currently stabilized (in outpatient or acute care settings) on any non-preferred medication in this class may receive prior authorization approval to continue on that medication. Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription. <u>Non-Preferred Products Newly Started for Treating Seizure Disorder or Convulsions:</u> Non-preferred medications newly started for members with a diagnosis of seizure disorder/convulsions may be approved if the following criteria are met: <ul style="list-style-type: none">• The requested medication is being prescribed by a practitioner who has sufficient education and experience to safely manage treatment AND• The request meets minimum age and maximum dose limits listed in Table 1 AND• For medications indicated for use as adjunctive therapy, the medication is being used in conjunction with another medication indicated for treatment of seizure disorder/convulsions AND• The request meets additional criteria listed for any of the following: APTiom (eslicarbazepine) <ul style="list-style-type: none">• Member has history of trial and failure‡ of any carbamazepine-containing product BRIVIACT (brivaracetam) <ul style="list-style-type: none">• Member has history of trial and failure‡ of any levetiracetam-containing product DIACOMIT (stiripentol) <ul style="list-style-type: none">• Member is concomitantly taking clobazam AND• Member has diagnosis of seizures associated with Dravet syndrome ELEPSIA XR (levetiracetam ER) tablet <ul style="list-style-type: none">• Member has history of trial and failure‡ of levetiracetam ER (KEPPRA XR) EPIDIOLEX (cannabidiol) <ul style="list-style-type: none">• Member has diagnosis of seizures associated with Lennox-Gastaut syndrome (LGS) or Dravet Syndrome OR• Member has a diagnosis of seizures associated with tuberous sclerosis complex (TSC). FINTEPLA (fenfluramine) <ul style="list-style-type: none">• Member has a diagnosis of seizures associated with Dravet syndrome or Lennox-Gastaut syndrome
Barbiturates		
Phenobarbital elixir, solution, tablet Primidone tablet	MYSOLINE (primidone) tablet	
Hydantoins		
DILANTIN (phenytoin) 30 mg capsules, Infatab, suspension PHENYTEK (phenytoin ER) capsule Phenytoin suspension, chewable, ER capsule	DILANTIN (phenytoin ER), 100 mg capsules	
Succinamides		
Ethosuximide capsule, solution	CELONTIN (methsuximide) Kapseal Methsuximide capsule ZARONTIN (ethosuximide) capsule, solution	
Benzodiazepines		
Clobazam oral syringe, tablet, suspension Clonazepam tablet, ODT	KLONOPIN (clonazepam) tablet ONFI (clobazam) suspension, tablet	

	SYMPAZAN (clobazam) SL film	<p>OXTELLAR XR (oxcarbazepine ER)</p> <ul style="list-style-type: none">Member is being treated for partial-onset seizures ANDMember has history of trial and failure‡ of any carbamazepine or oxcarbazepine-containing product <p>SPRITAM (levetiracetam) tablet for suspension</p> <ul style="list-style-type: none">Member has history of trial and failure‡ of levetiracetam solution <p>SYMPAZAN (clobazam) film</p> <ul style="list-style-type: none">Member has history of trial and failure‡ of clobazam tablet or solution ORProvider attests that member cannot take clobazam tablet or solution <p><u>Non-Preferred Products Newly Started for Non-Seizure Disorder Diagnoses:</u></p> <p>Non-preferred medications newly started for non-seizure disorder diagnoses may be approved if meeting the following criteria:</p> <ul style="list-style-type: none">Member has history of trial and failure‡ of two preferred agents ANDThe prescription meets minimum age and maximum dose limits listed in Table 1. <p>‡Failure is defined as lack of efficacy, allergy, intolerable side effects, significant drug-drug interaction, documented contraindication to therapy, or inability to take preferred formulation. Members identified as HLA-B*15:02 positive, carbamazepine and oxcarbazepine should be avoided per Clinical Pharmacogenetics Implementation Consortium Guideline. This may be considered a trial for prior authorization approvals of a non-preferred agent.</p>
Valproic Acid and Derivatives		
DEPAKOTE (divalproex DR) sprinkle capsule	DEPAKOTE (divalproex DR) tablet	
Divalproex sprinkle capsule, DR tablet, ER tablet	DEPAKOTE ER (divalproex ER) tablet	
Valproic acid capsule, solution		
Carbamazepine Derivatives		
Carbamazepine IR tablet, ER tablet, chewable, ER capsule, suspension	APTIOM (eslicarbazepine) tablet	
CARBATROL ER (carbamazepine) capsule	Eslicarbazepine tablet	
Oxcarbazepine tablet	EQUETRO (carbamazepine) capsule	
TEGRETOL (carbamazepine) suspension, tablet	Oxcarbazepine suspension	
TEGRETOL XR (carbamazepine ER) tablet	Oxcarbazepine ER (generic Oxtellar XR) tablet	
TRILEPTAL ^{BNR} (oxcarbazepine) suspension	OXTELLAR XR (oxcarbazepine) tablet	
	TRILEPTAL (oxcarbazepine) tablet	
Lamotrigines		
Lamotrigine IR tablet, ER tablet, chewable/dispersible tablet, ODT	LAMICTAL (lamotrigine) chewable/dispersible dose pack, tablet	

Table 1: Non-preferred Product Minimum Age and Maximum Dose		
	Minimum Age**	Maximum Dose**
Barbiturates		
primidone (MYSOLINE)		2,000 mg per day
Benzodiazepines		
clobazam (ONFI) suspension, tablet	2 years	40 mg per day
clobazam film (SYMPAZAN)	2 years	40 mg per day
clonazepam (KLONOPIN)		20 mg per day
Brivaracetam/Levetiracetam		
brivaracetam (BRIVIACT)	1 month	200 mg per day
levetiracetam (KEPPRA)	1 month	3,000 mg per day
levetiracetam (SPRITAM)	4 years	3,000 mg per day
levetiracetam ER (ELEPSIA XR)	12 years	3,000 mg per day
levetiracetam ER (KEPPRA XR)	12 years	3,000 mg per day
Carbamazepine Derivatives		
carbamazepine (EPITOL)		1,600 mg per day
carbamazepine ER (EQUETRO)		1,600 mg per day
eslicarbazepine (APTIOM)	4 years	1,600 mg per day
oxcarbazepine ER (OXTELLAR XR)	6 years	2,400 mg per day

	LAMICTAL (lamotrigine) ODT, ODT dose pack LAMICTAL XR (lamotrigine ER) tablet, dose pack Lamotrigine ER/IR/ODT dose packs	Hydantoins phenytoin ER (DILANTIN) 100mg capsules, suspension, Infatab Lamotrigines lamotrigine IR (LAMICTAL) lamotrigine (LAMICTAL ODT) lamotrigine ER (LAMICTAL XR)		1,000 mg loading dose 600 mg/day maintenance dose
Topiramates		Succinamides ethosuximide (ZARONTIN) methsuximide (CELONTIN) Valproic Acid and Derivatives divalproex ER (DEPAKOTE ER)		
Topiramate tablet, sprinkle capsule	EPRONTIA (topiramate) solution QUDEXY XR (topiramate) capsule TOPAMAX (topiramate) tablet, sprinkle capsule Topiramate ER capsule, solution TROKENDI XR (topiramate ER) capsule	Topiramates topiramate (TOPAMAX) topiramate ER (QUDEXY XR) topiramate ER (TROKENDI XR) Other cannabidiol (EPIDIOLEX) cenobamate (XCOPRI) felbamate tablet, suspension fenfluramine (FINTEPLA) lacosamide (VIMPAT) perampanel (FYCOMPA) rufinamide (BANZEL) tablet and suspension stiripentol (DIACOMIT)	3 years 10 years 2 years 2 years 6 years	1,500 mg/day Not listed 60 mg/kg/day 400 mg per day 400 mg per day 400 mg per day
Brivaracetam/Levetiracetam		tiagabine tiagabine (GABITRIL) vigabatrin vigabatrin (SABRIL) vigabatrin (VIGADRUNE) powder packet zonisamide (ZONEGRAN)	1 year 18 years 2 years 2 years 1 month 4 years 1 year 6 months (weighing \geq 7 kg)	25 mg/kg/day 400 mg per day 3,600 mg per day 26 mg per day 400 mg per day 12 mg per day 3,200 mg per day 3,000 mg per day
Levetiracetam IR tablet, ER tablet, solution	BRIVIACT (brivaracetam) solution, tablet ELEPSIA XR (levetiracetam ER) tablet KEPPRA (levetiracetam) tablet, solution KEPRA XR (levetiracetam ER) tablet Levetiracetam 250mg tablets for suspension SPRITAM (levetiracetam) tablet	**Limits based on data from FDA package insert. Approval for age/dosing that falls outside of the indicated range may be evaluated on a case-by-case basis.		
Other				
*Felbamate suspension				

FELBATOL (felbamate) suspension	BANZEL (rufinamide) suspension, tablet	
FELBATOL (felbamate) ^{BNR} tablet	DIACOMIT (stiripentol) capsule, powder packet	
Lacosamide solution, tablet	EPIDIOLEX (cannabidiol) solution	
Zonisamide capsule	Felbamate tablet	
	FINTEPLA (fenfluramine) solution	
	FYCOMPA (perampanel) suspension, tablet	
	GABITRIL (tiagabine) tablet	
	Lacosamide UD solution	
	MOTPOLY XR (lacosamide) capsule	
	Perampanel tablet	
	Rufinamide suspension, tablet	
	SABRIL (vigabatrin) powder packet, tablet	
	Tiagabine tablet	
	Vigabatrin tablet, powder packet	
	VIGAFYDE (vigabatrin) solution	
	VIMPAT (lacosamide) solution, kit, tablet	
	XCOPRI (cenobamate) tablet, pack	
	ZONISADE (zonisamide) suspension	
	ZTALMY (ganaxolone) suspension	

No PA Required	PA Required	
Bupropion IR, SR, XL tablet	<i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i>	Non-preferred products may be approved for members who have failed adequate trial with two preferred newer generation anti-depressant products. If two preferred newer generation anti-depressant products are not available for indication being treated, approval of prior authorization for non-preferred products will require adequate trial of all preferred products FDA approved for that indication (failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction).
Citalopram solution, tablet		Zurzuvae (zuranolone) may be approved if meeting the following criteria:
Desvenlafaxine succinate ER (generic Pristiq) tablet		<ul style="list-style-type: none">Member is ≥ 18 years of age ANDMember has a diagnosis of postpartum depression based on Diagnostic and Statistical Manual of Mental Disorders (DSM-5) criteria for a major depressive episode ANDMember is not currently pregnant ANDPrescriber attests that the member has been counseled and has been engaged in shared decision making with regard to:<ul style="list-style-type: none">The importance of effective contraception during zuranolone treatment, as zuranolone may cause fetal harm ANDZuranolone is present in low levels in human breast milk and there are limited data on its effects on a breastfed infant ANDConsideration for the favorable long-term safety data associated with use of SSRIs as first-line, recommended therapies for perinatal depressive disorders by the American College of Obstetricians and Gynecologists (ACOG) or SNRIs as reasonable ACOG-recommended alternatives
Duloxetine (generic Cymbalta) capsule		AND
Escitalopram tablet	APLENZIN (bupropion ER) tablet	<ul style="list-style-type: none">Prescriber attests that the member has been counseled to refrain from engaging in potentially hazardous activities requiring mental alertness, including driving, for ≥ 12 hours after each zuranolone dose ANDThe member has been counseled to take the medication with 400 to 1,000 calories of food containing 25% to 50% fat ANDPrescriber verifies that concomitant medications have been assessed for potential drug interactions (CNS depressants, CYP3A4 inhibitors, CYP3A4 inducers) and any needed dosage adjustments for zuranolone have been made in accordance with package labeling ANDBaseline renal and hepatic function have been assessed and prescriber verifies that dosing is appropriate in accordance with package labeling.
Fluoxetine capsule, solution, 60 mg tablet	AUVELITY ER	<u>Quantity Limit:</u>
Fluvoxamine tablet	(dextromethorphan/bupropion) tablet	<ul style="list-style-type: none">Zurzuvae 20 mg and 25 mg: 28 capsules/14 daysZurzuvae 30 mg: 14 capsules/14 days
Mirtazapine tablet, ODT	Bupropion XL (generic Forfivo XL) tablet	<u>Maximum dose:</u> 50 mg once daily
Paroxetine IR tablet	CELEXA (citalopram) tablet	<u>Duration of Approval:</u> Approval will allow 30 days to fill for one 14-day course of treatment per postpartum period
Sertraline solution, tablet	Citalopram hydrobromide capsule	
Trazodone tablet	CYMBALTA (duloxetine) capsule	
Venlafaxine IR tablet	Desvenlafaxine fumarate ER tablet	
Venlafaxine ER capsules	DRIZALMA (duloxetine) sprinkle capsule	
Vilazodone tablet	EFFEXOR XR (venlafaxine ER) capsule	
	Escitalopram solution	
	FETZIMA (levomilnacipran ER) capsule, titration pack	
	Fluoxetine IR tablet, DR capsule	
	Fluvoxamine ER capsule	
	FORFIVO XL (bupropion ER) tablet	
	LEXAPRO (escitalopram) tablet	
	Nefazodone tablet	
	Paroxetine CR/ER tablet, suspension	
	Paroxetine mesylate capsule	

	<p>PAXIL (paroxetine) tablet, suspension</p> <p>PAXIL CR (paroxetine ER) tablet</p> <p>PEXEVA (paroxetine mesylate) tablet</p> <p>PRISTIQ (desvenlafaxine succinate ER) tablet</p> <p>PROZAC (fluoxetine) Pulvule</p> <p>RALDESY (trazodone) solution</p> <p>REMERON (mirtazapine) Soltab (ODT), tablet</p> <p>Sertraline capsule</p> <p>TRINTELLIX (vortioxetine) tablet</p> <p>Venlafaxine ER tablet</p> <p>Venlafaxine besylate ER tablet</p> <p>VIIBRYD (vilazodone) tablet, dose pack</p> <p>WELLBUTRIN SR, XL (bupropion) tablet</p> <p>ZOLOFT (sertraline) tablet, oral concentrate</p> <p>ZURZUVAE (zuranolone) capsule</p>	<p>Citalopram doses higher than 40mg/day for ≤60 years of age and 20mg/day for >60 years of age will require prior authorization. Please see the FDA guidance at: https://www.fda.gov/drugs/drugsafety/ucm297391.htm for important safety information.</p> <p>Members currently stabilized on a non-preferred newer generation antidepressant may receive approval to continue on that agent for one year if medically necessary. Verification may be provided from the prescriber or the pharmacy.</p>
Therapeutic Drug Class: MONOAMINE OXIDASE INHIBITORS (MAOIs) – <i>Effective 4/1/2025</i>		
	<p>PA Required</p> <p>EMSAM (selegiline) patch</p> <p>MARPLAN (isocarboxazid) tablet</p> <p>NARDIL (phenelzine) tablet</p> <p>Phenelzine tablet</p> <p>Tranylcypromine tablet</p>	<p>Non-preferred products may be approved for members who have failed adequate trial (8 weeks) with two preferred anti-depressant products. If two preferred anti-depressant products are not available for indication being treated, approval of prior authorization for non-preferred products will require adequate trial of all preferred anti-depressant products FDA approved for that indication. (Failure is defined as: lack of efficacy after 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction)</p> <p>Members currently stabilized on a Non-preferred MAOi antidepressant may receive approval to continue that agent for one year if medically necessary. Verification may be provided from the prescriber or the pharmacy.</p>

Therapeutic Drug Class: TRICYCLIC ANTI-DEPRESSANTS (TCAs) – <i>Effective 4/1/2025</i>		
No PA Required Amitriptyline tablet Clomipramine capsule Desipramine tablet Doxepin 10mg, 25mg, 50mg, 75mg, 100mg, 150mg capsule, oral concentrate Imipramine HCl tablet Nortriptyline capsule	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i> Amoxapine tablet ANAFRANIL (clomipramine) capsule Imipramine pamoate capsule NORPRAMIN (desipramine) tablet Nortriptyline solution PAMELOR (nortriptyline) capsule Protriptyline tablet Trimipramine capsule	<p>Non-preferred products may be approved for members who have failed adequate trial (8 weeks) with three preferred tricyclic products. If three preferred products are not available for indication being treated, approval of prior authorization for non-preferred products will require adequate trial of all tricyclic preferred products FDA approved for that indication. (Failure is defined as: lack of efficacy after 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction)</p> <p>Members currently stabilized on a non-preferred tricyclic antidepressant may receive approval to continue on that agent for one year if medically necessary. Verification may be provided from the prescriber or the pharmacy.</p>
Therapeutic Drug Class: ANTI-PARKINSON’S AGENTS – <i>Effective 4/1/2025</i>		
Dopa decarboxylase inhibitors, dopamine precursors and combinations		
No PA Required Carbidopa/Levodopa IR, ER tablet Carbidopa/Levodopa/Entacapone tablet	PA Required Carbidopa IR Carbidopa/Levodopa ODT CREXONT ER (carbidopa/levodopa) capsule DHIVY (carbidopa/levodopa) tablet DUOPA (carbidopa/levodopa) suspension INBRIJA (levodopa) capsule for inhalation	<p>Non-preferred agents may be approved with adequate trial and failure of carbidopa-levodopa IR and ER formulations (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>Carbidopa or levodopa single agent products may be approved for members with diagnosis of Parkinson’s Disease as add-on therapy to carbidopa-levodopa.</p> <p>Non-preferred medications that <u>are not</u> prescribed for Parkinson’s Disease (or an indication related to Parkinson’s Disease) may receive approval for other FDA-labeled indications without meeting trial and failure step therapy criteria.</p> <p>Members with history of trial and failure of a non-preferred Parkinson’s Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred.</p>

	LODOSYN (carbidopa) tablet RYTARY ER (carbidopa/levodopa) capsule SINEMET (carbidopa/levodopa) IR tablet STALEVO (carbidopa/levodopa/entacapone) tablet	Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.
MAO-B inhibitors		
No PA Required Rasagiline tablet Selegiline capsule, tablet	PA Required AZILECT (rasagiline) tablet XADAGO (safinamide) tablet ZELAPAR (selegiline) ODT	Non-preferred agents may be approved with adequate trial and failure of selegiline capsule or tablet (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects or significant drug-drug interactions). Non-preferred medications that are not prescribed for Parkinson’s Disease (or an indication related to Parkinson’s Disease) may receive approval for other FDA-labeled indications without meeting trial and failure step therapy criteria. Members with history of trial and failure of a non-preferred Parkinson’s Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred. Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.
Dopamine Agonists		
No PA Required Pramipexole IR tablet Ropinirole IR tablet	PA Required APOKYN (apomorphine) SC cartridge Apomorphine SC cartridge Bromocriptine capsule, tablet MIRAPEX (pramipexole) ER tablet NEUPRO (rotigotine) patch PARLODEL (bromocriptine) capsule, tablet Pramipexole ER tablet Ropinirole ER tablet	Non-preferred agents may be approved with adequate trial and failure of ropinirole IR AND pramipexole IR (failure is defined as lack of efficacy with 4-week trial, documented contraindication to therapy, allergy, intolerable side effects or significant drug-drug interactions). APOKYN (apomorphine subcutaneous cartridge) may be approved if meeting the following: <ul style="list-style-type: none">• APOKYN (apomorphine) is being used as an adjunct to other medications for acute, intermittent treatment of hypomobility, “off” episodes ("end-of-dose wearing off" and unpredictable "on/off" episodes) in patients with advanced Parkinson’s disease AND• Due to the risk of profound hypotension and loss of consciousness, member is not concomitantly using a 5HT3 antagonist such as ondansetron, granisetron, dolasetron, palonosetron or alosetron. Maximum dose: 6mg (0.6mL) three times per day KYNMOBI (apomorphine sublingual film) may be approved if meeting the following: <ul style="list-style-type: none">• KYNMOBI (apomorphine) is being used for the acute, intermittent treatment of “off” episodes in patients with Parkinson's disease AND• Due to the risk of profound hypotension and loss of consciousness, member must not be concomitantly using a 5HT3 antagonist such as ondansetron, granisetron, dolasetron, palonosetron or alosetron. Maximum dose: 30mg five times per day

		<p>Non-preferred medications that <u>are not</u> prescribed for Parkinson’s Disease (or an indication related to Parkinson’s Disease) may receive approval for other FDA-labeled indications without meeting trial and failure step therapy criteria.</p> <p>Members with history of trial and failure of a non-preferred Parkinson’s Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred.</p> <p>Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.</p>
Other Parkinson’s agents		
<p>No PA Required</p> <p>Amantadine capsule, solution/syrup</p> <p>Benztropine tablet</p> <p>Trihexyphenidyl tablet, elixir</p>	<p>PA Required</p> <p>Amantadine tablet</p> <p>COMTAN (entacapone) tablet</p> <p>Entacapone tablet</p> <p>GOCOVRI ER (amantadine ER) capsule</p> <p>NOURIANZ (istradefylline) tablet</p> <p>ONGENTYS (opicapone) capsule</p> <p>OSMOLEX ER (amantadine) tablet</p> <p>TASMAR (tolcapone) tablet</p> <p>Tolcapone tablet</p>	<p>Non-preferred agents may be approved with adequate trial and failure of two preferred agents (failure is defined as lack of efficacy with 4-week trial, documented contraindication to therapy, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>Non-preferred medications that <u>are not</u> prescribed for Parkinson’s Disease (or an indication related to Parkinson’s Disease) may receive approval for other FDA-labeled indications without meeting trial and failure step therapy criteria.</p> <p>Members with history of trial and failure of a non-preferred Parkinson’s Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred.</p> <p>Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.</p>
Therapeutic Drug Class: BENZODIAZEPINES (NON-SEDATIVE HYPNOTIC) – <i>Effective 4/1/2025</i>		
<p>No PA Required (*may be subject to age limitations)</p> <p>Alprazolam IR, ER tablet*</p> <p>Chlordiazepoxide capsule*</p> <p>Clonazepam tablet, ODT</p> <p>Clorazepate tablet*</p> <p>Diazepam tablet*, solution</p> <p>Lorazepam tablet*, oral concentrate</p> <p>Oxazepam capsule*</p>	<p>PA Required</p> <p>Alprazolam ODT, oral concentrate</p> <p>ATIVAN (lorazepam) tablet</p> <p>Diazepam Intensol</p> <p>KLONOPIN (clonazepam) tablet</p> <p>LOREEV (lorazepam ER) capsule</p> <p>XANAX (alprazolam) tablet</p> <p>XANAX XR (alprazolam ER) tablet</p>	<p>Non-preferred products may be approved following trial and failure of three preferred agents. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions.</p> <p><u>Children:</u> Prior authorization will be required for all agents when prescribed for children <18 years of age (with the exception of oral solution products) and may be approved with prescriber verification of necessity of use for member age.</p> <p>Diazepam Intensol may be approved following trial and failure of the preferred 5 mg/5 mL oral solution. Failure is defined as intolerable side effects, drug-drug interaction, or lack of efficacy.</p> <p>All benzodiazepine anxiolytics will require prior authorization for members ≥ 65 years of age when exceeding 90 days of therapy.</p> <p>Continuation of Therapy:</p> <ul style="list-style-type: none">Members < 65 years of age who are currently stabilized on a non-preferred benzodiazepine medication may receive approval to continue that medication.

		<ul style="list-style-type: none">Members < 18 years of age who are currently stabilized on a non-preferred oral solution product may receive authorization to continue that medication. <p>Prior authorization will be required for prescribed doses that exceed the maximum (Table 1).</p> <table><tr><th colspan="3">Table 1 Maximum Doses</th></tr><tr><th>Product</th><th>Maximum Daily Dose</th><th>Maximum Monthly Dose</th></tr><tr><td>Alprazolam tablet</td><td rowspan="6"><u>Adults ≥ 18 years:</u> 10 mg/day</td><td rowspan="6">Total of 300 mg from all dosage forms per 30 days</td></tr><tr><td>Alprazolam ER tablet</td></tr><tr><td>Alprazolam ODT</td></tr><tr><td>XANAX (alprazolam) tablet</td></tr><tr><td>XANAX XR (alprazolam ER) tablet</td></tr><tr><td>Alprazolam Intensol oral concentrate 1 mg/mL</td></tr><tr><td>Clorazepate tablet</td><td rowspan="2"><u>>12 years:</u> 90 mg/day <u>Children 9-12 years:</u> up to 60 mg/day</td><td rowspan="2">Total of 2,700 mg (adults) and 1,800 mg (children) from all tablet strengths per 30 days</td></tr><tr><td>TRANXENE (clorazepate) T-Tab</td></tr><tr><td>Chlordiazepoxide capsule</td><td><u>Adults ≥ 18 years:</u> 300 mg/day <u>Children 6-17 years:</u> up to 40 mg/day (pre-operative apprehension and anxiety)</td><td>Total of 9,000 mg (adults) and 120 mg (children, pre-op therapy) from all tablet strengths per 30 days</td></tr><tr><td>Diazepam Intensol oral concentrate 5 mg/mL</td><td rowspan="3"><u>Adults ≥ 18 years:</u> 40 mg/day <u>Members age 6 months to 17 years:</u> up to 10 mg/day</td><td rowspan="3">Total of 1200 mg (adults) and 300 mg (pediatrics) from all dosage forms per 30 days</td></tr><tr><td>Diazepam solution 5 mg/5 mL</td></tr><tr><td>Diazepam tablet</td></tr><tr><td>ATIVAN (lorazepam) Intensol concentrate 2 mg/mL</td><td rowspan="4"><u>Adults ≥ 18 years:</u> 10 mg/day <u>Children:</u> N/A</td><td rowspan="4">Total of 300 mg from all dosage forms per 30 days</td></tr><tr><td>ATIVAN (lorazepam) tablet</td></tr><tr><td>Lorazepam oral concentrated soln 2 mg/mL</td></tr><tr><td>Lorazepam tablet</td></tr><tr><td>Oxazepam capsule</td><td><u>Adults ≥ 18 years:</u> 120 mg/day <u>Children 6-18 years:</u></td><td>Total of 3600 mg from all dosage forms per 30 days</td></tr></table>	Table 1 Maximum Doses			Product	Maximum Daily Dose	Maximum Monthly Dose	Alprazolam tablet	<u>Adults ≥ 18 years:</u> 10 mg/day	Total of 300 mg from all dosage forms per 30 days	Alprazolam ER tablet	Alprazolam ODT	XANAX (alprazolam) tablet	XANAX XR (alprazolam ER) tablet	Alprazolam Intensol oral concentrate 1 mg/mL	Clorazepate tablet	<u>>12 years:</u> 90 mg/day <u>Children 9-12 years:</u> up to 60 mg/day	Total of 2,700 mg (adults) and 1,800 mg (children) from all tablet strengths per 30 days	TRANXENE (clorazepate) T-Tab	Chlordiazepoxide capsule	<u>Adults ≥ 18 years:</u> 300 mg/day <u>Children 6-17 years:</u> up to 40 mg/day (pre-operative apprehension and anxiety)	Total of 9,000 mg (adults) and 120 mg (children, pre-op therapy) from all tablet strengths per 30 days	Diazepam Intensol oral concentrate 5 mg/mL	<u>Adults ≥ 18 years:</u> 40 mg/day <u>Members age 6 months to 17 years:</u> up to 10 mg/day	Total of 1200 mg (adults) and 300 mg (pediatrics) from all dosage forms per 30 days	Diazepam solution 5 mg/5 mL	Diazepam tablet	ATIVAN (lorazepam) Intensol concentrate 2 mg/mL	<u>Adults ≥ 18 years:</u> 10 mg/day <u>Children:</u> N/A	Total of 300 mg from all dosage forms per 30 days	ATIVAN (lorazepam) tablet	Lorazepam oral concentrated soln 2 mg/mL	Lorazepam tablet	Oxazepam capsule	<u>Adults ≥ 18 years:</u> 120 mg/day <u>Children 6-18 years:</u>	Total of 3600 mg from all dosage forms per 30 days
Table 1 Maximum Doses																																					
Product	Maximum Daily Dose	Maximum Monthly Dose																																			
Alprazolam tablet	<u>Adults ≥ 18 years:</u> 10 mg/day	Total of 300 mg from all dosage forms per 30 days																																			
Alprazolam ER tablet																																					
Alprazolam ODT																																					
XANAX (alprazolam) tablet																																					
XANAX XR (alprazolam ER) tablet																																					
Alprazolam Intensol oral concentrate 1 mg/mL																																					
Clorazepate tablet	<u>>12 years:</u> 90 mg/day <u>Children 9-12 years:</u> up to 60 mg/day	Total of 2,700 mg (adults) and 1,800 mg (children) from all tablet strengths per 30 days																																			
TRANXENE (clorazepate) T-Tab																																					
Chlordiazepoxide capsule	<u>Adults ≥ 18 years:</u> 300 mg/day <u>Children 6-17 years:</u> up to 40 mg/day (pre-operative apprehension and anxiety)	Total of 9,000 mg (adults) and 120 mg (children, pre-op therapy) from all tablet strengths per 30 days																																			
Diazepam Intensol oral concentrate 5 mg/mL	<u>Adults ≥ 18 years:</u> 40 mg/day <u>Members age 6 months to 17 years:</u> up to 10 mg/day	Total of 1200 mg (adults) and 300 mg (pediatrics) from all dosage forms per 30 days																																			
Diazepam solution 5 mg/5 mL																																					
Diazepam tablet																																					
ATIVAN (lorazepam) Intensol concentrate 2 mg/mL	<u>Adults ≥ 18 years:</u> 10 mg/day <u>Children:</u> N/A	Total of 300 mg from all dosage forms per 30 days																																			
ATIVAN (lorazepam) tablet																																					
Lorazepam oral concentrated soln 2 mg/mL																																					
Lorazepam tablet																																					
Oxazepam capsule	<u>Adults ≥ 18 years:</u> 120 mg/day <u>Children 6-18 years:</u>	Total of 3600 mg from all dosage forms per 30 days																																			

			absolute dosage not established	
Therapeutic Drug Class: ANXIOLYTIC, NON- BENZODIAZEPINES – <i>Effective 4/1/2025</i>				
No PA Required Buspirone tablet		Non-preferred products may be approved following trial and failure of buspirone. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions.		
Therapeutic Drug Class: ATYPICAL ANTI-PSYCHOTICS - Oral and Topical – <i>Effective 4/1/2025</i>				
No PA Required (unless indicated by * in criteria; all products subject to dose and age limitations)	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i>	*Vraylar (cariprazine) or Rexulti (brexpiprazole) may be approved for members after trial and failure of one preferred agent. Failure is defined as contraindication, lack of efficacy with 6-week trial, allergy, intolerable side effects, significant drug-drug interactions, or known interacting genetic polymorphism that prevents safe preferred product dosing.		
Aripiprazole tablet		Non-preferred products may be approved for members meeting all of the following:		
Asenapine SL tablet		<ul style="list-style-type: none">• Medication is being prescribed for an FDA-Approved indication AND		
Clozapine tablet		<ul style="list-style-type: none">• Prescription meets dose and age limitations (Table 1) AND		
Lurasidone tablet	ABILIFY (aripiprazole) tablet, MyCite	<ul style="list-style-type: none">• Request meets one of the following:<ul style="list-style-type: none">○ Member has history of trial and failure of two preferred products with FDA approval for use for the prescribed indication (failure defined as lack of efficacy with 6-week trial, allergy, intolerable side effects (including rapid weight gain), contraindication, significant drug-drug interactions, or known interacting genetic polymorphism that prevents safe preferred product dosing) OR		
Olanzapine tablet, ODT	Aripiprazole oral solution, ODT	<ul style="list-style-type: none">○ Prescriber attests that within the last year (365 days) the member has trialed and failed (been unsuccessfully treated with) a preferred antipsychotic medication that was used to treat the member’s diagnosis (failure defined as lack of efficacy with 6-week trial, allergy, intolerable side effects (including rapid weight gain), significant drug-drug interactions, or known interacting genetic polymorphism that prevents safe preferred product dosing). Treatment must be under an FDA approved indication for a mental health condition or disorder.		
Paliperidone ER tablet	CAPLYTA (lumateperone) capsule			
Quetiapine IR tablet**	COBENFY (xanomeline/trospium) capsule, starter pack			
Quetiapine ER tablet				
REXULTI (brexpiprazole) dose pack, tablet*	Clozapine ODT	Age Limits: All products including preferred products will require a PA for members younger than the FDA approved age for the agent (Table 1). Members younger than the FDA approved age for the agent who are currently stabilized on an atypical antipsychotic will be eligible for approval.		
Risperidone ODT, oral solution, tablet	CLOZARIL (clozapine) tablet, ODT	Atypical Antipsychotic prescriptions for members under 5 years of age may require a provider-provider telephone consult with a child and adolescent psychiatrist (provided at no cost to provider or member).		
VRAYLAR (cariprazine) capsule*	FANAPT (iloperidone tablet, titration pack)			
Ziprasidone capsule	GEODON (ziprasidone) capsule	**Quetiapine IR when given at subtherapeutic doses may be restricted for therapy. Low-dose quetiapine (<150mg/day) is only FDA approved as part of a drug titration schedule to aid patients in getting to the target quetiapine dose. PA will be required for quetiapine < 150mg per day except for utilization (when appropriate) in members 65 years or older. PA will be approved for members 10-17 years of age with approved diagnosis (Table 1) stabilized on <150mg quetiapine IR per day.		
	INVEGA ER (paliperidone) tablet			
	LATUDA (lurasidone) tablet			
	LYBALVI (olanzapine/samidorphan) tablet	Aripiprazole solution: Aripiprazole <u>tablet</u> quantity limit is 2 tablets/day for pediatric members to allow for incremental dose titration and use of the preferred tablet formulation should be considered for dose titrations when possible and clinically appropriate. If incremental dose cannot be achieved with titration of the aripiprazole tablet for members < 18 years of age OR for members unable to swallow solid tablet dosage form, aripiprazole solution may be approved. For all other cases, aripiprazole solution is subject to meeting non-preferred product approval criteria listed above.		

	<p>NUPLAZID (pimavanserin) capsule, tablet</p> <p>Olanzapine/Fluoxetine capsule</p> <p>OPIPZA (aripiprazole) film</p> <p>RISPERDAL (risperidone) tablet, oral solution</p> <p>SAPHRIS (asenapine) SL tablet</p> <p>SECUADO (asenapine) patch</p> <p>SEROQUEL IR (quetiapine IR) tablet***</p> <p>SEROQUEL XR (quetiapine ER) tablet</p> <p>SYMBYAX (olanzapine/fluoxetine) capsule</p> <p>VERSACLOZ (clozapine) suspension</p> <p>ZYPREXA (olanzapine) tablet</p> <p>ZYPREXA ZYDIS (olanzapine) ODT</p>	<p>Nuplazid (pimavanserin tartrate) may be approved for the treatment of hallucinations and delusions associated with Parkinson’s Disease psychosis AND following trial and failure of therapy with quetiapine or clozapine, or clinical rationale is provided supporting why these medications cannot be trialed. Failure will be defined as contraindication, intolerable side effects, drug-drug interaction, or lack of efficacy.</p> <p>Abilify MyCite may be approved if meeting all of the following:</p> <ul style="list-style-type: none">• Member has history of adequate trial and failure of 5 preferred agents (one trial must include aripiprazole tablet). Failure is defined as lack of efficacy with 6-week trial on maximally tolerated dose, allergy, intolerable side effects, significant drug-drug interactions AND• Information is provided regarding adherence measures being recommended by provider and followed by member (such as medication organizer or digital medication reminders) AND• Member has history of adequate trial and failure of 3 long-acting injectable formulations of atypical antipsychotics, one of which must contain aripiprazole (failure is defined as lack of efficacy with 8-week trial, contraindication, allergy, intolerable side effects, significant drug-drug interactions) AND• Abilify MyCite is being used with a MyCite patch and member is using a compatible mobile application. AND• Medication adherence information is being shared with their provider via a web portal or dashboard. <p><u>Quantity Limits:</u> Quantity limits will be applied to all products (Table 1). In order to receive approval for off-label dosing, the member must have an FDA approved indication and must have tried and failed on the FDA approved dosing regimen.</p> <p>Members currently stabilized on a non-preferred atypical antipsychotic may receive approval to continue therapy with that agent for one year.</p>
Therapeutic Drug Class: ATYPICAL ANTI-PSYCHOTICS – Injectables – <i>Effective 10/1/2025</i>		

No PA Required	PA Required																																											
ABILIFY ASIMTUFII (aripiprazole) syringe, vial	<i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i>	Preferred products do not require prior authorization. All products are subject to meeting FDA-labeled dosing quantity limits listed in Table 1.																																										
ABILIFY MAINTENA (aripiprazole) syringe, vial		Non-preferred products may be approved for members meeting the following: <ul style="list-style-type: none">Medication is being prescribed for an FDA-Approved indication ANDPrescription meets dose limitations (Table 1) ANDMember has history of trial and failure of one preferred product with FDA approval for use for the prescribed indication (failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, contraindication, significant drug-drug interactions, or known interacting genetic polymorphism that prevents safe preferred product dosing).																																										
ARISTADA ER (aripiprazole lauroxil) syringe																																												
ARISTADA INITIO (aripiprazole lauroxil) syringe																																												
Chlorpromazine ampule, vial																																												
Fluphenazine vial																																												
Fluphenazine decanoate vial																																												
HALDOL (haloperidol decanoate) ampule																																												
Haloperidol decanoate ampule, vial																																												
Haloperidol lactate syringe, vial																																												
INVEGA HAFYERA (paliperidone palmitate) syringe																																												
INVEGA SUSTENNA (paliperidone palmitate) syringe																																												
INVEGA TRINZA (paliperidone palmitate) syringe																																												
Olanzapine vial																																												
PERSERIS ER (risperidone) syringe, syringe kit																																												
RISPERDAL CONSTA ^{BNR} (risperidone microspheres) syringe, vial																																												
UZEDY (risperidone) syringe																																												
Ziprasidone																																												
ZYPREXA RELPREVV (olanzapine pamoate) Vial kit																																												
		<table><tr><th colspan="3">Table 1: FDA-Labeled Dosing Quantity Limits</th></tr><tr><th>Long-Acting injectable</th><th>Route</th><th>Quantity Limit</th></tr><tr><td>ABILIFY ASIMTUFII (aripiprazole)</td><td>IM</td><td>1 pack/2 months (56 days)</td></tr><tr><td>ABILIFY MAINTENA (aripiprazole)</td><td>IM</td><td>1 pack/28 days</td></tr><tr><td>ARISTADA ER (aripiprazole)</td><td>IM</td><td>1,064 mg: 1 pack/2 months (56 days) All other strengths: 1 pack/28 days</td></tr><tr><td>ARISTADA INITIO (aripiprazole)</td><td>IM</td><td>1 pack/7 weeks (49 days)</td></tr><tr><td>INVEGA HAFYERA (paliperidone)</td><td>IM</td><td>1 pack/6 months (168 days)</td></tr><tr><td>INVEGA SUSTENNA (paliperidone)</td><td>IM</td><td>156 mg: 2 packs/5 weeks (35 days) All other strengths: 1 pack/28 days</td></tr><tr><td>INVEGA TRINZA (paliperidone)</td><td>IM</td><td>1 pack/3 months (84 days)</td></tr><tr><td>PERSERIS ER (risperidone)</td><td>Subcutaneous</td><td>1 pack/28 days</td></tr><tr><td>RISPERDAL CONSTA (risperidone)</td><td>IM</td><td>2 packs/28 days</td></tr><tr><td>UZEDY ER (risperidone)</td><td>Subcutaneous</td><td>150 mg, 200 mg and 250 mg: 1 pack/2 months (56 days) All other strengths: 1 pack/28 days</td></tr><tr><td>ZYPREXA RELPREVV (olanzapine)</td><td>IM</td><td>405 mg: 1 pack/28 days All other strengths: 1 pack/14 days</td></tr><tr><td colspan="3">*Requests for dosing regimens exceeding maximum may be approved for one year with prescriber attestation that the member is stabilized on the requested dose and schedule.</td></tr></table>	Table 1: FDA-Labeled Dosing Quantity Limits			Long-Acting injectable	Route	Quantity Limit	ABILIFY ASIMTUFII (aripiprazole)	IM	1 pack/2 months (56 days)	ABILIFY MAINTENA (aripiprazole)	IM	1 pack/28 days	ARISTADA ER (aripiprazole)	IM	1,064 mg: 1 pack/2 months (56 days) All other strengths: 1 pack/28 days	ARISTADA INITIO (aripiprazole)	IM	1 pack/7 weeks (49 days)	INVEGA HAFYERA (paliperidone)	IM	1 pack/6 months (168 days)	INVEGA SUSTENNA (paliperidone)	IM	156 mg: 2 packs/5 weeks (35 days) All other strengths: 1 pack/28 days	INVEGA TRINZA (paliperidone)	IM	1 pack/3 months (84 days)	PERSERIS ER (risperidone)	Subcutaneous	1 pack/28 days	RISPERDAL CONSTA (risperidone)	IM	2 packs/28 days	UZEDY ER (risperidone)	Subcutaneous	150 mg, 200 mg and 250 mg: 1 pack/2 months (56 days) All other strengths: 1 pack/28 days	ZYPREXA RELPREVV (olanzapine)	IM	405 mg: 1 pack/28 days All other strengths: 1 pack/14 days	*Requests for dosing regimens exceeding maximum may be approved for one year with prescriber attestation that the member is stabilized on the requested dose and schedule.		
Table 1: FDA-Labeled Dosing Quantity Limits																																												
Long-Acting injectable	Route	Quantity Limit																																										
ABILIFY ASIMTUFII (aripiprazole)	IM	1 pack/2 months (56 days)																																										
ABILIFY MAINTENA (aripiprazole)	IM	1 pack/28 days																																										
ARISTADA ER (aripiprazole)	IM	1,064 mg: 1 pack/2 months (56 days) All other strengths: 1 pack/28 days																																										
ARISTADA INITIO (aripiprazole)	IM	1 pack/7 weeks (49 days)																																										
INVEGA HAFYERA (paliperidone)	IM	1 pack/6 months (168 days)																																										
INVEGA SUSTENNA (paliperidone)	IM	156 mg: 2 packs/5 weeks (35 days) All other strengths: 1 pack/28 days																																										
INVEGA TRINZA (paliperidone)	IM	1 pack/3 months (84 days)																																										
PERSERIS ER (risperidone)	Subcutaneous	1 pack/28 days																																										
RISPERDAL CONSTA (risperidone)	IM	2 packs/28 days																																										
UZEDY ER (risperidone)	Subcutaneous	150 mg, 200 mg and 250 mg: 1 pack/2 months (56 days) All other strengths: 1 pack/28 days																																										
ZYPREXA RELPREVV (olanzapine)	IM	405 mg: 1 pack/28 days All other strengths: 1 pack/14 days																																										
*Requests for dosing regimens exceeding maximum may be approved for one year with prescriber attestation that the member is stabilized on the requested dose and schedule.																																												

		<i>Note: Effective January 14, 2022, no place of service prior authorization is required for extended-release injectable medications (LAIs) used for the treatment of mental health or substance use disorders (SUD), when administered by a healthcare professional and billed under the pharmacy benefit. In addition, LAIs may be administered in any setting (pharmacy, clinic, medical office or member home) and billed to the pharmacy or medical benefit as most appropriate and in accordance with all Health First Colorado billing policies.</i>
--	--	---

Table 1 Atypical Antipsychotics – FDA Approved Indication, Age Range, Quantity and Maximum Dose					
Brand	Generic	Approved Indications	Age Range	Maximum Daily Dose by Age/Indication	Quantity and Maximum Dose Limitations
ABILIFY	aripiprazole	Schizophrenia Bipolar I Disorder Bipolar I Disorder Irritability w/autistic disorder Tourette’s disorder Adjunctive treatment of MDD	≥ 13 years ≥ 18 years 10-17 years 6-17 years 6-18 years ≥ 18 years	30 mg 30 mg 30 mg 15 mg 20 mg (weight-based) 15 mg	Maximum one tablet per day (maximum of two tablets per day allowable for members < 18 years of age to accommodate for incremental dose changes)
CAPLYTA	lumateperone	Schizophrenia Bipolar I Disorder Bipolar II Disorder	≥ 18 years	42 mg	Maximum dosage of 42mg per day
CLOZARIL	clozapine	Treatment-resistant schizophrenia Recurrent suicidal behavior in schizophrenia or schizoaffective disorder	≥ 18 years	900 mg	Maximum dosage of 900mg per day
COBENFY	xanomeline and trospium	Schizophrenia	≥ 18 years	250 mg xanomeline and 60 mg trospium	Maximum two capsules per day
FANAPT	iloperidone	Schizophrenia Bipolar I Disorder	≥ 18 years	24 mg	Maximum two tablets per day
GEODON	ziprasidone	Schizophrenia Bipolar I Disorder	≥ 18 years ≥ 18 years	200 mg 160 mg	Maximum two capsules per day

	INVEGA ER	paliperidone	Schizophrenia & schizoaffective disorder	≥ 12 years and weight ≥ 51 kg ≥ 12 years and weight < 51 kg	12 mg 6 mg	Maximum two 6mg tablets per day; all other strengths 1 tablet per day	
	LATUDA	lurasidone	Schizophrenia Schizophrenia Bipolar I disorder Bipolar I disorder	≥ 18 years 13-17 years ≥ 18 years 10–17 years	160 mg 80 mg 120 mg 80 mg	Maximum one tablet per day (If dosing 160mg for schizophrenia, then max of two tablets per day)	
	LYBALVI	olanzapine and samidorphan	Schizophrenia in adults Bipolar I disorder in adults	≥ 18 years ≥ 18 years	20 mg olanzapine and 10 mg samidorphan	Maximum one tablet per day	
	NUPLAZID	pimavanserin	Parkinson's disease psychosis	≥ 18 years	34 mg	Maximum dosage of 34mg per day	
	RISPERDAL	risperidone	Schizophrenia Schizophrenia Bipolar mania Irritability w/autistic disorder	≥ 18 years 13-17 years ≥ 10 years 5–17 years	16 mg 6 mg 6 mg 3 mg	Maximum dosage of 16mg/day (4 tablet/day limitation applied in claims system to allow for dose escalation and tapering)	
	REXULTI	brexpiprazole	Schizophrenia Adjunctive treatment of MDD Agitation associated with Alzheimer's disease (AD)	≥ 13 years ≥ 18 years	4 mg 3 mg	Maximum of 3mg/day for MDD adjunctive therapy, and agitation due to AD, Maximum of 4mg/day for schizophrenia	
	SAPHRIS	asenapine	Schizophrenia Bipolar mania or mixed episodes	≥ 18 years ≥ 10 years	20 mg 20 mg	Maximum two tablets per day	
	SECUADO	asenapine patch	Schizophrenia	≥ 18 years	7.6 mg/ 24 hours	Maximum 1 patch per day	
	SEROQUEL	quetiapine	Schizophrenia Schizophrenia Bipolar I mania or mixed Bipolar I mania or mixed Bipolar I depression Bipolar I Disorder Maintenance	≥ 18 years 13-17 years ≥ 18 years 10-17 years ≥ 18 years ≥ 18 years	750 mg 800 mg 800 mg 600 mg 300 mg 800 mg	Maximum three tablets per day	
	SEROQUEL XR	quetiapine ER	Schizophrenia Bipolar I mania Bipolar I mania Bipolar I depression Adjunctive treatment of MDD	≥ 13 years ≥ 18 years 10-17 years ≥ 18 years ≥ 18 years	800 mg 800 mg 600 mg 300 mg 300 mg	Maximum one tablet per day (for 300mg & 400mg tablets max 2 tablets per day)	
	SYMBYAX	olanzapine/ fluoxetine	Acute depression in Bipolar I Disorder Treatment resistant depression (MDD)	≥ 10 years	12 mg olanzapine/ 50 mg fluoxetine	Maximum three capsules per day (18mg olanzapine/75mg fluoxetine)	
	VERSACLOZ	clozapine	Treatment-resistant schizophrenia Recurrent suicidal behavior in schizophrenia or schizoaffective disorder	≥ 18 years ≥ 18 years	900 mg	Maximum dosage of 900 mg per day	

	VRAYLAR	cariprazine	Schizophrenia Acute manic or mixed episodes with Bipolar I disorder Depressive episodes with Bipolar I disorder Adjunctive treatment of MDD	≥ 18 years ≥ 18 years ≥ 18 years ≥ 18 years	6 mg 6 mg 3 mg 3 mg	Maximum dosage of 6mg/day	
	ZYPREXA ZYPREXA ZYDIS	olanzapine	Schizophrenia Acute manic or mixed episodes with Bipolar I disorder	≥ 13 years	20 mg	Maximum one tablet per day	
Therapeutic Drug Class: CALCITONIN GENE – RELATED PEPTIDE INHIBITORS (CGRPi)s – <i>Effective 4/1/2025</i>							
PA Required for all agents			*Preferred agents may be approved if meeting the following criteria:				
Preferred		Non-Preferred	<u>Preferred Medications for Migraine Prevention (must meet all of the following):</u> <ul style="list-style-type: none">• The requested medication is being used as preventive therapy for episodic or chronic migraine AND• Member has diagnosis of migraine with or without aura AND• Member has tried and failed 2 oral preventive pharmacological agents listed as Level A per the most current American Headache Society/American Academy of Neurology guidelines (such as divalproex, topiramate, metoprolol, propranolol). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction OR• If the prescribed medication is Nurtec, the member has tried and failed two preferred injectable product formulations. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, significant drug-drug interaction, severe needle phobia, or member (or parent/caregiver) is unable to administer preferred CGRP inhibitor injectable formulation due to limited functional ability (such as vision impairment, limited manual dexterity and/or limited hand strength). <u>Preferred Medications for Acute Migraine Treatment (must meet all of the following):</u> <ul style="list-style-type: none">• The requested medication is being used as acute treatment for migraine headache AND• Member has history of trial and failure of two triptans (failure is defined as lack of efficacy with 4-week trial, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction, severe needle phobia, or member (or parent/caregiver) is unable to administer preferred triptan injectable formulation due to limited functional ability (such as vision impairment, limited manual dexterity and/or limited hand strength). <u>Non-Preferred Medications for Migraine Prevention (must meet all of the following):</u> <ul style="list-style-type: none">• The requested medication is being used as preventive therapy for episodic or chronic migraine AND• Member has diagnosis of migraine with or without aura AND• Member has tried and failed two oral preventive pharmacological agents listed as Level A per the most current American Headache Society/American Academy of Neurology guidelines (such as divalproex, topiramate, metoprolol, propranolol). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND• The requested medication is not being used in combination with another CGRP medication AND				
* AIMOVIG (erenumab-aooe) auto-injector		EMGALITY (galcanezumab-gnlm) 100 mg syringe					
* AJOVY (fremanezumab-vfrm) auto-injector, syringe							
* EMGALITY (galcanezumab-gnlm) pen, 120 mg syringe		QULIPTA (atogepant) tablet					
* NURTEC (rimegepant) ODT		ZAVZPRET (zavegepant) nasal					
* UBRELVY (ubrogepant) tablet							

		<ul style="list-style-type: none">• The member has history of adequate trial and failure of three preferred products indicated for preventive therapy (failure is defined as lack of efficacy with 4-week trial, contraindication to therapy, allergy, intolerable side effects, significant drug-drug interaction, severe needle phobia, or member (or parent/caregiver) is unable to administer preferred triptan injectable formulation due to limited functional ability (such as vision impairment, limited manual dexterity and/or limited hand strength). <p><u>Non-Preferred Medications for Acute Migraine Treatment (must meet all of the following):</u></p> <ul style="list-style-type: none">• Member is 18 years of age or older AND• Medication is being prescribed to treat migraine headache with moderate to severe pain AND• The requested medication is not being used in combination with another CGRP medication AND• Member has history of trial and failure with <u>all</u> of the following (failure is defined as lack of efficacy with 4-week trial, allergy, contraindication, intolerable side effects, or significant drug-drug interaction):<ul style="list-style-type: none">○ Two triptans AND○ One NSAID agent AND○ One preferred agent indicated for acute migraine treatment <p><u>Non-Preferred Medications for Treatment of Episodic Cluster Headache (must meet all of the following):</u></p> <ul style="list-style-type: none">• Member is 19-65 years of age AND• Member meets diagnostic criteria for episodic cluster headache (has had no more than 8 attacks per day, a minimum of one attack every other day, and at least 4 attacks during the week prior to this medication being prescribed) AND• Member is not taking other preventive medications to reduce the frequency of cluster headache attacks AND• Member has history of trial and failure of all of the following (failure is defined as lack of efficacy with 4-week trial, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction):<ul style="list-style-type: none">○ Oxygen therapy AND○ Sumatriptan subcutaneous or intranasal OR zolmitriptan intranasal• Initial authorization will be limited to 8 weeks. Continuation (12-month authorization) will require documentation of clinically relevant improvement with no less than 30% reduction in headache frequency in a 4-week period. <p><u>Age Limitations:</u></p> <p>All products: ≥ 18 years</p> <table><tr><th colspan="2">Table 1. Calcitonin Gene-Related Peptide Inhibitor Quantity Limits</th></tr><tr><th>Drug Name</th><th>Maximum Dosing</th></tr><tr><td>Aimovig (erenumab)</td><td>one 140 mg autoinjector per 30 days</td></tr><tr><td>Ajovy (fremanezumab)</td><td>one 225 mg autoinjector or syringe per 30 days or three 225 mg autoinjectors or syringes every 90 days</td></tr><tr><td>Emgality 100mg (galcanezumab)</td><td>three 100 mg prefilled syringes per 30 days</td></tr><tr><td>Emgality 120 mg (galcanezumab)</td><td>two 120 mg pens or prefilled syringes once as first loading dose then one 120 mg pen or prefilled syringe per 30 days</td></tr><tr><td>Nurtec (rimegepant)</td><td>Prevention: 16 tablets/30 days; Acute Treatment: 8 tablets/30 days</td></tr><tr><td>Qulipta (atogepant)</td><td>30 tablets/30 days</td></tr><tr><td>Ubrelvy 50 mg (ubrogepant)</td><td>16 tablets/30 days</td></tr></table>	Table 1. Calcitonin Gene-Related Peptide Inhibitor Quantity Limits		Drug Name	Maximum Dosing	Aimovig (erenumab)	one 140 mg autoinjector per 30 days	Ajovy (fremanezumab)	one 225 mg autoinjector or syringe per 30 days or three 225 mg autoinjectors or syringes every 90 days	Emgality 100mg (galcanezumab)	three 100 mg prefilled syringes per 30 days	Emgality 120 mg (galcanezumab)	two 120 mg pens or prefilled syringes once as first loading dose then one 120 mg pen or prefilled syringe per 30 days	Nurtec (rimegepant)	Prevention: 16 tablets/30 days; Acute Treatment: 8 tablets/30 days	Qulipta (atogepant)	30 tablets/30 days	Ubrelvy 50 mg (ubrogepant)	16 tablets/30 days
Table 1. Calcitonin Gene-Related Peptide Inhibitor Quantity Limits																				
Drug Name	Maximum Dosing																			
Aimovig (erenumab)	one 140 mg autoinjector per 30 days																			
Ajovy (fremanezumab)	one 225 mg autoinjector or syringe per 30 days or three 225 mg autoinjectors or syringes every 90 days																			
Emgality 100mg (galcanezumab)	three 100 mg prefilled syringes per 30 days																			
Emgality 120 mg (galcanezumab)	two 120 mg pens or prefilled syringes once as first loading dose then one 120 mg pen or prefilled syringe per 30 days																			
Nurtec (rimegepant)	Prevention: 16 tablets/30 days; Acute Treatment: 8 tablets/30 days																			
Qulipta (atogepant)	30 tablets/30 days																			
Ubrelvy 50 mg (ubrogepant)	16 tablets/30 days																			

		Ubrelvy 100 mg (ubrogepant)	16 tablets/30 days
		ZAVZPRET (zavegepant)	6 unit-dose nasal spray devices per 30 days
		Members with current prior authorization approval on file for a preferred agent may receive approval for continuation of therapy with the preferred agent.	
Therapeutic Drug Class: LITHIUM AGENTS – <i>Effective 4/1/2025</i>			
No PA Required Lithium carbonate capsule, tablet Lithium citrate solution Lithium ER tablet	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i> LITHOBID ER (lithium ER) tablet	Non-preferred products may be approved with trial and failure of one preferred agent (failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, significant drug-drug interactions, intolerance to dosage form). Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.	
Therapeutic Drug Class: NEUROCOGNITIVE DISORDER AGENTS – <i>Effective 4/1/2025</i>			
Preferred *Must meet eligibility criteria *Donepezil 5mg, 10mg tablet *Donepezil ODT *Galantamine IR tablet *Memantine IR tablet, dose pack *Memantine ER capsule *Rivastigmine capsule, patch	Non-Preferred PA Required ADLARITY (donepezil) patch ARICEPT (donepezil) tablet Donepezil 23mg tablet EXELON (rivastigmine) patch Galantamine solution, ER capsule Memantine IR solution MESTINON (pyridostigmine) IR/ER tablet, syrup Nemantine/donepezil ER capsule, NAMZARIC (memantine/donepezil ER) capsule, dose pack Pyridostigmine syrup, IR/ER tablet	*Eligibility criteria for Preferred Agents – Preferred products may be approved for a diagnosis of neurocognitive disorder (eligible for AutoPA automated approval). Non-preferred products may be approved if the member has failed treatment with one of the preferred products in the last 12 months. (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions) Members currently stabilized on a non-preferred product may receive approval to continue on that agent for one year if medically necessary and if there is a diagnosis of neurocognitive disorder.	

Therapeutic Drug Class: SEDATIVE HYPNOTICS – <i>Effective 4/1/2025</i>		
Non-Benzodiazepines		
<p>Preferred No PA Required* (Unless age, dose, or duplication criteria apply)</p> <p>Eszopiclone tablet</p> <p>Ramelteon tablet</p> <p>Zaleplon capsule</p> <p>Zolpidem IR, ER tablet</p>	<p>Non-Preferred PA Required</p> <p>AMBIEN (zolpidem) tablet</p> <p>AMBIEN CR (zolpidem ER) tablet</p> <p>BELSOMRA (suvorexant) tablet</p> <p>DAYVIGO (lemoborexant) tablet</p> <p>Doxepin tablet</p> <p>EDLUAR (zolpidem) SL tablet</p> <p>HETLIOZ (tasimelteon) capsule</p> <p>HETLIOZ LQ (tasimelteon) suspension</p> <p>LUNESTA (eszopiclone) tablet</p> <p>QUVIVIQ (daridorexant) tablet</p> <p>ROZEREM (ramelteon) tablet</p> <p>SILENOR (doxepin) tablet</p> <p>Tasimelteon capsule</p> <p>Zolpidem capsule, SL tablet</p>	<p>Non-preferred non-benzodiazepine sedative hypnotics may be approved for members who have failed treatment with two preferred non-benzodiazepine agents (failure is defined as lack of efficacy with a 2-week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p><u>Children:</u> Prior authorization will be required for all agents for members < 18 years of age.</p> <p><u>Duplications:</u> Only one agent in the sedative hypnotic drug class will be approved at a time (concomitant use of agents in the same sedative hypnotic class or differing classes will not be approved).</p> <p>All sedative hypnotics will require prior authorization for members ≥ 65 years of age when exceeding 90 days of therapy.</p> <p>Belsomra (suvorexant) may be approved for adult members that meet the following:</p> <ul style="list-style-type: none">• Member has trialed and failed therapy with two preferred agents (failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) AND• Member is not receiving strong CYP3A4 inhibitors (such as erythromycin, clarithromycin, telithromycin, itraconazole, ketoconazole, posaconazole, fluconazole, voriconazole, delavirdine, and milk thistle) or strong CYP3A4 inducers (such as carbamazepine, oxcarbazepine, phenobarbital, phenytoin, rifampin, rifabutin, rifapentine, dexamethasone, efavirenz, etravirine, nevirapine, darunavir/ritonavir, ritonavir, and St John’s Wort) AND• Member does not have a diagnosis of narcolepsy <p>Dayvigo (lemborexant) may be approved for adult member that meet the following:</p> <ul style="list-style-type: none">• Member has trialed and failed therapy with two preferred agents AND Belsomra (surovexant). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND• Member is not receiving strong CYP3A4 inhibitors (such as erythromycin, clarithromycin, telithromycin, itraconazole, ketoconazole, posaconazole, fluconazole, voriconazole, delavirdine, and milk thistle) or strong CYP3A4 inducers (such as carbamazepine, oxcarbazepine, phenobarbital, phenytoin, rifampin, rifabutin, rifapentine, dexamethasone, efavirenz, etravirine, nevirapine, darunavir/ritonavir, ritonavir, and St John’s Wort) AND• Member does not have a diagnosis of narcolepsy <p>Hetlioz (tasimelteon) capsules may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none">• Member is ≥18 years of age and has a documented diagnosis of Non-24-hour sleep wake disorder (Non-24) OR• Member is ≥16 years of age and has a documented diagnosis of nighttime sleep disturbances in Smith-Magenis syndrome (SMS) AND• The requested medication is being prescribed by a sleep specialist or a practitioner who has sufficient education and experience to safely prescribe tasimelteon <p>Hetlioz LQ (tasimelteon) oral suspension may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none">• Member is 3 to 15 years of age and has a documented diagnosis of nighttime sleep disturbances in Smith-Magenis Syndrome (SMS)

		<ul style="list-style-type: none">AND the requested medication is being prescribed by a sleep specialist or a practitioner who has sufficient education and experience to safely prescribe tasimelteon. <p>Silenor (doxepin) may be approved for adult members that meet ONE of the following criteria:</p> <ul style="list-style-type: none">Member has tried and failed two preferred oral sedative hypnotics (Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) ORProvider attests to the medical necessity of prescribing individual doxepin doses of less than 10 mg, ORMember’s age is ≥ 65 years <p>Prior authorization will be required for prescribed doses exceeding maximum (Table 1) below.</p>
--	--	---

Benzodiazepines		
<p>Preferred No PA Required* (Unless age, dose, or duplication criteria apply)</p> <p>Temazepam 15mg, 30mg capsule</p> <p>Triazolam tablet</p>	<p>Non-Preferred PA Required</p> <p>DORAL (quazepam) tablet</p> <p>Estazolam tablet</p> <p>Flurazepam capsule</p> <p>HALCION (triazolam) tablet</p> <p>Quazepam tablet</p> <p>RESTORIL (temazepam) capsule</p> <p>Temazepam 7.5mg, 22.5mg capsule</p>	<p>Non-preferred benzodiazepine sedative hypnotics may be approved for members who have trialed and failed therapy with two preferred benzodiazepine agents (failure is defined as lack of efficacy with a 2-week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>Temazepam 22.5 mg may be approved if the member has trialed and failed temazepam 15mg or 30mg AND one other preferred product (failure is defined as lack of efficacy with a 2-week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>Temazepam 7.5 mg may be approved if provider attests to the medical necessity of prescribing individual temazepam doses of less than 15 mg.</p> <p><u>Children:</u> Prior authorization will be required for all sedative hypnotic agents when prescribed for members < 18 years of age.</p> <p><u>Duplications:</u> Only one agent in the sedative hypnotic drug class will be approved at a time (concomitant use of agents in the same sedative hypnotic class or differing classes will not be approved).</p> <p>All sedative hypnotics will require prior authorization for member’s ≥ 65 years of age when exceeding 90 days of therapy.</p> <p>Members currently stabilized on a non-preferred benzodiazepine medication may receive authorization to continue that medication.</p> <p>Prior authorization will be required for prescribed doses exceeding maximum (Table 1).</p>

Table 1: Sedative Hypnotic Maximum Dosing		
Brand	Generic	Maximum Dose
Non-Benzodiazepine		
Ambien CR	Zolpidem CR	12.5 mg/day
Ambien IR	Zolpidem IR	10 mg/day
Belsomra	Suvorexant	20 mg/day
Dayvigo	Lemborexant	10 mg/day
Edluar	Zolpidem sublingual	10 mg/day

		<table><tr><td>-</td><td>Zolpidem sublingual</td><td>Men: 3.5mg/day Women: 1.75 mg/day</td></tr><tr><td>Hetlioz</td><td>Tasimelteon capsule</td><td>20 mg/day</td></tr><tr><td>Hetlioz LQ</td><td>Tasimelteon liquid</td><td>≤ 28 kg: 0.7 mg/kg/day > 28 kg : 20 mg/day</td></tr><tr><td>Lunesta</td><td>Eszopiclone</td><td>3 mg/day</td></tr><tr><td>Quviviq</td><td>Daridorexant</td><td>50 mg/day</td></tr><tr><td>-</td><td>Zaleplon</td><td>20 mg/day</td></tr><tr><td>Rozerem</td><td>Ramelteon</td><td>8 mg/day</td></tr><tr><td colspan="3">Benzodiazepine</td></tr><tr><td>Halcion</td><td>Triazolam</td><td>0.5 mg/day</td></tr><tr><td>Restoril</td><td>Temazepam</td><td>30 mg/day</td></tr><tr><td>Silenor</td><td>Doxepin</td><td>6mg/day</td></tr><tr><td>-</td><td>Estazolam</td><td>2 mg/day</td></tr><tr><td>-</td><td>Flurazepam</td><td>30 mg/day</td></tr><tr><td>Doral</td><td>Quazepam</td><td>15 mg/day</td></tr></table>	-	Zolpidem sublingual	Men: 3.5mg/day Women: 1.75 mg/day	Hetlioz	Tasimelteon capsule	20 mg/day	Hetlioz LQ	Tasimelteon liquid	≤ 28 kg: 0.7 mg/kg/day > 28 kg : 20 mg/day	Lunesta	Eszopiclone	3 mg/day	Quviviq	Daridorexant	50 mg/day	-	Zaleplon	20 mg/day	Rozerem	Ramelteon	8 mg/day	Benzodiazepine			Halcion	Triazolam	0.5 mg/day	Restoril	Temazepam	30 mg/day	Silenor	Doxepin	6mg/day	-	Estazolam	2 mg/day	-	Flurazepam	30 mg/day	Doral	Quazepam	15 mg/day
-	Zolpidem sublingual	Men: 3.5mg/day Women: 1.75 mg/day																																										
Hetlioz	Tasimelteon capsule	20 mg/day																																										
Hetlioz LQ	Tasimelteon liquid	≤ 28 kg: 0.7 mg/kg/day > 28 kg : 20 mg/day																																										
Lunesta	Eszopiclone	3 mg/day																																										
Quviviq	Daridorexant	50 mg/day																																										
-	Zaleplon	20 mg/day																																										
Rozerem	Ramelteon	8 mg/day																																										
Benzodiazepine																																												
Halcion	Triazolam	0.5 mg/day																																										
Restoril	Temazepam	30 mg/day																																										
Silenor	Doxepin	6mg/day																																										
-	Estazolam	2 mg/day																																										
-	Flurazepam	30 mg/day																																										
Doral	Quazepam	15 mg/day																																										
Therapeutic Drug Class: SKELETAL MUSCLE RELAXANTS – <i>Effective 4/1/2025</i>																																												
No PA Required (*if under 65 years of age)	PA Required																																											
Baclofen tablet	AMRIX ER (cyclobenzaprine ER) capsule	<p>All agents in this class will require a PA for members 65 years of age and older. The maximum allowable approval will be for a 7-day supply.</p> <p>Authorization for any CARISOPRODOL product will be given for a maximum 3-week one-time authorization for members with acute, painful musculoskeletal conditions who have failed treatment with three preferred products within the last 6 months.</p> <p>*Dantrolene may be approved for members who have trialed and failed‡ one preferred agent and meet the following criteria:</p> <ul style="list-style-type: none">● Documentation of age-appropriate liver function tests AND● One of following diagnoses: Multiple Sclerosis, Cerebral Palsy, stroke, upper motor neuron disorder, or spinal cord injury● Dantrolene will be approved for the period of one year● If a member is stabilized on dantrolene, they may continue to receive approval <p>All other non-preferred skeletal muscle relaxants may be approved for members who have trialed and failed‡ three preferred agents. ‡Failure is defined as: lack of efficacy with 14-day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions.</p>																																										
Cyclobenzaprine tablet	Baclofen solution, suspension																																											
Methocarbamol tablet	Carisoprodol tablet																																											
Tizanidine tablet	Carisoprodol/Aspirin tablet																																											
	Chlorzoxazone tablet																																											
	Cyclobenzaprine ER capsule																																											
	DANTRIUM (dantrolene) capsule																																											
	*Dantrolene capsule																																											
	FEXMID (cyclobenzaprine) tablet																																											
	FLEQSUVY (baclofen) solution																																											
	LORZONE (chlorzoxazone) tablet																																											
	LYVISPAH (baclofen) granules																																											

	Metaxalone tablet NORGESIC/NORGESIC FORTE (orphenadrine/aspirin/ caffeine) tablet Orphenadrine ER tablet Orphenadrine/Aspirin/Caffeine tablet SOMA (carisoprodol) tablet Tizanidine capsule ZANAFLEX (tizanidine) capsule, tablet	
Therapeutic Drug Class: STIMULANTS AND RELATED AGENTS – <i>Effective 4/1/2025</i>		
<p>Preferred *No PA Required (if age, max daily dose, and diagnosis met)</p> <p><i>Brand/generic changes effective 08/08/2024</i></p> <p>Amphetamine salts, mixed ER (generic Adderall XR) capsule</p> <p>Amphetamine salts, mixed (generic Adderall IR) tablet</p> <p>Armodafinil tablet</p> <p>Atomoxetine capsule</p> <p>Clonidine ER tablet</p> <p>DAYTRANA^{BNR} (methylphenidate) patch</p> <p>Dexmethylphenidate IR tablet</p> <p>Dexmethylphenidate ER capsule</p> <p>Guanfacine ER tablet</p> <p>Methylphenidate (generic Methylin/Ritalin) solution, tablet</p> <p>Methylphenidate ER tablet (generic Concerta)</p>	<p>Non-Preferred PA Required</p> <p>ADDERALL IR (amphetamine salts, mixed IR) tablet</p> <p>ADDERALL XR (amphetamine salts, mixed ER) capsule</p> <p>ADZENYS XR-ODT (amphetamine)</p> <p>Amphetamine tablet (generic Evekeo)</p> <p>APTENSIO XR (methylphenidate ER) capsule</p> <p>AZSTARYS (serdexmethylphenidate/dexmethylphenidate) capsule</p> <p>CONCERTA (methylphenidate ER) tablet</p> <p>COTEMPLA XR-ODT (methylphenidate ER)</p>	<p>*Preferred medications may be approved through AutoPA for indications listed in Table 1 (preferred medications may also receive approval for off-label use for fatigue associated with multiple sclerosis).</p> <p>Non-preferred medications may be approved for members meeting the following criteria (for Sunosi (solriamfetol) and Wakix (pitolisant), refer to specific criteria listed below):</p> <ul style="list-style-type: none">• Prescription meets indication/age limitation criteria (Table 1) AND• <u>If member is ≥ 6 years of age:</u><ul style="list-style-type: none">○ Has documented trial and failure‡ with three preferred products in the last 24 months AND○ If the member is unable to swallow solid oral dosage forms, two of the trials must be methylphenidate solution, dexmethylphenidate ER, Vyvanse, Adderall XR, or any other preferred product that can be taken without the need to swallow a whole capsule. <p>OR</p> <ul style="list-style-type: none">• <u>If member is 3–5 years of age:</u><ul style="list-style-type: none">○ Has documented trial and failure‡ with one preferred product in the last 24 months AND○ If the member is unable to swallow solid oral dosage forms, the trial must be methylphenidate solution, dexmethylphenidate ER, Vyvanse, Adderall XR, or any other preferred product that can be taken without the need to swallow a whole capsule. <p>SUNOSI (solriamfetol) prior authorization may be approved if member meets the following criteria:</p> <ul style="list-style-type: none">• Member is 18 years of age or older AND• Member has diagnosis of either narcolepsy or obstructive sleep apnea (OSA) and is experiencing excessive daytime sleepiness AND• Member does not have end stage renal disease AND• If Sunosi is being prescribed for OSA, member has 1 month trial of CPAP AND

Modafinil tablet VYVANSE ^{BNR} (lisdexamfetamine) capsule	DESOXYN (methamphetamine) tablet	<ul style="list-style-type: none">Member has trial and failure[‡] of modafinil AND armodafinil AND one other agent in stimulant PDL class. <p>WAKIX (pitolisant) prior authorization may be approved if member meets the following criteria:</p> <ul style="list-style-type: none">Member is 6 years of age or older ANDMember has diagnosis of narcolepsy and is experiencing excessive daytime sleepiness ANDMember does not have end stage renal disease (eGFR <15 mL/minute) ANDMember does not have severe hepatic impairment ANDMember has trial and failure[‡] of modafinil AND armodafinil AND one other agent in the stimulant PDL class ANDMember has been counseled that Wakix may reduce the efficacy of hormonal contraceptives and counseled regarding use of an alternative non-hormonal method of contraception during Wakix therapy and for at least 21 days after discontinuing treatment. <p>Maximum Dose (all products): See Table 2</p> <p>Exceeding Maximum Dose: Prior authorization may be approved for doses that are higher than the listed maximum dose (Table 2) for members meeting the following criteria:</p> <ul style="list-style-type: none">Member is taking medication for indicated use listed in Table 1 ANDMember has 30-day trial and failure[‡] of three different preferred or non-preferred agents at maximum doses listed in Table 2 ANDDocumentation of member’s symptom response to maximum doses of three other agents is provided ANDMember is not taking a sedative hypnotic medication (such as temazepam, triazolam, or zolpidem from the Sedative Hypnotic PDL class). <p>[‡]Failure is defined as: lack of efficacy with 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p>
	DEXEDRINE (dextroamphetamine) Spansule	
	Dextroamphetamine ER capsule, solution, tablet	
	DYANAVEL XR (amphetamine) suspension, tablet	
	EVEKEO (amphetamine) ODT, tablet	
	FOCALIN (dexmethylphenidate) tablet, XR capsule	
	INTUNIV (guanfacine ER) tablet	
	JORNAY PM (methylphenidate) capsule	
	Lisdexamfetamine capsule, chewable tablet	
	Methamphetamine tablet	
	METHYLIN (methylphenidate) solution	
	Methylphenidate CD/ER/LA capsule, chewable tablet, ER tablet (generic Relexxi/Ritalin), patch	
	MYDAYIS ER (dextroamphetamine/amphetamine) capsule	
	NUVIGIL (armodafinil) tablet	

	<div>PROVIGIL (modafinil) tablet</div> <div>QELBREE (viloxazine ER) capsule</div> <div>QUILLICHEW ER (methylphenidate) chewable tablet, XR suspension</div> <div>RELEXXII (methylphenidate ER) tablet</div> <div>RITALIN (methylphenidate) IR/ER tablet, ER capsule</div> <div>STRATTERA (atomoxetine) capsule</div> <div>SUNOSI (solriamfetol) tablet</div> <div>VYVANSE (lisdexamfetamine) chewable tablet</div> <div>WAKIX (pitolisant) tablet</div> <div>XELSTRYM (dextroamphetamine) patch</div> <div>ZENZEDI (dextroamphetamine) tablet</div>	
--	---	--

Table 1: Diagnosis and Age Limitations	
<ul style="list-style-type: none">Approval for medically accepted indications <u>not</u> listed in Table 1 may be given with prior authorization review and may require submission of peer-reviewed literature or medical compendia showing safety and efficacy of the medication used for the prescribed indication.Preferred medications may also receive approval for off-label use for fatigue associated with multiple sclerosis if meeting all other criteria for approval.Bolded drug names are preferred (subject to preferential coverage changes for brand/generic equivalents)	
Drug	Diagnosis and Age Limitations
Stimulants–Immediate Release	
Amphetamine sulfate (EVEKEO)	ADHD (Age ≥ 3 years), Narcolepsy (Age ≥ 6 years)
Dexmethylphenidate IR (FOCALIN)	ADHD (Age ≥ 6 years)
Dextroamphetamine IR tablet (ZENZEDI)	ADHD (Age 3 to16 years), Narcolepsy (Age ≥ 6 years)
Dextroamphetamine solution (PROCENTRA)	ADHD (Age 3 to 16 years), Narcolepsy (Age ≥ 6 years)
Methamphetamine (DESOXYN)	ADHD (Age ≥ 6 years)
methylphenidate IR (generic METHYLIN, RITALIN)	ADHD (Age ≥ 6 years [†]), Narcolepsy (Age ≥ 6 years), OSA.

		<p>†Prior Authorization for members 3-6 years of age with a diagnosis of ADHD may be approved with prescriber attestation to the following:</p> <ul style="list-style-type: none">• Member’s symptoms have not significantly improved despite adequate behavior interventions AND• Member experiences moderate-to-severe continued disturbance in functioning AND• Prescriber has determined that the potential benefits of starting methylphenidate before the age of 6 years outweigh the potential harm of delaying treatment.	
	Mixed amphetamine salts IR (generic ADDERALL)	ADHD (Age ≥ 3 years), Narcolepsy (Age ≥ 6 years)	
	Stimulants –Extended-Release		
	Amphetamine ER (ADZENYS XR-ODT and ADZENYS ER suspension)	ADHD (Age ≥ 6 years)	
	Amphetamine ER (DYANAVEL XR)	ADHD (Age ≥ 6 years)	
	Mixedamphetamine salts ER (ADDERALL XR)	ADHD (Age ≥ 6 years)	
	Dexmethylphenidate ER (generic Focalin XR)	ADHD (Age ≥ 6 years)	
	Dextroamphetamine ER (DEXEDRINE)	ADHD (Age 6 to 16 years), Narcolepsy (Age ≥ 6 years)	
	Dextroamphetamine ER/amphetamine ER (MYDAYIS ER)	ADHD (Age ≥ 13 years)	
	Dextroamphetamine ER patch (XELSTRYM)	ADHD (Age ≥ 6 years)	
	Lisdexamfetamine dimesylate (VYVANSE capsule, Vyvanse chewable)	ADHD (Age ≥ 6 years), Moderate to severe binge eating disorder in adults (Age ≥ 18 years)	
	Methylphenidate ER OROS (CONCERTA)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years), OSA	
	Methylphenidate patch (DAYTRANA)	ADHD (Age ≥ 6 years)	
	Methylphenidate SR (METADATE ER)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years)	
	Methylphenidate ER (METADATE CD)	ADHD (Age ≥ 6 years)	
	Methylphenidate ER (QUILLICHEW ER)	ADHD (Age 6 years to ≤ 65 years), Narcolepsy (Age ≥ 6 years)	
	Methylphenidate ER (QUILLIVANT XR)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years)	
	Methylphenidate ER (RELEXXI ER)	ADHD (Age 6 to 65 years)	
	Methylphenidate ER (RITALIN LA)	<p>ADHD (Age ≥ 6 years)</p> <p>†Prior Authorization for members 4-6 years of age with a diagnosis of ADHD may be approved with prescriber attestation to the following:</p> <ul style="list-style-type: none">• Member’s symptoms have not significantly improved despite adequate behavior interventions AND• Member experiences moderate-to-severe continued disturbance in functioning AND <p>Prescriber has determined that the potential benefits of starting methylphenidate before the age of 6 years outweigh the potential harm of delaying treatment.</p>	
	Methylphenidate ER (ADHANSIA XR)	ADHD (Age ≥ 6 years)	
	Methylphenidate ER (JORNAY PM)	ADHD (Age ≥ 6 years)	
	Methylphenidate XR (APTENSIO XR)	ADHD (Age ≥ 6 years)	

Methylphenidate XR ODT (COTEMPLA XR-ODT)	ADHD (Age 6 to 17 years)
Serdexmethylphenidate/dexmethylphenidate (AZSTARYS)	ADHD (Age ≥ 6 years)
Non-Stimulants	
Atomoxetine (generic STRATTERA)	ADHD (Age ≥ 6 years)
Clonidine ER	ADHD as monotherapy or adjunctive therapy to stimulants (Age ≥ 6 years)
Guanfacine ER (generic INTUNIV)	ADHD as monotherapy or adjunctive therapy to stimulants (Age ≥ 6 years)
Viloxazine ER (QELBREE)	ADHD (Age ≥ 6 years)
Wakefulness-promoting Agents	
Armodafinil (generic NUVIGIL)	Excessive sleepiness associated with narcolepsy, OSA, SWD, and adjunct therapy to treat fatigue and sleepiness in patients with major depressive disorder (MDD) (Age ≥ 18 years)
Modafinil (PROVIGIL)	Excessive sleepiness associated with narcolepsy, OSA, SWD, and adjunct therapy to treat fatigue and sleepiness in patients with major depressive disorder (MDD), antipsychotic medication-related fatigue (Age ≥ 18 years)
Pitolisant (WAKIX)	Excessive sleepiness associated with narcolepsy (Age ≥ 6 years)
Solriamfetol (SUNOSI)	Excessive sleepiness associated with narcolepsy, OSA (Age ≥ 18 years)
KEY: ADHD —attention-deficit/hyperactivity disorder, OSA —obstructive sleep apnea, SWD —shift work disorder	

Table 2: Maximum Dose	
Drug	Maximum Daily Dose
ADDERALL	60 mg
ADDERALL XR	60 mg
ADHANSIA XR	85 mg
ADZENYS XR ODT	18.8 mg (age 6-12)
ADZENYS ER SUSPENSION	12.5 mg (age ≥ 13)
AMPHETAMINE SALTS	40 mg
APTENSIO XR	60 mg
CONCERTA	54 mg (age 6-12) or 72 mg (≥ age 13)
AZSTARYS	52.3 mg serdexmethylphenidate and 10.4 mg dexmethylphenidate
CLONIDINE ER	0.4 mg
COTEMPLA XR-ODT	51.8 mg
DEXTROAMPHETAMINE ER	60 mg
DAYTRANA	30 mg/9 hour patch (3.3 mg/hr)
DESOXYN	25 mg
DEXEDRINE	60 mg
DYANAVEL XR	20 mg
EVEKEO	60 mg
FOCALIN	20 mg
FOCALIN XR	40 mg
GUANFACINE ER	4 mg (age 6-12) or 7 mg (age ≥ 13)

	INTUNIV ER	4 mg (age 6-12) or 7 mg (age ≥ 13)					
	JORNAY PM	100 mg					
	METADATE CD	60 mg					
	METADATE ER	60 mg					
	METHYLIN	60 mg					
	METHYLIN ER	60 mg					
	METHYLIN SUSPENSION	60 mg					
	METHYLPHENIDATE	60 mg					
	METHYLPHENIDATE ER	60 mg					
	MYDAYIS ER	25 mg (age 13-17) or 50 mg (age ≥ 18)					
	NUVIGIL	250 mg					
	PROCENTRA	60 mg					
	PROVIGIL	400 mg					
	QELBREE	400 mg (age 6-17) or 600 mg (age ≥ 18)					
	QUILLICHEW ER	60 mg					
	QUILLIVANT XR	60 mg					
	RELEXXII	54 mg (ages 6-12) or 72 mg (≥ age 13)					
	RITALIN IR	60 mg					
	RITALIN SR	60 mg					
	RITALIN LA	60 mg					
	STRATTERA	100mg					
	SUNOSI	150 mg					
	VYVANSE CAPSULES AND CHEWABLE TABLETS	70 mg					
	WAKIX	35.6 mg					
	XELSTRYM ER PATCH	18 mg/9 hours					
	ZENZEDI	60 mg					
Therapeutic Drug Class: TRIPTANS, DITANS AND OTHER MIGRAINE TREATMENTS - Oral – <i>Effective 4/1/2025</i>							
No PA Required (Quantity limits may apply)	PA Required	Reyvow (lasmiditan) may be approved if meeting the following: <ul style="list-style-type: none">Member has trialed and failed three preferred products OR member is unable to use triptan therapy due to cardiovascular risk factors AND <ul style="list-style-type: none">Member has trialed and failed two preferred agents in the CGRP Inhibitors drug class indicated for the acute treatment of migraine. All other non-preferred oral products may be approved for members who have trialed and failed three preferred oral products. Failure is defined as lack of efficacy with 4-week trial, allergy, documented contraindication to therapy, intolerable side effects, or significant drug-drug interaction. Quantity Limits: <table><tr><td>Amerge (naratriptan), Frova (frovatriptan), Imitrex (sumatriptan), Zomig (zolmitriptan)</td><td>9 tabs/30 days</td></tr><tr><td>Treximet (sumatriptan/naproxen)</td><td>9 tabs/30 days</td></tr></table>		Amerge (naratriptan), Frova (frovatriptan), Imitrex (sumatriptan), Zomig (zolmitriptan)	9 tabs/30 days	Treximet (sumatriptan/naproxen)	9 tabs/30 days
Amerge (naratriptan), Frova (frovatriptan), Imitrex (sumatriptan), Zomig (zolmitriptan)	9 tabs/30 days						
Treximet (sumatriptan/naproxen)	9 tabs/30 days						
Eletriptan tablet (generic Relpax)	Almotriptan tablet						
Naratriptan tablet (generic Amerge)	FROVA (frovatriptan) tablet						
Rizatriptan tablet, ODT (generic Maxalt)	Frovatriptan tablet						
Sumatriptan tablet (generic Imitrex)	IMITREX (sumatriptan) tablet						
Zolmitriptan tablet (generic Zomig)	MAXALT/MAXALT MLT (rizatriptan) tablet, ODT						
	RELPAx (eletriptan) tablet						

	REYVOW (lasmiditan) tablet	Axert (almotriptan) and Relpax (eletriptan)	6 tabs/30 days	
	Sumatriptan/Naproxen tablet	Maxalt (rizatriptan)	12 tabs/30 days	
	SYMBRAVO (rizatriptan/meloxicam) tablet	Reyvow (lasmiditan)	8 tabs/30 days	
	Zolmitriptan ODT			
	ZOMIG (zolmitriptan) tablet			

Therapeutic Drug Class: **TRIPTANS, DITANS, AND OTHER MIGRAINE TREATMENTS - Non-Oral** – *Effective 4/1/2025*

No PA Required (Quantity limits may apply)	PA Required																	
Dihydroergotamine nasal spray	Dihydroergotamine injection	<p>Zembrace Symtouch injection, Tosymra nasal spray, or Onzetra Xsail nasal powder may be approved for members who have trialed and failed one preferred non-oral triptan products AND two oral triptan agents with different active ingredients. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects, significant drug-drug interaction, or documented inability to take alternative dosage form.</p> <p>All other non-preferred products may be approved for members who have trialed and failed one preferred non-oral triptan product AND one preferred oral triptan product. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions, documented inability to tolerate dosage form.</p> <p>Quantity Limits:</p> <table><tr><td>Dihydroergotamine mesylate vial 1mg/mL</td><td>24 vials/ 28 days</td></tr><tr><td>Imitrex (sumatriptan) injection</td><td>4 injectors / 30 days</td></tr><tr><td>Imitrex (sumatriptan) nasal spray</td><td>6 inhalers / 30 days</td></tr><tr><td>Migranal (dihydroergotamine mesylate) nasal spray</td><td>8 nasal spray devices/ 30 days</td></tr><tr><td>Onzetra Xsail (sumatriptan) nasal powder</td><td>16 nosepieces / 30 days</td></tr><tr><td>Tosymra (sumatriptan) nasal spray</td><td>12 nasal spray devices / 30 days</td></tr><tr><td>Zembrace Symtouch (sumatriptan) injection</td><td>36mg / 30 days</td></tr><tr><td>Zomig (zolmitriptan) nasal spray</td><td>6 inhalers / 30 days</td></tr></table> <p>Members currently utilizing a non-oral dihydroergotamine product formulation (based on recent claims history) may receive one year approval to continue therapy with that medication.</p>	Dihydroergotamine mesylate vial 1mg/mL	24 vials/ 28 days	Imitrex (sumatriptan) injection	4 injectors / 30 days	Imitrex (sumatriptan) nasal spray	6 inhalers / 30 days	Migranal (dihydroergotamine mesylate) nasal spray	8 nasal spray devices/ 30 days	Onzetra Xsail (sumatriptan) nasal powder	16 nosepieces / 30 days	Tosymra (sumatriptan) nasal spray	12 nasal spray devices / 30 days	Zembrace Symtouch (sumatriptan) injection	36mg / 30 days	Zomig (zolmitriptan) nasal spray	6 inhalers / 30 days
Dihydroergotamine mesylate vial 1mg/mL	24 vials/ 28 days																	
Imitrex (sumatriptan) injection	4 injectors / 30 days																	
Imitrex (sumatriptan) nasal spray	6 inhalers / 30 days																	
Migranal (dihydroergotamine mesylate) nasal spray	8 nasal spray devices/ 30 days																	
Onzetra Xsail (sumatriptan) nasal powder	16 nosepieces / 30 days																	
Tosymra (sumatriptan) nasal spray	12 nasal spray devices / 30 days																	
Zembrace Symtouch (sumatriptan) injection	36mg / 30 days																	
Zomig (zolmitriptan) nasal spray	6 inhalers / 30 days																	
IMITREX (sumatriptan) nasal spray	IMITREX (sumatriptan) cartridge, pen injector																	
Sumatriptan cartridge, pen injector	TOSYMRA (sumatriptan) nasal spray																	
MIGRANAL ^{BNR} (dihydroergotamine) nasal spray	TRUDHESA (dihydroergotamine) nasal spray																	
Sumatriptan nasal spray*, vial	ZEMBRACE SYMTOUCH (sumatriptan) auto-injector																	
	Zolmitriptan nasal spray																	
	ZOMIG (zolmitriptan) nasal spray																	

V. Dermatological

Therapeutic Drug Class: **ACNE AGENTS– Topical** – *Effective 10/1/2025*

Preferred No PA Required (if age and diagnosis criteria are met*)	Non-Preferred PA Required	Authorization will not be approved for acne agents prescribed solely for cosmetic purposes.
*Adapalene gel		Preferred topical clindamycin and erythromycin products may be approved by AutoPA verification of ICD-10 diagnosis code for acne vulgaris, psoriasis, cystic acne, comedonal acne, disorders of keratinization, neoplasms, folliculitis,

<div>*Adapalene/benzoyl peroxide gel (generic Epiduo), gel pump (generic Epiduo Forte)</div> <div>*Clindamycin phosphate gel, lotion, solution, medicated swab/pledget</div> <div>*Clindamycin/benzoyl peroxide gel jar (generic Benzaclin)</div> <div>*Clindamycin/benzoyl peroxide gel tube (generic Duac)</div> <div>*Dapsone gel</div> <div>*Erythromycin solution</div> <div>*Erythromycin/Benzoyl peroxide gel (generic Benzamycin)</div> <div>*Sulfacetamide sodium suspension</div> <div>*Tretinoin cream</div> <div>*Tretinoin gel (Mylan only)</div>	<div>ACANYA (clindamycin/benzoyl peroxide) gel, pump</div> <div>Adapalene cream, gel pump, solution</div> <div>ALTRENO (tretinoin) lotion</div> <div>ARAZLO (tazarotene) lotion</div> <div>ATRALIN (tretinoin) gel</div> <div>BENZAMYCIN (erythromycin/benzoyl peroxide) gel</div> <div>BP (sulfacetamide sodium/sulfur/urea) cleansing wash</div> <div>CABTREO (adapalene/benzoyl peroxide/clindamycin) gel</div> <div>CLEOCIN-T (clindamycin) lotion</div> <div>CLINDACIN ETZ/PAC (clindamycin phosphate) kit</div> <div>CLINDAGEL gel</div> <div>Clindamycin phosphate foam</div> <div>Clindamycin/Benzoyl peroxide gel pump</div> <div>Clindamycin/tretinoin gel</div> <div>Dapsone gel pump</div> <div>ERY/ERYGEL (erythromycin/ethanol) gel, medicated swabs/pads</div> <div>Erythromycin gel</div> <div>EVOCLIN (clindamycin) foam</div>	<div>hidradenitis suppurativa, or perioral dermatitis (erythromycin only). Approval of preferred topical clindamycin and erythromycin products for other medically accepted indications may be considered following clinical prior authorization review by a call center pharmacist.</div> <div>All other preferred topical acne agents may be approved if meeting the following criteria:</div> <div><ul style="list-style-type: none">For members > 25 years of age, may be approved following prescriber verification that the medication is not being utilized for cosmetic purposes AND prescriber verification that the indicated use is for acne vulgaris, psoriasis, cystic acne, disorders of keratinization, neoplasms, or comedonal acne. These medications are only eligible for prior authorization approval for the aforementioned diagnoses.For members ≤ 25 years of age, may be approved for a diagnosis of acne vulgaris, psoriasis, cystic acne, disorders of keratinization, neoplasms, or comedonal acne. Diagnosis will be verified through automated verification (AutoPA) of the appropriate corresponding ICD-10 diagnosis code related to the indicated use of the medication.</div> <div>Non-preferred topical products may be approved for members meeting all of the following criteria:</div> <div><ul style="list-style-type: none">Member has trialed/failed three preferred topical products with different mechanisms (such as tretinoin, antibiotic). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction ANDPrescriber verification that the medication is being prescribed for one of the following diagnoses: acne vulgaris, psoriasis, cystic acne, disorders of keratinization, neoplasms, or comedonal acne.</div>
--	---	--

	<p>FABIOR (tazarotene) foam</p> <p>KLARON (sulfacetamide) suspension</p> <p>NEUAC (clindamycin/benzoyl peroxide/emollient) kit</p> <p>ONEXTON (clindamycin/benzoyl peroxide) gel, gel pump</p> <p>RETIN-A MICRO (tretinoin) (all products)</p> <p>ROSULA (sulfacetamide sodium/sulfur) cloths, wash</p> <p>SSS 10-5 (sulfacetamide sodium/sulfur) foam</p> <p>Sulfacetamide sodium cleanser, cleansing gel, lotion, shampoo, wash</p> <p>Sulfacetamide sodium/sulfur cleanser, cream, pad, suspension, wash</p> <p>SUMADAN/XLT (sulfacetamide sodium/sulfur) kit, wash</p> <p>SUMAXIN/ CP/TS (sulfacetamide sodium/sulfur) kit, pads, suspension, wash</p> <p>Tazarotene cream, foam, gel</p> <p>Tretinoin gel (all other manufacturers)</p> <p>Tretinoin microspheres (all products)</p> <p>WINLEVI (clascoterone) cream</p> <p>ZIANA (clindamycin/tretinoin) gel</p>	
--	--	--

Therapeutic Drug Class: ACNE AGENTS– ORAL ISOTRETINOIN – <i>Effective 7/1/2025</i>		
PA Required for all agents		Preferred products may be approved for adults and children ≥ 12 years of age for treating severe acne vulgaris or for treating moderate acne vulgaris in members unresponsive to conventional therapy. Non-preferred products may be approved for members meeting the following: <ul style="list-style-type: none">• Member has trialed/failed one preferred agent (failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) AND• Member is an adult or child ≥ 12 years of age with severe, recalcitrant nodulocystic acne and has been unresponsive to conventional therapy.
Preferred AMNESTEEM capsule CLARAVIS capsule Isotretinoin 10 mg, 20 mg, 30 mg, 40 mg capsule (<i>Mayne-Pharma, Upsher-Smith, Zydus only</i>) ZENATANE capsule	Non-Preferred ABSORICA capsule ABSORICA LD capsule Isotretinoin 10 mg, 20 mg, 30 mg, 40 mg capsule (<i>All manufacturers except Mayne-Pharma, Upsher-Smith, Zydus</i>) Isotretinoin 25 mg, 35 mg capsule MYORISAN capsule	
Therapeutic Drug Class: ANTI-PSORIATICS - Oral – <i>Effective 7/1/2025</i>		
No PA Required Acitretin capsule	PA Required Methoxsalen capsule	Prior authorization for non-preferred oral agents may be approved with failure of two preferred anti-psoriatic agents, one of which must be a preferred oral agent. Failure is defined as lack of efficacy of a 4-week trial, allergy, intolerable side effects or significant drug-drug interaction.
Therapeutic Drug Class: ANTI-PSORIATICS -Topical – <i>Effective 7/1/2025</i>		
No PA Required Calcipotriene cream, foam, ointment, solution Calcipotriene/betamethasone dipropionate ointment TACLONEX SCALP ^{BNR} (calcipotriene/betamethasone) suspension TACLONEX (calcipotriene/betamethasone) ointment	PA Required Calcipotriene/betamethasone dipropionate suspension Calcitriol ointment DUOBRII (halobetasol/tazarotene) lotion ENSTILAR (calcipotriene/betamethasone) foam SORILUX (calcipotriene) foam	Preferred and non-preferred products that contain a corticosteroid ingredient (such as betamethasone) will be limited to 4 weeks of therapy. Continued use will require one week of steroid-free time in between treatment periods. Non-preferred topical agents may be approved with failure of two preferred topical agents. If non-preferred topical agent being requested is a combination product, trial of two preferred agents must include a preferred combination agent. Failure is defined as lack of efficacy of a 4-week trial, allergy, intolerable side effects or significant drug-drug interaction. Members with >30% of their body surface area affected may not use Enstilar (calcipotriene/betamethasone DP) foam or Taclonex (calcipotriene/betamethasone DP) ointment products as safety and efficacy have not been established. ZORYVE (roflumilast) 0.3% cream may receive approval if meeting the following based on prescribed indication: <u>Plaque psoriasis</u> (0.3% cream formulation only): <ul style="list-style-type: none">• Member is ≥ 6 years of age AND• Member has a diagnosis of plaque psoriasis AND

	<p>VTAMA (tapinarof) cream</p> <p>ZORYVE 0.3% (roflumilast) cream, 0.3% foam</p>	<ul style="list-style-type: none">• Member has body surface area (BSA) involvement of $\leq 20\%$ AND• Member does not have moderate or severe hepatic impairment (Child-Pugh B or C) AND• Medication is being prescribed by or in consultation with a dermatologist AND• <u>If the affected area is limited to the scalp:</u><ul style="list-style-type: none">○ Prescriber attests that member has been counseled regarding alternative treatment options, including over-the-counter (OTC) emollients, vitamin D analogs, and coal tar shampoo when appropriate <p>AND</p> <ul style="list-style-type: none">○ Member has documented trial and failure (with a minimum 2-week treatment period) of a topical corticosteroid. Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction.• <u>If the affected area includes the face or body:</u><ul style="list-style-type: none">○ Member has documented trial and failure (with a minimum 2-week treatment period) of at least one product from ALL of the following categories. (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction):<ul style="list-style-type: none">▪ Topical corticosteroid▪ Topical calcineurin inhibitor (such as pimecrolimus, tacrolimus) <p>Members may not apply Zoryve (roflumilast) cream to $>20\%$ of affected body surface area, as safety and efficacy have not been established.</p> <p><u>Quantity limit:</u> 60 grams/30 days</p> <p><u>Initial approval:</u> 8 weeks</p> <p><u>Reauthorization:</u> Reauthorization for one year may be approved based on provider attestation that member’s symptoms improved during the initial 8 weeks of treatment and continuation of therapy is justified.</p> <p><u>Plaque psoriasis (0.3% foam formulation only):</u></p> <ul style="list-style-type: none">• Member is ≥ 12 years of age AND• Member has a diagnosis of plaque psoriasis AND• Member has body surface area (BSA) involvement of $\leq 20\%$ AND• Member does not have moderate or severe hepatic impairment (Child-Pugh B or C) AND• Medication is being prescribed by or in consultation with a dermatologist AND• If the affected area is limited to the scalp:<ul style="list-style-type: none">○ Prescriber attests that member has been counseled regarding alternative treatment options, including over-the-counter (OTC) emollients, vitamin D analogs, and coal tar shampoo when appropriate AND○ Member has documented trial and failure (with a minimum 2-week treatment period) of a topical corticosteroid. Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction.• If the affected area includes the face or body:<ul style="list-style-type: none">○ Member has documented trial and failure (with a minimum 2-week treatment period) of at least one product from ALL of the following categories (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction):<ul style="list-style-type: none">▪ Topical corticosteroid
--	--	---

		<div>▪ Topical calcineurin inhibitor (such as pimecrolimus, tacrolimus)</div> <div><u>Quantity limit:</u> 60 grams/30 days</div> <div><u>Initial approval:</u> 8 weeks</div> <div><u>Reauthorization:</u> Reauthorization for one year may be approved based on provider attestation that member’s symptoms improved during the initial 8 weeks of treatment and continuation of therapy is justified.</div>
Therapeutic Drug Class: IMMUNOMODULATORS, TOPICAL – <i>Effective 7/15/2025</i>		
Atopic Dermatitis		
<div>No PA Required (Unless indicated*)</div> <div>ELIDEL (pimecrolimus) cream</div> <div>*EUCRISA (crisaborole) ointment</div> <div>*OPZELURA (ruxolitinib) cream</div> <div>Pimecrolimus cream</div> <div>Tacrolimus ointment</div>	<div>PA Required</div> <div>ANZUPGO (delgocitinib) cream</div> <div>VTAMA (tapinarof) 1% cream</div> <div>ZORYVE (roflumilast) 0.15% cream, 0.3% foam</div>	<div>EUCRISA (crisaborole) may be approved if the following criteria are met:</div> <div><ul style="list-style-type: none">• Member is ≥ 3 months of age AND• Member has a diagnosis of mild to moderate atopic dermatitis AND• Member tried and failed‡ one preferred agent OR one medium-to-very high potency topical corticosteroid AND• Eucrisa (crisaborole) is being prescribed by or in consultation with a dermatologist or allergist/immunologist.</div> <div>OPZELURA (ruxolitinib) cream may be approved if the following criteria are met based on prescribed indication:</div> <div><u>Atopic Dermatitis</u><div><ul style="list-style-type: none">• Member is ≥ 2 years of age AND• Member has a diagnosis of mild to moderate atopic dermatitis AND• Medication is being prescribed by or in consultation with a dermatologist or allergist/immunologist AND• Member has trialed and failed‡ one preferred agent OR one medium potency to very high potency topical corticosteroid (such as mometasone furoate, betamethasone dipropionate, or fluocinonide) or prescriber verifies that member is not a candidate for topical corticosteroids.</div><div><u>Nonsegmental Vitiligo</u><div><ul style="list-style-type: none">• Member is ≥ 12 years of age AND• Member is immunocompetent AND• Member has a diagnosis of stable nonsegmental vitiligo, defined as no increase in the size of existing lesions and the absence of new lesions in the previous 3 to 6 months, AND• Medication is being prescribed by or in consultation with a dermatologist AND• Member has trialed and failed‡ one preferred agent AND one medium potency to very high potency topical corticosteroid (such as mometasone furoate, betamethasone dipropionate, or fluocinonide) or prescriber verifies that member is not a candidate for topical corticosteroids</div></div><div><u>Quantity limit:</u> 60 grams/week</div></div>

		<p>Non-preferred topical immunomodulator products may be approved for atopic dermatitis following adequate trial and failure‡ of one prescription topical corticosteroid AND two preferred agents.</p> <p>ZORYVE (roflumilast) 0.15% cream and 0.3% foam may receive approval if meeting the following based on prescribed indication:</p> <p><u>Atopic dermatitis (0.15% cream formulation only):</u></p> <ul style="list-style-type: none">• 6 years of age and older AND• Member has a diagnosis of mild atopic dermatitis in adult and pediatric patients AND• Request meets trial and failure criteria for non-preferred agents listed above <p><u>Seborrheic dermatitis (0.3% foam formulation only):</u></p> <ul style="list-style-type: none">• Member is \geq 9 years of age AND• Member has a diagnosis of seborrheic dermatitis AND• Member does not have moderate or severe hepatic impairment (Child-Pugh B or C) AND• Medication is being prescribed by or in consultation with a dermatologist AND• Member has been counseled that Zoryve foam is flammable. Fire, flame, or smoking during and immediately following application must be avoided.• <u>If the affected area is limited to the scalp:</u><ul style="list-style-type: none">○ Prescriber attests that member has been counseled regarding alternative treatment options, including over-the-counter (OTC) antifungal shampoo (such as selenium sulfide, zinc pyrithione) and OTC coal tar shampoo, when appropriate) <p>AND</p> <ul style="list-style-type: none">○ Member has documented trial and failure (with a minimum 2-week treatment period) of at least one prescription product for seborrheic dermatitis, such as ketoconazole 2% antifungal shampoo or a topical corticosteroid. Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction. • <u>If the affected area includes the face or body:</u><ul style="list-style-type: none">○ Member has documented trial and failure (with a minimum 2-week treatment period) with at least one product from ALL of the following categories (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction):<ul style="list-style-type: none">▪ Topical antifungal (such as ketoconazole, ciclopirox)▪ Topical corticosteroid▪ Topical calcineurin inhibitor (such as pimecrolimus, tacrolimus) <p><u>Quantity limit:</u> 60 grams/30 days</p> <p><u>Initial approval:</u> 8 weeks</p>
--	--	--

		<p>Members may not apply Zoryve (roflumilast) cream to >20% of affected body surface area, as safety and efficacy have not been established.</p> <p><u>Reauthorization</u>: Reauthorization for one year may be approved based on provider attestation that member’s symptoms improved during the initial 8 weeks of treatment and continuation of therapy is justified.</p> <p>‡Failure is defined as a lack of efficacy with a 2-week trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction</p>
Antineoplastic Agents		
<p>Preferred No PA Required (Unless indicated*)</p> <p>*Diclofenac 3% gel (generic Solaraze)</p> <p>Fluorouracil 5% cream (generic Efudex)</p> <p>Fluorouracil 2%, 5% solution</p>	<p>Non-Preferred PA Required</p> <p>Bexarotene gel</p> <p>CARAC (fluorouracil) cream</p> <p>EFUDEX (fluorouracil) cream</p> <p>Fluorouracil 0.5% (generic Carac) cream</p> <p>PANRETIN (alitretinoin) gel</p> <p>TARGRETIN (bexarotene) gel</p> <p>VALCHLOR (mechlorethamine) gel</p>	<p>*Diclofenac 3% gel (generic Solaraze) may be approved if the member has a diagnosis of actinic keratosis (AK).</p> <p>TARGRETIN (bexarotene) gel or VALCHLOR (mechlorethamine) gel may be approved for members who meet the following criteria:</p> <ul style="list-style-type: none">• Member is ≥ 18 years of age AND• Member has been diagnosed with Stage IA or IB cutaneous T-cell lymphoma (CTCL) AND• Member has refractory or persistent CTCL disease after other therapies OR has not tolerated other therapies AND• Member and partners have been counseled on appropriate use of contraception <p>Non-preferred agents may be approved for members who have failed an adequate trial of all preferred products FDA-approved for that indication. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.</p>
Other Agents		
<p>No PA Required</p> <p>Imiquimod (generic Aldara) cream</p> <p>Podofilox gel, solution</p>	<p>PA Required</p> <p>CONDYLOX (podofilox) gel</p> <p>HYFTOR (sirolimus) gel</p> <p>Imiquimod (generic Zyclara) cream, cream pump</p> <p>VEREGEN (sinecatechins) ointment</p> <p>ZYCLARA (imiquimod) cream, cream pump</p>	<p>Hyftor (sirolimus) gel</p> <ul style="list-style-type: none">• Member has a diagnosis of facial angiofibroma associated with tuberous sclerosis AND• Member is ≥ 6 years of age AND• Provider has evaluated, and member has received, all age-appropriate vaccinations as recommended by current immunization guidelines prior to initiating treatment with HYFTOR <p><u>Initial approval</u>: 6 months</p> <p><u>Reauthorization</u>: An additional 6 months may be approved based on provider attestation that symptoms improved during the initial 6 months of treatment and the provider has assessed use of all vaccinations recommended by current immunization guidelines.</p> <p><u>Maximum dose</u>: one 10-gram tube/28 days</p> <p>Veregen (sinecatechins) may be approved if the following criteria are met:</p> <ul style="list-style-type: none">• Member has a diagnosis of external genital and/or perianal warts (Condylomata acuminata) AND• Member is ≥ 18 years of age AND

		<p>Member is immunocompetent AND</p> <ul style="list-style-type: none">Member has tried and failed two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. <p>Zyclara (imiquimod) 2.5% cream may be approved if the following criteria are met:</p> <ul style="list-style-type: none">Member has a diagnosis of clinically typical visible or palpable actinic keratoses (AK) of the full face or balding scalp ANDMember is ≥ 18 years of age ANDMember is immunocompetent ANDMember has tried and failed one preferred product in the Antineoplastic Agents class (such as diclofenac gel or fluorouracil) AND the preferred imiquimod (generic Aldara) product. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. <p>Zyclara (imiquimod) 3.75% cream may be approved for:</p> <ul style="list-style-type: none">Treatment of clinically typical visible or palpable, actinic keratoses (AK) of the full face or balding scalp if the following criteria are met:<ul style="list-style-type: none">Member is ≥ 18 years of age ANDMember is immunocompetent ANDMember has tried and failed one preferred product from the Antineoplastic Agents class (such as diclofenac gel or fluorouracil) AND the preferred imiquimod (generic Aldara) product. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. <p>OR</p> <ul style="list-style-type: none">Treatment of external genital and/or perianal warts (Condylomata acuminata) if the following criteria are met:<ul style="list-style-type: none">Member is ≥ 12 years of age ANDMember has tried and failed two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. <p>All other non-preferred products may be approved for members who have trialed and failed all preferred products that are FDA-approved for use for the prescribed indication. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.</p> <p><u>Quantity Limits:</u> Aldara (imiquimod) cream has a quantity limit of 12 packets/28 days.</p>
Therapeutic Drug Class: ROSACEA AGENTS – <i>Effective 7/1/2025</i>		
<p>No PA Required</p> <p>Azelaic acid gel</p> <p>FINACEA (azelaic acid) gel</p> <p>FINACEA (azelaic acid) foam</p>	<p>PA Required</p> <p>Brimonidine gel pump</p> <p>*Doxycycline monohydrate DR capsule (generic Oracea)</p> <p>Ivermectin cream</p>	<p>Prior authorization for non-preferred products in this class may be approved if meeting the following criteria for the prescribed diagnosis:</p> <p><u>Rosacea:</u></p> <ul style="list-style-type: none">Member has a diagnosis of persistent (non-transient) facial erythema with inflammatory papules and pustules due to rosacea ANDPrescriber attests that medication is not being used solely for cosmetic purposes ANDMember has tried and failed two preferred agents of different mechanisms of action (Failure is defined as lack of efficacy with 4-week trial, allergy, contraindication, or intolerable side effects)

Metronidazole cream, lotion	Metronidazole 1% gel, gel pump	<u>Demodex Blepharitis:</u> <ul style="list-style-type: none">Requests for non-preferred topical ivermectin cream may be approved for treatment of moderate to severe Demodex blepharitis *Doxycycline monohydrate DR (generic Oracea) may be approved if the following criteria are met: <ul style="list-style-type: none">Member has taken generic doxycycline for a minimum of three months and failed therapy in the last 6 months. Failure is defined as: lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions ANDMember has history of an adequate trial/failure (8 weeks) of 2 other preferred agents (oral or topical). Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions ANDMember is ≥ 18 years of age and has been diagnosed with rosacea with inflammatory lesions (papules and pustules)
Metronidazole 0.75% gel	MIRVASO (Brimonidine gel pump)	
	NORITATE (metronidazole) cream	
	RHOFADE (oxymetazoline) cream	
	ROSADAN (metronidazole/skin cleanser) cream kit, gel kit	
Therapeutic Drug Class: TOPICAL STEROIDS – <i>Effective 7/1/2025</i>		
Low potency		
No PA Required DERMA-SMOOTH-FS (fluocinolone) 0.01% body oil/scalp oil ^{BNR} Desonide 0.05% cream, ointment Fluocinolone 0.01% cream, 0.01% solution Hydrocortisone (Rx) cream, lotion, ointment	PA Required Alclometasone 0.05% cream, ointment CAPEX (fluocinolone) 0.01% shampoo Desonide 0.05% lotion Fluocinolone 0.01% body oil, 0.01% scalp oil PROCTOCORT (hydrocortisone) (Rx) 1% cream SYNALAR (fluocinolone) 0.01% solution SYNALAR TS (fluocinolone/skin cleanser) Kit TEXACORT (hydrocortisone) 2.5% solution	Non-preferred Low Potency topical corticosteroids may be approved following adequate trial and failure of two preferred agents in the Low Potency class (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Medium potency		

No PA Required	PA Required	
Betamethasone dipropionate 0.05% cream, lotion, ointment	BESER (fluticasone) lotion, emollient kit	Non-preferred Medium Potency topical corticosteroids may be approved following adequate trial and failure of two preferred agents in the Medium Potency class (failure is defined as: lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Betamethasone valerate 0.1% cream, ointment	Betamethasone valerate 0.1% lotion, 0.12% foam	
Fluocinolone 0.025% cream, 0.05% cream, 0.005% ointment	Clocortolone 0.1% cream, cream pump	
Fluticasone cream, ointment	CLODERM (clocortolone) 0.1% cream, cream pump	
Hydrocortisone valerate 0.2% cream	CUTIVATE (fluticasone) 0.05% cream, lotion	
Mometasone 0.1% cream, 0.1% ointment, 0.1% solution	Diflorasone 0.05% cream	
Triamcinolone acetonide 0.025% cream, 0.1% cream, 0.025% ointment, 0.05% ointment, 0.1% ointment, 0.025% lotion, 0.1% lotion	Fluocinolone 0.025% ointment	
Triamcinolone 0.1% dental paste	Fluocinonide-E 0.05% cream	
	Flurandrenolide 0.05% cream, lotion, ointment	
	Fluticasone 0.05% lotion	
	Hydrocortisone butyrate 0.1% cream, lotion, solution, ointment, lipid/lipocream	
	Hydrocortisone valerate 0.2% ointment	
	KENALOG (triamcinolone) spray	
	LOCOID (hydrocortisone butyrate) 0.1% lotion	
	LOCOID LIPOCREAM (hydrocortisone butyrate-emollient) 0.1% cream	
	LUXIQ (betamethasone valerate) 0.12% foam	

	<div>ORALONE (Triamcinolone) 0.1% dental paste</div> <div>PANDEL (hydrocortisone probutate) 0.1% cream</div> <div>Prednicarbate 0.1% cream, ointment</div> <div>PSORCON (diflorasone) 0.05% cream</div> <div>SYNALAR (fluocinolone) 0.025% cream/kit, ointment/kit</div> <div>Triamcinolone 0.147 mg/gm spray</div>	
High potency		
<div>No PA Required (*unless exceeds duration of therapy)</div> <div>* Betamethasone dipropionate 0.05% ointment</div> <div>*Betamethasone dipropionate/propylene glycol (augmented) 0.05% cream</div> <div>*Fluocinonide 0.05% cream, 0.05% gel, 0.05% ointment, 0.05% solution</div> <div>*Triamcinolone acetonide 0.5% cream, 0.5% ointment</div>	<div>PA Required</div> <div>Amcinonide 0.1% cream, lotion</div> <div>APEXICON-E (diflorasone/emollient) 0.05% cream</div> <div>Desoximetasone 0.05%, 0.25% cream, 0.05% gel, 0.05%, 0.25% ointment</div> <div>Diflorasone 0.05% ointment</div> <div>Halcinonide 0.1% cream</div> <div>HALOG (halcinonide) 0.1% cream, ointment, solution</div> <div>TOPICORT (desoximetasone) 0.05%, 0.25% cream, 0.05% gel, 0.05%, 0.25% ointment</div>	<div>Non-preferred High Potency topical corticosteroids may be approved following adequate trial and failure of two preferred agents in the High Potency class (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).</div> <div>*All High Potency topical corticosteroids will require prior authorization beyond 4 weeks of therapy. The provider will be encouraged to transition to a medium or low potency topical steroid after this time has elapsed.</div> <div>Claims for compounded products containing high-potency topical steroids will be limited to a maximum of 60 grams or 60 mL of a high-potency ingredient per 4-week treatment period. Claims exceeding this quantity limit will require prior authorization with prescriber’s justification for use of the product at the prescribed dose.</div>
Very high potency		
<div>No PA Required (Unless exceeds duration of therapy*)</div> <div>*Betamethasone dipropionate/propylene glycol (augmented), 0.05% lotion 0.05% ointment</div>	<div>PA Required</div> <div>Betamethasone dipropionate/propylene glycol (augmented) 0.05% gel</div>	<div>Non-preferred Very High Potency topical corticosteroids may be approved following adequate trial and failure of clobetasol propionate in the same formulation as the product being requested (if the formulation of the requested non-preferred product is not available in preferred clobetasol product options, then trial and failure of any preferred clobetasol product formulation</div>

<p>*Clobetasol 0.05% cream, 0.05% gel, 0.05% ointment, 0.05% solution</p> <p>*Fluocinonide 0.1% cream</p>	<p>BRYHALI (halobetasol) 0.01% lotion</p> <p>Clobetasol emollient/emulsion 0.05% cream, foam</p> <p>Clobetasol 0.05% lotion, foam, spray, shampoo</p> <p>CLODAN (clobetasol) 0.05% cleanser kit</p> <p>Desoximetasone 0.25% spray</p> <p>DIPROLENE (betamethasone dipropionate/propylene glycol, augmented) 0.05% ointment</p> <p>Halobetasol 0.05% cream, foam, ointment</p> <p>IMPEKLO (clobetasol) 0.05% lotion</p> <p>LEXETTE (halobetasol) 0.05% foam</p> <p>OLUX (clobetasol) 0.05% foam</p> <p>TOPICORT (desoximetasone) 0.25% spray</p> <p>TOVET EMOLLIENT (clobetasol) 0.05% foam</p> <p>ULTRAVATE (halobetasol) 0.05% lotion</p> <p>VANOS (fluocinonide) 0.1% cream</p>	<p>will be required). Failure is defined as lack of efficacy with 2-week trial, allergy, intolerable side effects or significant drug-drug interactions.</p> <p>*All Very High Potency topical corticosteroids will require prior authorization beyond 2 weeks of therapy. If clobetasol propionate shampoo is being used to treat plaque psoriasis, then prior authorization will be required beyond 4 weeks of therapy. The provider will be encouraged to transition to a medium or low potency topical steroid after this time has elapsed.</p>
---	---	---

<h2>VI. Endocrine</h2>		
Therapeutic Drug Class: ANDROGENIC AGENTS, Topical, Injectable, Oral – <i>Effective 10/1/2025</i>		
PA Required for all agents in this class		
<p>Preferred</p> <p>Testosterone cypionate IM injection</p> <p>Testosterone gel packet</p> <p>Testosterone 1.62% gel pump</p> <p><i>Injectable testosterone cypionate is a pharmacy benefit when self-administered. Administration in an office setting is a medical benefit.</i></p>	<p>Non-Preferred</p> <p>ANDROGEL (testosterone) gel packet</p> <p>ANDROGEL (testosterone) gel 1.62% pump</p> <p>DEPO-TESTOSTERONE (testosterone cypionate) IM injection</p> <p>JATENZO (testosterone undecanoate) capsule</p> <p>KYZATREX (testosterone undecanoate) capsule</p> <p>METHITEST (methyltestosterone) tablet</p> <p>Methyltestosterone capsule</p> <p>NATESTO (testosterone) nasal spray</p> <p>TESTIM (testosterone) gel</p> <p>Testosterone 1% gel tube, 30 mg/1.5 ml pump</p> <p>Testosterone enanthate IM injection</p> <p>TLANDO (testosterone undecanoate) capsule</p> <p>UNDECATREX (testosterone undecanoate) capsule</p>	<p><u>Hypogonadotropic or Primary Hypogonadism (may be secondary to Klinefelter Syndrome):</u></p> <p>Preferred products may be approved for members meeting the following:</p> <ul style="list-style-type: none">Member is a male patient ≥ 16 years of age with a documented diagnosis of hypogonadotropic or primary hypogonadism OR ≥ 12 years of age with a diagnosis of hypogonadotropic or hypogonadism secondary to Klinefelter Syndrome (all other diagnoses will require manual review) ANDMember has two documented low serum testosterone levels below the lower limit of normal range for testing laboratory prior to initiation of therapy ANDMember does not have a diagnosis of breast or prostate cancer ANDIf the member is > 40 years of age, has prostate-specific antigen (PSA) < 4 ng/mL or has no palpable prostate nodule ANDMember has baseline hematocrit < 50% <p>Reauthorization Criteria (requests for renewal of a currently expiring prior authorization for a preferred product may be approved for members meeting the following criteria):</p> <ul style="list-style-type: none">Member is a male patient ≥ 16 years of age with a documented diagnosis of hypogonadotropic or primary hypogonadism OR ≥ 12 years of age with a diagnosis of hypogonadotropic or hypogonadism secondary to Klinefelter Syndrome ANDSerum testosterone is being regularly monitored (at least annually) to achieve total testosterone level in the middle tertile of the normal reference range ANDMember does not have a diagnosis of breast or prostate cancer ANDMember has a hematocrit < 54% <p><u>Gender Transition/Affirming Hormone Therapy:</u></p> <p>Preferred androgenic drugs may be approved for members meeting the following:</p> <ol style="list-style-type: none">Female sex assigned at birth and has reached Tanner stage 2 of puberty ANDIs undergoing female to male transition ANDHas a negative pregnancy test prior to initiation ANDHematocrit (or hemoglobin) is being monitored. <p>Non-Preferred Products:</p> <p>Non-preferred topical androgenic agents may be approved for members meeting the above criteria with trial and failed‡ therapy with two preferred topical androgen formulations.</p>

	XYOSTED (testosterone enanthate) SC injection	<p>Non-preferred injectable androgenic agents may be approved for members meeting the above criteria with trial and failed‡ therapy with a preferred injectable androgenic drug.</p> <p>Prior authorization for oral androgen agents (tablet, capsule, buccal) may be approved if member has trialed and failed‡ therapy with a preferred topical agent AND testosterone cypionate injection.</p> <p>‡Failure is defined as lack of efficacy with 8 week trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction.</p> <p>For all agents and diagnoses, members < 16 years of age will require a manual prior authorization review by a pharmacist (with exception of members ≥ 12 years of age with a diagnosis of hypogonadotropic or hypogonadism secondary to Klinefelter Syndrome).</p>
Therapeutic Drug Class: BONE RESORPTION SUPPRESSION AND RELATED AGENTS – <i>Effective 10/1/2025</i>		
Bisphosphonates		
No PA Required Alendronate solution, tablet Ibandronate tablet Risedronate tablet	PA Required ACTONEL (risedronate) tablet ATELVIA (risedronate) tablet BINOSTO (alendronate) effervescent tablet FOSAMAX (alendronate) tablet FOSAMAX plus D (alendronate/vit D) tablet	<p>Non-preferred bisphosphonates may be approved for members who have failed treatment with one preferred product at treatment dose. Failure is defined as lack of efficacy with a 12-month trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>For members who have a low risk of fracture, discontinuation of bisphosphonate therapy and drug holiday should be considered following 5 years of treatment. Low risk is defined as having a bone mineral density, based on the most recent T-score, of greater than (better than) -2.5 AND no history of low trauma or fragility fracture.</p>
Non-Bisphosphonates		
PREFERRED FORTEO (teriparatide) SC pen ^{BNR*} Raloxifene tablet	Non-Preferred BONSITY (teriparatide) SC pen Calcitonin salmon nasal spray EVISTA (raloxifene) tablet Teriparatide SC pen TYMLOS (abaloparatide) SC pen	<p>*FORTEO (teriparatide) or generic teriparatide may be approved if the following criteria are met:</p> <ul style="list-style-type: none">Member has one of the following diagnoses:<ul style="list-style-type: none">Male primary or hypogonadal osteoporosis (BMD T-score of -2.5 or less)Osteoporosis due to corticosteroid usePostmenopausal osteoporosis <p>AND</p> <ul style="list-style-type: none">Member is at very high risk for fracture† OR member has history of trial and failure of one preferred bisphosphonate. Failure is defined as lack of efficacy with a 12-month trial, allergy, intolerable side effects, or significant drug-drug interaction.Prior authorization will be given for one year and total exposure of parathyroid hormone analogs (teriparatide and abaloparatide) shall not exceed two years <p>TYMLOS (abaloparatide) may be approved if the member meets the following criteria:</p> <ul style="list-style-type: none">Member has a diagnosis of postmenopausal osteoporosis (BMD T-score of -2.5 or less) AND

		<ul style="list-style-type: none">Member is post-menopausal with very high risk for fracture† OR member has history of trial and failure of FORTEO (teriparatide). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. ANDPrior authorization will be given for one year and total exposure of parathyroid hormone analogs (teriparatide and abaloparatide) shall not exceed two years. <p>All other non-preferred non-bisphosphonates may be approved for FDA-labeled indications for members who have failed treatment with one preferred bisphosphonate or non-bisphosphonate product at treatment dose. Failure is defined as lack of efficacy with a 12-month trial, allergy, unable to use oral therapy, intolerable side effects, or significant drug-drug interaction.</p> <p>†Members at very high risk for fracture: Members will be considered at very high risk for fracture if they meet <u>one</u> of the following:</p> <ul style="list-style-type: none">A history of fracture within the past 12 months ORFractures experienced while receiving guideline-supported osteoporosis therapy ORA history of multiple fractures ORA history of fractures experienced while receiving medications that cause skeletal harm (such as long-term glucocorticoids) ORA very low T-score (less than -3.0) ORA high risk for falls or a history of injurious falls ORA very high fracture probability by FRAX (> 30% for a major osteoporosis fracture or > 4.5% for hip fracture). <table><tr><th>Non-Bisphosphonate Product</th><th>FDA-approved Maximum Dose</th></tr><tr><td>Calcitonin salmon nasal spray</td><td>1 metered dose spray (200 units) daily</td></tr><tr><td>Evista (raloxifene) oral tablet</td><td>60 mg daily</td></tr><tr><td>Forteo (teriparatide) subcutaneous injection</td><td>20 mcg daily</td></tr><tr><td>Tymlos (abaloparatide) subcutaneous injection</td><td>80 mcg daily</td></tr></table> <p><i>Note: Prior authorization criteria for Prolia (denosumab) and other injectable bone resorption and related agents are listed on Appendix P.</i></p>	Non-Bisphosphonate Product	FDA-approved Maximum Dose	Calcitonin salmon nasal spray	1 metered dose spray (200 units) daily	Evista (raloxifene) oral tablet	60 mg daily	Forteo (teriparatide) subcutaneous injection	20 mcg daily	Tymlos (abaloparatide) subcutaneous injection	80 mcg daily
Non-Bisphosphonate Product	FDA-approved Maximum Dose											
Calcitonin salmon nasal spray	1 metered dose spray (200 units) daily											
Evista (raloxifene) oral tablet	60 mg daily											
Forteo (teriparatide) subcutaneous injection	20 mcg daily											
Tymlos (abaloparatide) subcutaneous injection	80 mcg daily											
Therapeutic Drug Class: CONTRACEPTIVES - Topical – <i>Effective 07/10/2025</i>												
Effective 01/14/22, topical contraceptive patch products are eligible for coverage with a written prescription by an enrolled pharmacist. Additional information regarding pharmacist enrollment can be found at https://hcpf.colorado.gov/pharm-serv .												
No PA Required	PA Required											
ANNOVERA (segesterone acetate/EE) vaginal ring		Non-preferred topical contraceptive products may be approved following a trial and failure of one preferred topical										

Etonorgestrel/EE vaginal ring (<i>Prasco Labs</i>)	Etonorgestrel/EE vaginal ring (<i>all other manufacturers</i>)	contraceptive product. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.
*PHEXXI (lactic acid/citric/potassium) vaginal gel	Norelgestromin/EE TD patch (generic XULANE)	* PHEXXI (lactic acid/citric/potassium) vaginal gel quantity limit: 120 grams per 30 days
TWIRLA (levonorgestrel/EE) TD patch	NUVARING (etonorgestrel/EE) vaginal ring	Effective 7/1/2022: Prescriptions are eligible to be filled for up to a twelve-month supply.
XULANE (norelgestromin/EE) TD patch	ZAFEMY (norelgestromin/EE) TD patch	<i>Note: IUD and select depot product formulations are billed through the medical benefit</i>
Therapeutic Drug Class: DIABETES MANAGEMENT CLASSES, INSULINS – Effective 02/27/2025		
Rapid-Acting		
No PA Required	PA Required	All non-preferred products may be approved following trial and failure of treatment with two preferred products, one of which is the same rapid-acting insulin analog (lispro or aspart) as the non-preferred product being requested. (Failure is defined as allergy [hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, and angioedema] or intolerable side effects).
HUMALOG (insulin lispro) cartridge, vial	ADMELOG (insulin lispro) Solostar pen, vial	Afrezza (human insulin) may be approved if meeting the following criteria: <ul style="list-style-type: none">• Member is 18 years or older AND• Member has trialed and failed treatment with two preferred products (failure is defined as allergy [hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, or angioedema] or intolerable side effects) AND• Member must not have chronic lung disease such as COPD or asthma AND• If member has type 1 diabetes, must use in conjunction with long-acting insulin AND• Prescriber acknowledges that Afrezza is not recommended in patients who smoke or have recently stopped smoking.
Insulin aspart cartridge, pen, vial	AFREZZA (regular insulin) cartridge, unit	
Insulin lispro Kwikpen, Jr. Kwikpen, vial (<i>Eli Lilly</i>)	APIDRA (insulin glulisine) Solostar pen, vial	
NOVOLOG (insulin aspart) cartridge, FlexPen, vial	FIASP (insulin aspart) FlexPen, PenFill, pump cartridge, vial	
	HUMALOG (insulin lispro) 200 U/mL Kwikpen	
	HUMALOG Tempo Pen 100 U/mL	
	HUMALOG 100U/mL KwikPen, vial	
	HUMALOG Jr. (insulin lispro) KwikPen	
	Insulin lispro 100 U/mL vial (<i>all other manufacturers</i>)	
	KIRSTY (insulin aspart-xjhz) Kwikpen, vial, Tempo pen	

	LYUMJEV (insulin lispro-aabc) Kwikpen, vial, Tempo pen MERILOG (insulin aspart-szjj) pen, vial	
Short-Acting		
No PA Required HUMULIN R U-100 (insulin regular) vial (OTC) NOVOLIN R U-100 (insulin regular) FlexPen (OTC)	PA Required NOVOLIN R U-100 (insulin regular) vial (OTC)	Non-preferred products may be approved following trial and failure of treatment with one preferred product (failure is defined as allergy or intolerable side effects).
Intermediate-Acting		
No PA Required HUMULIN N U-100 (insulin NPH), KwikPen (OTC), vial (OTC) NOVOLIN N U-100 (insulin NPH) FlexPen (OTC)	PA Required NOVOLIN N U-100 (insulin NPH) vial (OTC)	Non-preferred products may be approved following trial and failure of treatment with one preferred product (failure is defined as allergy or intolerable side effects).
Long-Acting		
Preferred LANTUS ^{BNR} (insulin glargine) Solostar, vial TRESIBA ^{BNR} (insulin degludec)* FlexTouch, vial	Non-Preferred BASAGLAR (insulin glargine) Kwikpen, Tempo pen Insulin degludec FlexTouch, vial Insulin glargine solostar, vial Insulin glargine MAX solostar Insulin glargine-yfgn pen, vial LEVEMIR (insulin detemir) FlexTouch, vial REZVOGLAR (insulin glargine-aglr) Kwikpen SEMGLEE (insulin glargine-yfgn) pen, vial	*Preferred Tresiba pen and vial formulations may be approved for members who have trialed and failed‡ Lantus. Non-preferred products may be approved if the member has tried and failed‡ treatment with Lantus AND a preferred insulin degludec product. ‡Failure is defined as lack of efficacy, allergy, or intolerable side effects.

	<div>TOUJEO (insulin glargine) Solostar</div> <div>TOUJEO MAX (insulin glargine) Solostar</div>	
Concentrated		
<div>No PA Required</div> <div>HUMULIN R U-500 (insulin regular) concentrated vial, Kwikpen</div>	<div>PA Required</div>	<div>Non-preferred products may be approved following trial and failure of treatment with one preferred product (failure is defined as allergy or intolerable side effects).</div>
Mixtures		
<div>No PA Required</div> <div>HUMALOG MIX 50/50 Kwikpen</div> <div>HUMALOG MIX 75/25 vial</div> <div>HUMULIN 70/30 (OTC) Kwikpen, vial</div> <div>Insulin aspart protamine/insulin aspart 70/30 FlexPen, vial (generic Novolog Mix)</div> <div>Insulin lispro protamine/insulin lispro 75/25 Kwikpen (generic Humalog Mix)</div> <div>NOVOLOG MIX 70/30 FlexPen, vial</div>	<div>PA Required</div> <div>HUMALOG MIX 75/25 Kwikpen</div> <div>NOVOLIN 70/30 FlexPen, vial (OTC)</div>	<div>Non-preferred products may be approved if the member has failed treatment with two of the preferred products (failure is defined as: allergy or intolerable side effects).</div>
Therapeutic Drug Class: DIABETES MANAGEMENT CLASSES, NON- INSULINS – 1/1/2026		
Amylin		
	<div>PA Required</div> <div>SYMLIN (pramlintide) pen</div>	<div>SYMLIN (pramlintide) may be approved following trial and failure of metformin AND trial and failure of a DPP-4 inhibitor or GLP-1 analogue. Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen) following 3-month trial, allergy, intolerable side effects, or a significant drug-drug interaction. Prior authorization may be approved for Symlin (pramlintide) products for members with a diagnosis of Type 1 diabetes without requiring trial and failure of other products.</div> <div>Maximum Dose: Prior authorization will be required for doses exceeding FDA-approved dosing listed in product package labeling.</div>
Biguanides		
<div>No PA Required</div> <div>Metformin IR tablets</div>	<div>PA Required</div> <div>GLUMETZA ER (metformin) tablet</div>	<div>Non-preferred products may be approved for members who have failed treatment with two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.</div>

Metformin ER 500mg, 750mg tablets (generic Glucophage XR)	Metformin 625 mg tablets	Liquid metformin may be approved for members that are unable to use a solid oral dosage form.														
	Metformin ER (generic Fortamet, Glumetza, <i>Bayshore Pharma</i>)															
	Metformin solution (generic Riomet)															
	RIOMET (metformin) solution															
	RIOMET ER (metformin) suspension															
Dipeptidyl Peptidase 4 Enzyme inhibitors (DPP-4 Inhibitors)																
Preferred	Non-Preferred PA Required	<p>Non-preferred DPP-4 inhibitors may be approved after a member has failed a 3-month trial of one preferred product. Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen), allergy, contraindication, intolerable side effects, or a significant drug-drug interaction.</p> <p><u>Continuation of therapy:</u> Members currently stabilized on Januvia (sitagliptin) may receive approval for continuation of therapy with that agent.</p> <p><u>Maximum Dose:</u> Prior authorization will be required for doses exceeding the FDA-approved maximum dosing listed in the following table:</p> <table><tr><th>DPP-4 Inhibitor</th><th>FDA-Approved Maximum Daily Dose</th></tr><tr><td>Alogliptin (generic Nesina)</td><td>25 mg/day</td></tr><tr><td>Januvia (sitagliptin)</td><td>100 mg/day</td></tr><tr><td>Nesina (alogliptin)</td><td>25 mg/day</td></tr><tr><td>Onglyza (saxagliptin)</td><td>5 mg/day</td></tr><tr><td>Tradjenta (linagliptin)</td><td>5 mg/day</td></tr><tr><td>Zituvio (sitagliptin)</td><td>100 mg/day</td></tr></table>	DPP-4 Inhibitor	FDA-Approved Maximum Daily Dose	Alogliptin (generic Nesina)	25 mg/day	Januvia (sitagliptin)	100 mg/day	Nesina (alogliptin)	25 mg/day	Onglyza (saxagliptin)	5 mg/day	Tradjenta (linagliptin)	5 mg/day	Zituvio (sitagliptin)	100 mg/day
DPP-4 Inhibitor	FDA-Approved Maximum Daily Dose															
Alogliptin (generic Nesina)	25 mg/day															
Januvia (sitagliptin)	100 mg/day															
Nesina (alogliptin)	25 mg/day															
Onglyza (saxagliptin)	5 mg/day															
Tradjenta (linagliptin)	5 mg/day															
Zituvio (sitagliptin)	100 mg/day															
TRADJENTA (linagliptin) tablet	Alogliptin tablet															
	BRYNOVIN (Sitagliptin) tablet for suspension															
	JANUVIA (sitagliptin) tablet															
	NESINA (alogliptin) tablet															
	ONGLYZA (saxagliptin) tablet															
	Saxagliptin tablet															
	Sitagliptin (generic Zituvio)															
	ZITUVIO (sitagliptin tablet)															
DPP-4 Inhibitors – Combination with Metformin																
Preferred	Non-Preferred PA Required															

JENTADUETO ^{BNR} (linagliptin/metformin) tablet		Alogliptin/metformin tablet	<p>Non-preferred combination products may be approved for members who have been stable on the two individual ingredients of the requested combination for three months AND have had adequate three-month trial and failure of a preferred combination agent. Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen), contraindication, allergy, intolerable side effects, or a significant drug-drug interaction.</p> <p><u>Continuation of therapy:</u> Members currently stabilized on Janumet (sitagliptin/metformin) or Janumet XR (sitagliptin ER/metformin ER) may receive approval for continuation of therapy with those agents.</p> <p><u>Maximum Dose:</u> Prior authorization will be required for doses exceeding the FDA-approved maximum dosing listed in the following table:</p> <table><tr><th>DPP-4 Inhibitor Combination</th><th>FDA Approved Maximum Daily Dose</th></tr><tr><td>Alogliptin/metformin tablet</td><td>25 mg alogliptin/2,000 mg metformin</td></tr><tr><td>Janumet and Janumet XR (sitagliptin/metformin)</td><td>100 mg sitagliptin/2,000 mg of metformin</td></tr><tr><td>Jentadueto and Jentadueto XR (linagliptin/metformin)</td><td>5 mg linagliptin/2,000 mg metformin</td></tr><tr><td>Kazano (alogliptin/metformin)</td><td>25 mg alogliptin/ 2,000 mg metformin</td></tr><tr><td>Kombiglyze XR (saxagliptin ER/metformin ER) tablet</td><td>5 mg saxagliptin/2,000 mg metformin</td></tr></table>	DPP-4 Inhibitor Combination	FDA Approved Maximum Daily Dose	Alogliptin/metformin tablet	25 mg alogliptin/2,000 mg metformin	Janumet and Janumet XR (sitagliptin/metformin)	100 mg sitagliptin/2,000 mg of metformin	Jentadueto and Jentadueto XR (linagliptin/metformin)	5 mg linagliptin/2,000 mg metformin	Kazano (alogliptin/metformin)	25 mg alogliptin/ 2,000 mg metformin	Kombiglyze XR (saxagliptin ER/metformin ER) tablet	5 mg saxagliptin/2,000 mg metformin
DPP-4 Inhibitor Combination	FDA Approved Maximum Daily Dose														
Alogliptin/metformin tablet	25 mg alogliptin/2,000 mg metformin														
Janumet and Janumet XR (sitagliptin/metformin)	100 mg sitagliptin/2,000 mg of metformin														
Jentadueto and Jentadueto XR (linagliptin/metformin)	5 mg linagliptin/2,000 mg metformin														
Kazano (alogliptin/metformin)	25 mg alogliptin/ 2,000 mg metformin														
Kombiglyze XR (saxagliptin ER/metformin ER) tablet	5 mg saxagliptin/2,000 mg metformin														
JENTADUETO XR (linagliptin/metformin) tablet		JANUMET (sitagliptin/metformin) tablet													
		JANUMET XR (sitagliptin/metformin) tablet													
		KAZANO (alogliptin/metformin) tablet													
		KOMBIGLYZE XR (saxagliptin/metformin) tablet													
		Linagliptin/metformin tablet													
		Saxagliptin/metformin tablet													
		Sitagliptin/metformin (generic Zituvimet)													

Glucagon-like Peptide-1 Receptor Agonists (GLP-1 Analogues)		
<p>Preferred *Must meet eligibility criteria</p> <p>*Liraglutide pen (<i>Teva</i>)</p> <p>*OZEMPIC (semaglutide) pen</p> <p>*TRULICITY (dulaglutide) pen</p> <p>*VICTOZA (liraglutide) pen</p> <p>**WEGOVY (Semaglutide) pen</p>	<p>Non-Preferred PA Required</p> <p>Exenatide pen</p> <p>Liraglutide pen (<i>all other manufacturers</i>)</p> <p>MOUNJARO (tirzepatide) pen</p>	<p>*Preferred products may be approved for members with a diagnosis of type 2 diabetes.</p> <p>**WEGOVY (semaglutide) may be approved if meeting the following criteria:</p> <ul style="list-style-type: none">Member is 18 years of age or older ANDMember has established cardiovascular disease (history of myocardial infarction, stroke, or symptomatic peripheral arterial disease) and either obesity or overweight (defined as a BMI ≥25 kg/m²) ANDMember does not have a diagnosis of Type 2 diabetes ANDWegovy (semaglutide) is being prescribed to decrease the risk of adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) ANDMember has been counseled regarding implementation of lifestyle interventions (diet modification and exercise) to promote weight loss. <p>ZEPBOUND (tirzepatide) may be approved if meeting the following criteria:</p> <ul style="list-style-type: none">Member is 18 years of age or older ANDMember has a documented diagnosis of moderate to severe obstructive sleep apnea (OSA) AND

	<div><div><div>RYBELSUS (semaglutide) oral tablet</div><div>ZEPBOUND (tirzepatide)</div></div></div>	<div><ul style="list-style-type: none">• Member has a BMI ≥30 kg/m² indicating obesity documented in medical chart notes AND• Diagnosis of OSA is confirmed by a sleep test that is approved by the Food and Drug Administration (FDA) as a diagnostic device AND• A polysomnogram has been performed at baseline with a documented result of Apnea-Hypopnea Index (AHI) ≥15 events/hour (submission of sleep study documentation required) AND• Member is not pregnant or planning to become pregnant AND• Member has been counseled regarding the risk of medullary thyroid cancer (MTC) with the use of Zepbound (tirzepatide) and does not have a personal or family history of MTC or Multiple Endocrine Neoplasia syndrome type 2 (MEN 2) AND• The requested medication is being prescribed by or in consultation with a neurologist, pulmonologist, otolaryngologist, or other sleep medicine specialist AND• Member has been counseled regarding and is engaged in implementation of lifestyle interventions (diet modification and exercise) to promote weight loss AND• Member has failed a 6-month trial of continuous positive airway pressure (CPAP) or has a contraindication to the use of PAP therapy.<div><u>Reauthorization:</u> Reauthorization for one year may be approved if meeting the following criteria:<ul style="list-style-type: none">• Member has previous PA approval on file (requests for members that do not have a historic PA approval on file will be subject to meeting “Initial Authorization” criteria listed above) AND• Prescriber attests that an in-person clinical re-evaluation of OSA from baseline has been performed by the treating practitioner AND• Clinical improvement in OSA symptoms has been documented in clinical chart notes AND• Adherence to use of Zepbound (tirzepatide) regimen has been evaluated by the treating practitioner.</div><div>Requests for GLP-1 analogues that are FDA-indicated for the treatment of metabolic dysfunction-associated steatohepatitis (MASH) may be approved if meeting the following:<ul style="list-style-type: none">• Member has a diagnosis of MASH with stage F2 to F3 fibrosis that has been confirmed by clinical presentation along with liver biopsy or imaging results AND• Member meets the FDA-labeled minimum age requirement for the prescribed product AND• Member does not have cirrhosis or significant liver disease other than MASH AND• The requested medication is being prescribed for use for the FDA-labeled indication and as outlined in product package labeling AND• Medication is prescribed by or in consultation with a gastroenterologist, endocrinologist, obesity medicine specialist, hepatologist, or liver transplant provider AND• Requests for non-preferred agents will be subject to meeting non-preferred criteria listed below.</div><div>All other non-preferred products may be approved for members with an FDA-labeled diagnosis (excluding labeled use solely for weight loss) following a trial and failure‡ of three preferred agents that are FDA-labeled for use for the prescribed indication</div><div><u>Continuation of therapy:</u> Members that are currently stabilized on therapy with Mounjaro (tirzepatide) 7.5 mg, 10 mg, 12.5 mg, or 15 mg strengths may receive approval for continuation of therapy with that product strength.</div><div><u>Maximum Dose:</u> Prior authorization is required for all products exceeding maximum dose listed in product package labeling.</div><table><tr><th colspan="2">Table 1: GLP-1 Analogue Maximum Dose</th></tr><tr><td>Mounjaro (tirzepatide)</td><td>15 mg weekly</td></tr></table></div>	Table 1: GLP-1 Analogue Maximum Dose		Mounjaro (tirzepatide)	15 mg weekly
Table 1: GLP-1 Analogue Maximum Dose						
Mounjaro (tirzepatide)	15 mg weekly					

			Ozempic (semaglutide)	2 mg weekly	
			Rybelsus (semaglutide)	14 mg daily	
			Trulicity (dulaglutide)	4.5 mg weekly	
			Victoza (liraglutide)	1.8 mg daily	
			Wegovy (semaglutide)	2.4 mg weekly	
			‡Failure is defined as lack of efficacy with a 3-month trial (such as not meeting hemoglobin A1C goal despite adherence to regimen), allergy, intolerable side effects, limited dexterity resulting in the inability to administer doses of a preferred product, or a significant drug-drug interaction.		
<i>Note: Prior Authorization for GLP-1 analogues prescribed solely for weight loss will not be approved.</i>					
Other Hypoglycemic Combinations					
	PA Required	Non-preferred products may be approved for members who have been stable on each of the individual ingredients in the requested combination for 3 months (including cases where the ingredients are taken as two separate 3-month trials or when taken in combination for at least 3 months). SOLIQUA (insulin glargine/lixisenatide) may be approved if member has had a trial and failure with one preferred GLP-1 AND one preferred insulin glargine product (Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen), allergy, intolerable side effects, or significant drug-drug interaction.)			
	Alogliptin/pioglitazone tablet				
	Glipizide/metformin tablet				
	Glyburide/metformin tablet				
	GLYXAMBI (empagliflozin/linagliptin) tablet				
	OSENI (alogliptin/pioglitazone) tablet				
	Pioglitazone/glimepiride tablet				
	QTERN (dapagliflozin/saxagliptin) tablet				
	SOLIQUA (insulin glargine/lixisenatide) pen				
	STEGLUJAN (ertugliflozin/sitagliptin) tablet				
	TRIJARDY XR tablet(empagliflozin/linagliptin/metformin)				

	XULTOPHY (insulin degludec/liraglutide) pen	
Meglitinides		
Repaglinide tablet	PA Required Nateglinide tablet	Non-preferred products may be approved for members who have failed treatment with one preferred product. Failure is defined as: lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen), allergy, intolerable side effects, or significant drug-drug interaction.
Meglitinides Combination with Metformin		
	PA Required Repaglinide/metformin	Non-preferred products may be approved for members who have been stable on the two individual ingredients of the requested combination for 3 months.
Sodium-Glucose Cotransporter Inhibitors (SGLT inhibitors)		
No PA Required FARXIGA ^{BNR} (dapagliflozin) tablet	PA Required Dapagliflozin tablet INPEFA (sotagliflozin) tablet INVOKANA (canagliflozin) tablet JARDIANCE (empagliflozin) tablet STEGLATRO (ertugliflozin) tablet	Non-preferred products may receive approval following trial and failure with one preferred product. Failure is defined as lack of efficacy with 3-month trial (such as not meeting hemoglobin A1C goal despite adherence to regimen), contraindication, allergy, intolerable side effects, or a significant drug-drug interaction. <u>Maximum Dose:</u> Prior authorization is required for all products exceeding maximum dose listed in product package labeling.
SGLT Inhibitor Combinations with Metformin		
No PA Required XIGDUO XR ^{BNR} (dapagliflozin/metformin) tablet	PA Required Dapagliflozin/Metformin XR tablet INVOKAMET (canagliflozin/metformin) tablet INVOKAMET XR (canagliflozin/metformin) tablet SEGLUROMET (ertugliflozin/metformin) tablet SYNJARDY (empagliflozin/metformin) tablet	Non-preferred products may be approved for members who have been stable on the two individual ingredients of the requested combination for 3 months. INVOKAMET, INVOKAMET XR, SEGLUROMET, SYNJARDY, SYNJARDY XR and XIGDUO XR are contraindicated in patients with an eGFR less than 30 mL/min/1.73 m ² or on dialysis.

	SYNJARDY XR (empagliflozin/metformin) tablet																					
Thiazolidinediones (TZDs)																						
No PA Required Pioglitazone tablet	PA Required ACTOS (pioglitazone) tablet	Non-preferred agents may be approved following trial and failure of one preferred product. Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen) with a 3-month trial, allergy, intolerable side effects, or a significant drug-drug interaction.																				
Thiazolidinediones Combination with Metformin																						
Pioglitazone/metformin tablet	PA Required ACTOPLUS MET (pioglitazone/metformin) TABLET	Non-preferred products may be approved for members who have been stable on the two individual ingredients of the requested combination for 3 months.																				
Therapeutic Drug Class: ESTROGEN AGENTS -Effective 10/1/2025																						
No PA Required	PA Required	Non-preferred parenteral estrogen agents may be approved with trial and failure of one preferred parenteral agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.																				
Parenteral																						
DELESTROGEN 10mg ^{BNR} (estradiol valerate) vial DELESTROGEN 20mg, 40mg (estradiol valerate) vial DEPO-ESTRODIOL (estradiol cypionate) vial Estradiol valerate 40mg/mL vial, 20mg/mL vial	Estradiol valerate 10mg/mL vial	Non-preferred oral estrogen agents may be approved with trial and failure of one preferred oral agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. Non-preferred transdermal estrogen agents may be approved with trial and failure of two preferred transdermal agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.																				
Oral/Transdermal		<table><tr><th colspan="2">Table 1: Transdermal Estrogen FDA-Labeled Dosing</th></tr><tr><td>ALORA (estradiol) patch</td><td>2/week</td></tr><tr><td>CLIMARA (estradiol) patch</td><td>1/week</td></tr><tr><td>DOTTI (estradiol) patch</td><td>2/week</td></tr><tr><td>Estradiol patch (once weekly)</td><td>1/week</td></tr><tr><td>Estradiol patch (twice weekly)</td><td>2/week</td></tr><tr><td>LYLLANA (estradiol) patch</td><td>2/week</td></tr><tr><td>MENOSTAR (estradiol) patch</td><td>1/week</td></tr><tr><td>MINIVELLE (estradiol) patch</td><td>2/week</td></tr><tr><td>VIVELLE-DOT (estradiol) patch</td><td>2/week</td></tr></table>	Table 1: Transdermal Estrogen FDA-Labeled Dosing		ALORA (estradiol) patch	2/week	CLIMARA (estradiol) patch	1/week	DOTTI (estradiol) patch	2/week	Estradiol patch (once weekly)	1/week	Estradiol patch (twice weekly)	2/week	LYLLANA (estradiol) patch	2/week	MENOSTAR (estradiol) patch	1/week	MINIVELLE (estradiol) patch	2/week	VIVELLE-DOT (estradiol) patch	2/week
Table 1: Transdermal Estrogen FDA-Labeled Dosing																						
ALORA (estradiol) patch	2/week																					
CLIMARA (estradiol) patch	1/week																					
DOTTI (estradiol) patch	2/week																					
Estradiol patch (once weekly)	1/week																					
Estradiol patch (twice weekly)	2/week																					
LYLLANA (estradiol) patch	2/week																					
MENOSTAR (estradiol) patch	1/week																					
MINIVELLE (estradiol) patch	2/week																					
VIVELLE-DOT (estradiol) patch	2/week																					
Estradiol oral tablet	CLIMARA (estradiol) patch																					
Estradiol (generic Climara) weekly patch	DOTTI (estradiol) patch																					
MINIVELLE ^{BNR} (estradiol) patch	ESTRACE (estradiol) oral tablet																					
VIVELLE-DOT ^{BNR} (estradiol) patch	Estradiol bi-weekly patch																					
	LYLLANA (estradiol) patch																					
	MENOSTAR (estradiol) patch																					

		<i>Note: Estrogen agents are a covered benefit for gender affirming hormone therapy and treating clinicians and mental health providers should be knowledgeable about the diagnostic criteria for gender-affirming hormone treatment and have sufficient training and experience in assessing related mental health conditions.</i>
Therapeutic Drug Class: GLUCAGON, SELF-ADMINISTERED – <i>Effective 11/8/2024</i>		
Preferred No PA Required BAQSIMI (glucagon) nasal spray Glucagon Emergency Kit (<i>Eli Lilly, Fresenius, Amphastar</i>) ZEGALOGUE (dasiglucagon) autoinjector	Non-Preferred PA Required GVOKE (glucagon) Hypopen, Syringe, vial ZEGALOGUE (dasiglucagon) syringe	Non-preferred products may be approved if the member has failed treatment with two preferred products (failure is defined as allergy to ingredients in product, intolerable side effects, contraindication, or inability to administer dosage form). Quantity limit for all products: 2 doses per year unless used/ damaged/ lost
Therapeutic Drug Class: GROWTH HORMONES – <i>Effective 10/1/2025</i>		
Preferred GENOTROPIN (somatropin) cartridge, Miniquick pen NGENLA (Somatrogon-ghla)* pen NORDITROPIN (somatropin) Flexpro pen SKYTROFA (lonapegsomatropin-tegd)* cartridge	Non-Preferred HUMATROPE (somatropin) cartridge NUTROPIN AQ (somatropin) Nuspín injector OMNITROPE (somatropin) cartridge, vial SAIZEN (somatropin) cartridge, vial SEROSTIM (somatropin) vial SOGROYA (somapacitan-beco) pen ZOMACTON (somatropin) vial	<p>All preferred products may be approved if the member has one of the qualifying diagnoses listed below (diagnosis may be verified through AutoPA) AND if prescription does not exceed limitations for maximum dosing (Table 1).</p> <p>*Second line preferred products (NGENLA, SKYTROFA) require trial and failure of Genotropin (somatropin) OR Norditropin (somatropin).</p> <p>Non-preferred Growth Hormone products may be approved if the following criteria are met:</p> <ul style="list-style-type: none">• Member failed treatment with one preferred growth hormone product (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions) AND• Member has a qualifying diagnosis that includes any of the following conditions:<ul style="list-style-type: none">▪ Prader-Willi Syndrome (PWS)▪ Chronic renal insufficiency/failure requiring transplantation (defined as Creatinine Clearance < 30mL/min)▪ Turner’s Syndrome▪ Hypopituitarism: as a result of pituitary disease, hypothalamic disease, surgery, radiation therapy or trauma verified by one of the following:<ul style="list-style-type: none">○ Has failed at least one GH stimulation test (peak GH level < 10 ng/mL)○ Has at least one documented low IGF-1 level (below normal range for patient’s age – refer to range on submitted lab document)○ Has deficiencies in ≥ 3 pituitary axes (such as TSH, LH, FSH, ACTH, ADH)▪ Cachexia associated with AIDS▪ Noonan Syndrome▪ Short bowel syndrome▪ Neonatal symptomatic growth hormone deficiency (limited to 3-month PA approval) <p>AND</p> <ul style="list-style-type: none">• Prescription does not exceed limitations for FDA-labeled maximum dosing for prescribed indication (Table 1) based on prescriber submission/verification of patient weight from most recent clinical documentation
Table 1: Growth Hormone Product Maximum Dosing*		

		<table><tr><th>Medication</th><th>Pediatric Maximum Dosing per week (age < 18 years)</th><th>Adult Maximum Dosing per week (age ≥ 18 years)</th></tr><tr><td>Genotropin</td><td>0.48 mg/kg/week</td><td>0.08 mg/kg/week</td></tr><tr><td>Humatrope</td><td>0.47 mg/kg/week</td><td>0.0875 mg/kg/week</td></tr><tr><td>Ngenla</td><td>0.66 mg/kg/week</td><td>Not Indicated</td></tr><tr><td>Norditropin Flexpro</td><td>0.47 mg/kg/week</td><td>0.112 mg/kg/week</td></tr><tr><td>Nutropin AQ Nuspin</td><td>0.7 mg/kg/week</td><td>0.175 mg/kg/week for ≤35 years of age 0.0875 mg/kg/week for >35 years of age</td></tr><tr><td>Omnitrope</td><td>0.48 mg/kg/week</td><td>0.08 mg/kg/week</td></tr><tr><td>Saizen</td><td>0.18 mg/kg/week</td><td>0.07 mg/kg/week</td></tr><tr><td>Serostim</td><td>Not Indicated</td><td>42 mg/week for HIV wasting or cachexia (in combination with antiretroviral therapy)</td></tr><tr><td>Skytrofa</td><td>1.68 mg/kg/week</td><td>Not Indicated</td></tr><tr><td>Sogroya</td><td>Dose Individualized for each patient, based on growth response</td><td>8 mg/week</td></tr><tr><td>Zomacton</td><td>0.47 mg/kg/week</td><td>0.0875 mg/kg/week</td></tr><tr><td>Zorbtive</td><td>Not Indicated</td><td>56 mg/week for up to 4 weeks for short bowel syndrome only</td></tr></table> <p>*Based on FDA labeled indications and dosing</p>	Medication	Pediatric Maximum Dosing per week (age < 18 years)	Adult Maximum Dosing per week (age ≥ 18 years)	Genotropin	0.48 mg/kg/week	0.08 mg/kg/week	Humatrope	0.47 mg/kg/week	0.0875 mg/kg/week	Ngenla	0.66 mg/kg/week	Not Indicated	Norditropin Flexpro	0.47 mg/kg/week	0.112 mg/kg/week	Nutropin AQ Nuspin	0.7 mg/kg/week	0.175 mg/kg/week for ≤35 years of age 0.0875 mg/kg/week for >35 years of age	Omnitrope	0.48 mg/kg/week	0.08 mg/kg/week	Saizen	0.18 mg/kg/week	0.07 mg/kg/week	Serostim	Not Indicated	42 mg/week for HIV wasting or cachexia (in combination with antiretroviral therapy)	Skytrofa	1.68 mg/kg/week	Not Indicated	Sogroya	Dose Individualized for each patient, based on growth response	8 mg/week	Zomacton	0.47 mg/kg/week	0.0875 mg/kg/week	Zorbtive	Not Indicated	56 mg/week for up to 4 weeks for short bowel syndrome only
Medication	Pediatric Maximum Dosing per week (age < 18 years)	Adult Maximum Dosing per week (age ≥ 18 years)																																							
Genotropin	0.48 mg/kg/week	0.08 mg/kg/week																																							
Humatrope	0.47 mg/kg/week	0.0875 mg/kg/week																																							
Ngenla	0.66 mg/kg/week	Not Indicated																																							
Norditropin Flexpro	0.47 mg/kg/week	0.112 mg/kg/week																																							
Nutropin AQ Nuspin	0.7 mg/kg/week	0.175 mg/kg/week for ≤35 years of age 0.0875 mg/kg/week for >35 years of age																																							
Omnitrope	0.48 mg/kg/week	0.08 mg/kg/week																																							
Saizen	0.18 mg/kg/week	0.07 mg/kg/week																																							
Serostim	Not Indicated	42 mg/week for HIV wasting or cachexia (in combination with antiretroviral therapy)																																							
Skytrofa	1.68 mg/kg/week	Not Indicated																																							
Sogroya	Dose Individualized for each patient, based on growth response	8 mg/week																																							
Zomacton	0.47 mg/kg/week	0.0875 mg/kg/week																																							
Zorbtive	Not Indicated	56 mg/week for up to 4 weeks for short bowel syndrome only																																							
<h2>VII. Gastrointestinal</h2>																																									
Therapeutic Drug Class: BILE SALTS – <i>Effective 7/1/2025</i>																																									
No PA Required Ursodiol capsule Ursodiol tablet	PA Required BYLVAY (odevixibat) capsule, pellet CHENODAL (chenodiol) tablet	Actigall (ursodiol) may be approved for members who meet the following criteria: <ul style="list-style-type: none">• Member is ≥ 18 years of age AND• Member has tried and failed therapy with a 12-month trial of a preferred ursodiol product (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions). Chenodal (chenodiol) may be approved for members who meet the following criteria:																																							

	<p>CHOLBAM (cholic acid) capsule</p> <p>LIVMARLI (maralixibat) solution, tablet</p> <p>OCALIVA (obeticholic acid) tablet</p> <p>RELTONE (ursodiol) capsule</p> <p>URSO (ursodiol) tablet</p> <p>URSO FORTE (ursodiol) tablet</p>	<ul style="list-style-type: none">• Member is > 18 years of age AND• Member has tried and failed therapy with a 12-month trial of a preferred ursodiol product (failure is defined as lack of efficacy, contraindication, allergy, intolerable side effects or significant drug-drug interactions). If chenodiol is being prescribed for treatment of cerebrotendinous xanthomatosis, no trial and failure of ursodiol is required. <p>Cholbam (cholic acid) may be approved for members who meet the following criteria:</p> <ul style="list-style-type: none">• Bile acid synthesis disorders:<ul style="list-style-type: none">○ Member age must be greater than 3 weeks old AND○ Member has a diagnosis for bile acid synthesis disorder due to single enzyme defect (Single Enzyme-Defect Disorders: Defective sterol nucleus synthesis, 3β-hydroxy-Δ-c27-steroid oxidoreductase deficiency, AKR1D1 deficiency, CYP7A1 deficiency, Defective side-chain synthesis, CYP27A1 deficiency (cerebrotendinous xanthomatosis), 2-methylacyl-CoA racemase deficiency (AMACR), 25-hydroxylation pathway (Smith–Lemli–Opitz).• Peroxisomal disorder including Zellweger spectrum disorders:<ul style="list-style-type: none">○ Member age must be greater than 3 weeks old AND○ Member has diagnosis of peroxisomal disorders (PDs) including Zellweger spectrum disorders AND○ Member has manifestations of liver disease, steatorrhea or complications from decreased fat-soluble vitamin absorption. <p>Ocaliva (obeticholic acid) may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none">• Member is ≥18 years of age AND• Medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider AND• Member has the diagnosis of primary biliary cholangitis without cirrhosis OR a diagnosis of primary biliary cholangitis with compensated cirrhosis with no evidence of portal hypertension AND• Member has failed treatment with a preferred ursodiol product for at least 6 months due to an inadequate response, intolerable side effects, drug-drug interaction, or allergy to preferred ursodiol formulations. <p>Reltone (ursodiol) may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none">• Member is ≥ 18 years of age AND• The requested medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider AND• The requested medication is being prescribed for one of the following:<ul style="list-style-type: none">○ Treatment of radiolucent, noncalcified gallbladder stones < 20 mm in greatest diameter AND elective cholecystectomy would be undertaken except for the presence of increased surgical risk due to systemic disease, advanced age, idiosyncratic reaction to general anesthesia, or for those patients who refuse surgery OR○ Prevention of gallstone formation in obese patients experiencing rapid weight lossAND• No compelling reasons for the member to undergo cholecystectomy exist, including unremitting acute cholecystitis, cholangitis, biliary obstruction, gallstone pancreatitis, or biliary-gastrointestinal fistula, AND• Member has trialed and failed treatment with a preferred ursodiol product for at least 6 months due to an inadequate response, intolerable side effects, drug-drug interaction, or allergy to inactive ingredients contained in the preferred ursodiol formulations. <p><u>Initial approval:</u> 1 year</p>
--	--	--

		<p><u>Reauthorization:</u> May be reauthorized for 1 additional year with provider attestation that partial or complete stone dissolution was observed after completion of the initial year of Reltone therapy. Maximum cumulative approval per member is 24 months.</p> <p>Urso (ursodiol) and Urso Forte (ursodiol) may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none">• Member is ≥ 18 years of age AND• Medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider AND• Member has the diagnosis of Primary Biliary Cholangitis as evidenced by two of the following at the time of diagnosis:<ul style="list-style-type: none">○ Evidence of cholestasis with an alkaline phosphatase elevation of at least 1.5 times the upper limit of normal○ Presence of antimitochondrial antibody with titer of 1:40 or higher○ Histologic evidence of nonsuppurative destruction cholangitis and destruction of interlobular bile ducts AND• Member has failed treatment with a preferred ursodiol product for at least 6 months due to an inadequate response, intolerable side effects, drug-drug interaction, or allergy to inactive ingredients contained in the preferred ursodiol formulations. <p>Requests for drug products that are FDA-indicated for the treatment of nonalcoholic steatohepatitis (NASH) may be approved if meeting the following:</p> <ul style="list-style-type: none">• A diagnosis of NASH has been confirmed through liver biopsy AND• Member meets the FDA-labeled minimum age requirement for the prescribed product AND• Member does not have significant liver disease other than NASH, AND• The requested medication is being prescribed for use for the FDA-labeled indication and as outlined in product package labeling AND• Medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider. <p>Non-preferred products prescribed for FDA-labeled indications not identified above may receive approval for use as outlined in product package labeling.</p>
Therapeutic Drug Class: ANTI-EMETICS, Oral – <i>Effective 7/1/2025</i>		
<p>No PA Required</p> <p>DICLEGIS DR^{BNR} tablet (doxylamine/pyridoxine)</p> <p>Meclizine (Rx) 12.5 mg, 25 mg tablet</p> <p>Metoclopramide solution, tablet</p> <p>Ondansetron ODT; 4mg, 8mg tablet</p>	<p>PA Required</p> <p>AKYNZEO (netupitant/palonosetron) capsule</p> <p>ANTIVERT (meclizine) 50 mg tablet</p> <p>ANZEMET (dolasetron) tablet</p>	<p>Emend (aprepitant) TriPack or Emend (aprepitant) powder kit may be approved following trial and failure of two preferred products AND Emend (aprepitant) <u>capsule</u>. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Doxylamine/pyridoxine tablet (generic) or Bonjesta (doxylamine/pyridoxine) may be approved for 9 months if meeting the following criteria:</p> <ul style="list-style-type: none">• Member has nausea and vomiting associated with pregnancy AND• Member has trialed and failed DICLEGIS DR tablet AND one of the following (failure is defined as lack of

Ondansetron oral suspension/ solution	Aprepitant capsule, tripack	<p>efficacy with a 7-day trial, allergy, intolerable side effects, or significant drug-drug interaction):</p> <ul style="list-style-type: none">○ Antihistamine (such as diphenhydramine, dimenhydrinate, meclizine) OR○ Dopamine antagonist (such as metoclopramide, prochlorperazine, promethazine) OR○ Serotonin antagonist (ondansetron, granisetron) <p>All other non-preferred products may be approved for members who have trialed and failed treatment with two preferred products. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Dronabinol prior authorization may be approved for members meeting above non-preferred criteria OR via AutoPA for members with documented HIV diagnosis.</p> <p>Promethazine product formulations require prior authorization for members < 2 years of age due to risk of fatal respiratory depression.</p>
Prochlorperazine tablet	BONJESTA ER (doxylamine/pyridoxine) tablet	
Promethazine syrup, tablet	Doxylamine/pyridoxine tablet (generic Diclegis)	
	Dronabinol capsule	
	EMEND (aprepitant) capsule, powder for suspension, dose/tri- pack	
	Granisetron tablet	
	MARINOL (dronabinol) capsule	
	Ondansetron 16mg tablet	
	REGLAN (metoclopramide) tablet	
	Trimethobenzamide capsule	
	ZOFRAN (ondansetron) tablet	
Therapeutic Drug Class: ANTI-EMETICS, Non-Oral – <i>Effective 7/1/2025</i>		
No PA Required	PA Required	Non-preferred products may be approved for members who have trialed and failed treatment with two preferred products. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.
Prochlorperazine 25 mg suppository	PROMETHEGAN 50 mg (Promethazine) suppository	
Promethazine 12.5 mg, 25 mg suppository	SANCUSO (granisetron) patch	
Scopolamine patch	TRANSDERM-SCOP (scopolamine) patch	
Therapeutic Drug Class: GI MOTILITY, CHRONIC – <i>Effective 7/1/2025</i>		
PA Required for all agents in this class		All agents will only be approved for FDA labeled indications and up to FDA approved maximum doses listed below.
Preferred	Non-Preferred	Preferred agents may be approved if the member meets the following criteria: <ul style="list-style-type: none">• Has diagnosis of Irritable Bowel Syndrome – Constipation (IBS-C), Chronic Idiopathic Constipation (CIC), Functional Constipation (FC), or Opioid Induced Constipation (OIC) in patients with opioids prescribed for noncancer pain AND• Member does not have a diagnosis of GI obstruction AND
LINZESS (linaclotide) capsule	Alosetron tablet	
Lubiprostone capsule	AMITIZA (lubiprostone) capsule	

MOVANTIK (naloxegol) tablet	IBSRELA tablet LOTRONEX (alosetron) tablet MOTEGRITY (prucalopride) tablet Prucalopride tablet RELISTOR (methylnaltrexone) syringe, tablet, vial SYMPROIC (naldemedine) tablet TRULANCE (plecanatide) tablet VIBERZI (eluxadoline) tablet	<ul style="list-style-type: none">For indication of OIC, member opioid use must exceed 4 weeks of treatment ANDFor indications of CIC, OIC, IBS-C; member must have documentation of adequate trial of two or more over-the-counter motility agents (polyethylene glycol, docusate or bisacodyl, for example). OR If the member cannot take oral medications, then the member must fail a 7-day trial with a nonphosphate enema (docusate or bisacodyl enema). Failure is defined as a lack of efficacy for a 7-day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction ANDFor indication of IBS-D, must have documentation of adequate trial and failure with loperamide and trial and failure with dicyclomine or hyoscyamine. Failure is defined as a lack of efficacy for a 7-day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction. <p>Non-preferred agents may be approved if the member meets the following criteria:</p> <ul style="list-style-type: none">Member meets all listed criteria for preferred agents ANDMember has trialed and failed two preferred agents OR if the indication is OIC caused by methadone, then a non-preferred agent may be approved after an adequate trial of MOVANTIK (naloxegol). Failure is defined as a lack of efficacy for a 7-day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction ANDIf prescribed Viberzi (eluxadoline) or Lotronex (alosetron), member meets the additional criteria for those agents listed below. <p>VIBERZI (eluxadoline) may be approved for members who meet the following additional criteria:</p> <ul style="list-style-type: none">Diagnosis of Irritable Bowel Syndrome – Diarrhea (IBS-D) ANDMember has a gallbladder ANDMember does not have severe hepatic impairment (Child-Pugh C), history of severe constipation, known mechanical gastrointestinal obstruction, biliary duct obstruction, history of pancreatitis or structural disease of the pancreas ANDMember does not drink more than 3 alcoholic drinks per day <p>LOTRONEX (alosetron) and generic alosetron may be approved for members who meet the following additional criteria:</p> <ul style="list-style-type: none">Member is a female with Irritable Bowel Syndrome – Diarrhea (IBS-D) with symptoms lasting 6 months or longer ANDMember does not have severe hepatic impairment (Child-Pugh C), history of severe constipation or ischemic colitis, hypercoagulable state, Crohn’s disease or ulcerative colitis, or known mechanical gastrointestinal obstruction.
-----------------------------	--	--

Medication	FDA approved indication	FDA Max Dose
Amitiza (lubiprostone)	IBS-C (females only), CIC, OIC (not caused by methadone)	48mcg/day
Linzess (linaclotide)	IBS-C, CIC (≥ 18 years)	290mcg/day
Movantik (naloxegol)	OIC, FC (6 to 17 years)	25mg/day (OIC), 72mcg/day (FC)
Viberzi (eluxadoline)	IBS-D	200mg/day
Relistor subcutaneous injection (methylnaltrexone)	OIC	12mg/day

	Relistor oral (methylnaltrexone)	OIC	450mg/day
	Lotronex (alosetron)	IBS-D (women only)	2mg/day (women only)
	Symproic (Naldemedine)	OIC	0.2mg/day
	Trulance (plecanatide)	CIC, IBS-C	3mg/day
	Motegrity (prucalopride)	CIC	2mg/day
CIC – chronic idiopathic constipation, FC – functional constipation, OIC – opioid induced constipation, IBS – irritable bowel syndrome, D – diarrhea predominant, C – constipation predominant			
Therapeutic Drug Class: H. PYLORI TREATMENTS – <i>Effective 7/1/2025</i>			
No PA Required PYLERA ^{BNR} capsule (bismuth subcitrate/metronidazole tetracycline)	PA Required Amoxicillin/lansoprazole/clarithromycin pack Bismuth subcitrate/metronidazole tetracycline capsule OMECLAMOX-PAK (amoxicillin/omeprazole/clarithromycin) TALICIA (omeprazole/amoxicillin/rifabutin) tablet VOQUEZNA DUAL (vonoprazan/amoxicillin) dose pack VOQUEZNA TRIPLE (vonoprazan/amoxicillin/clarithromycin dose pack	Non-preferred <i>H. pylori</i> treatments should be used as individual product ingredients unless one of the individual products is not commercially available, then a PA for the combination product may be given.	
Therapeutic Drug Class: HEMORRHOIDAL, ANORECTAL, AND RELATED TOPICAL ANESTHETIC AGENTS – <i>Effective 7/1/2025</i>			
Hydrocortisone single agent			
No PA Required ANUSOL-HC (hydrocortisone) 2.5% cream with applicator CORTIFOAM (hydrocortisone) 10% aerosol Hydrocortisone 1% cream with applicator	PA Required CORTENEMA (hydrocortisone) enema PROCORT cream	Non-preferred products may be approved following trial and failure of therapy with 3 preferred products (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).	

Hydrocortisone 2.5% cream with applicator		
Hydrocortisone enema		
Lidocaine single agent		
No PA Required Lidocaine 3% cream, 5% ointment	PA Required	
Other and Combinations		
No PA Required	PA Required	
Lidocaine-Hydrocortisone 3-0.5% cream with applicator	ANALPRAM HC (Hydrocortisone-Pramoxine) 1%-1% cream, 2.5%-1% cream	RECTIV (nitroglycerin) ointment may be approved if meeting the following: <ul style="list-style-type: none">• Member has a diagnosis of anal fissure AND• Prescriber attests that member has trialed and maximized use of appropriate supportive therapies including sitz bath, fiber, topical analgesics (such as lidocaine), and stool softeners/laxatives.
Lidocaine-Prilocaine Cream (<i>all other manufacturers</i>)	EPIFOAM (Hydrocortisone-Pramoxine) 1%-1% foam	
PROCTOFOAM-HC (hydrocortisone-pramoxine) 1%-1% foam	Hydrocortisone-Pramoxine 1%-1%, 2.5%-1% cream	
	Lidocaine-Hydrocortisone in Coleus 2%-2% cream kit	
	Lidocaine-Hydrocortisone 2.8%-0.55% gel	
	Lidocaine-Hydrocortisone 3%-0.5% cream w/o applicator, cream kit	
	Lidocaine-Hydrocortisone 3%-1% cream kit	
	Lidocaine-Hydrocortisone 3%-2.5% gel kit	
	Lidocaine-Prilocaine Cream (<i>Fougera only</i>)	
	PLIAGLIS (lidocaine-tetracaine) 7%-7% cream	

RECTIV (nitroglycerin) ointment may be approved if meeting the following:

- Member has a diagnosis of anal fissure **AND**
- Prescriber attests that member has trialed and maximized use of appropriate supportive therapies including sitz bath, fiber, topical analgesics (such as lidocaine), and stool softeners/laxatives.

	<div>PROCORT (Hydrocortisone- Pramoxine) 1.85%-1.15% cream</div> <div>RECTIV (nitroglycerin) 0.4% ointment</div>	
Therapeutic Drug Class: PANCREATIC ENZYMES – <i>Effective 7/1/2025</i>		
<div>No PA Required</div> <div>CREON (pancrelipase) capsule</div> <div>VIOKACE (pancrelipase) tablet</div> <div>ZENPEP (pancrelipase) capsule</div>	<div>PA Required</div> <div>PERTZYE (pancrelipase) capsule</div>	<div>Non-preferred products may be approved for members who have failed an adequate trial (4 weeks) with at least two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction.</div>
Therapeutic Drug Class: PROTON PUMP INHIBITORS – <i>Effective 7/1/2025</i>		
<div>No PA Required</div> <div>Esomeprazole DR packet for oral suspension, capsule (RX)</div> <div>Lansoprazole DR capsules (RX)</div> <div>Lansoprazole ODT (RX) <i>(for members under 2 years)</i></div> <div>Omeprazole DR capsule (RX)</div> <div>Pantoprazole tablet</div> <div>PROTONIX (pantoprazole DR) packet for oral suspension^{BNR}</div>	<div>PA Required</div> <div>ACIPHEX (rabeprazole) tablet, sprinkle capsule</div> <div>DEXILANT (dexlansoprazole) capsule</div> <div>Dexlansoprazole capsule</div> <div>Esomeprazole DR 49.3 capsule (RX), (OTC) capsule</div> <div>KONVOMEF (Omeprazole/Na bicarbonate) suspension</div> <div>Lansoprazole DR capsule OTC</div> <div>Lansoprazole ODT (OTC)</div> <div>NEXIUM (esomeprazole) capsule (RX), oral suspension packet, 24HR (OTC)</div> <div>Omeprazole/Na bicarbonate capsule, packet for oral suspension</div> <div>Omeprazole DR tablet (OTC), ODT (OTC)</div>	<div>For members treating GERD symptoms that are controlled on PPI therapy, it is recommended that the dose of the PPI be re-evaluated or step-down with an H2 blocker (such as famotidine) be trialed in order to reduce long-term PPI use. Prior authorization for non-preferred proton pump inhibitors may be approved if all of the following criteria are met:</div> <div><ul style="list-style-type: none">● Member has a qualifying diagnosis (below) AND● Member has trialed and failed therapy with three preferred agents within the last 24 months. (Failure is defined as: lack of efficacy following 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction) AND● Member has been diagnosed using one of the following diagnostic methods:<ul style="list-style-type: none">○ Diagnosis made by GI specialist○ Endoscopy○ X-ray○ Biopsy○ Blood test○ Breath Test</div> <div>Qualifying Diagnoses: Barrett’s esophagus, duodenal ulcer, erosive esophagitis, gastric ulcer, GERD, GI Bleed, H. pylori infection, hypersecretory conditions (Zollinger-Ellison), NSAID-induced ulcer, pediatric esophagitis, requiring mechanical ventilation, requiring a feeding tube</div> <div>Quantity Limits: All agents will be limited to once daily dosing except when used for the following diagnoses: Barrett’s esophagus, GI Bleed, H. pylori infection, hypersecretory conditions (Zollinger-Ellison), or members who have spinal cord injury with associated acid reflux.</div> <div>Adult members with GERD on once daily, high-dose PPI therapy who continue to experience symptoms may receive initial prior authorization approval for a 4-week trial of twice daily, high-dose PPI therapy. Continuation of the twice daily dosing regimen for GERD beyond 4 weeks will require additional prior authorization approval verifying adequate</div>

	<p>Pantoprazole packet for oral suspension</p> <p>PREVACID (lansoprazole) capsule, Solutab, suspension</p> <p>PRILOSEC (omeprazole) suspension</p> <p>PROTONIX (pantoprazole DR) tablet</p> <p>Rabeprazole tablet</p> <p>VOQUEZNA (vonoprazan) tablet</p> <p>ZEGERID (omeprazole/Na bicarbonate) capsule, packet for oral suspension</p>	<p>member response to the dosing regimen and approval may be placed for one year. If a member with symptomatic GERD does not respond to twice daily, high-dose PPI therapy, this should be considered a treatment failure.</p> <p>Pediatric members (< 18 years of age) on once daily dosing of a PPI who continue to experience symptoms may receive one-year prior authorization approval for twice daily PPI therapy.</p> <p>Age Limits: Nexium 24H and Zegerid will not be approved for members less than 18 years of age.</p> <p>Prevacid Solutab may be approved for members < 2 years of age OR for members ≥ 2 years of age with a feeding tube.</p> <p><u>Continuation of Care:</u> Members currently taking Dexilant (dexlansoprazole) capsules may continue to receive approval for that medication.</p>
Therapeutic Drug Class: NON-BIOLOGIC ULCERATIVE COLITIS AGENTS- Oral – <i>Effective 11/21/2025</i>		
<p>No PA Required</p> <p>APRISO (mesalamine ER) capsule</p> <p>Mesalamine DR tablet (generic Lialda) (<i>Takeda, Lannet, GSMS, and Bryant Ranch Manufacturers</i>)</p> <p>Mesalamine ER capsule (generic Apriso) (<i>Teva only</i>)</p> <p>PENTASA^{BNR} (mesalamine) capsule</p> <p>Sulfasalazine IR and DR tablet</p>	<p>PA Required</p> <p>AZULFIDINE (sulfasalazine) Entab, tablet</p> <p>Balsalazide capsule</p> <p>Budesonide DR tablet</p> <p>COLAZAL (balsalazide) capsule</p> <p>DELZICOL (mesalamine DR) capsule</p> <p>DIPENTUM (olsalazine) capsule</p> <p>LIALDA (mesalamine DR) tablet</p> <p>Mesalamine DR tablet (generic Asacol HD, Lialda – all other manufacturers)</p> <p>Mesalamine DR/ER capsule (generic Delzicol and Pentasa)</p>	<p>Prior authorization for non-preferred oral formulations will require trial and failure of two preferred oral products with different active ingredients AND one preferred rectal product. If inflammation is not within reach of topical therapy, trial of preferred rectal product is not required. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Uceris (budesonide) tablet: Prior authorization may be approved following trial and failure of one preferred oral product AND one preferred rectal product. If inflammation is not within reach of topical therapy, trial of preferred rectal product is not required. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. Approval will be placed for 8 weeks. Further prior authorization may be approved if 7 days of steroid-free time has elapsed, and member continues to meet the above criteria.</p>

	UCERIS (budesonide) tablet	
Therapeutic Drug Class: NON-BIOLOGIC ULCERATIVE COLITIS AGENTS- Rectal – <i>Effective 7/1/2025</i>		
No PA Required Mesalamine suppository Mesalamine 4gm/60 ml enema (generic SF ROWASA) SF ROWASA enema, kit (mesalamine)	PA Required Budesonide foam CANASA (mesalamine) suppository Mesalamine enema, kit ROWASA enema, kit (mesalamine) UCERIS (budesonide) foam	Prior authorization for non-preferred rectal formulations will require trial and failure of one preferred rectal formulation and one preferred oral formulation (Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction). Uceris (budesonide) foam: If the above criteria are met, Uceris (budesonide) foam prior authorization may be approved for 6 weeks. Further prior authorization may be approved if 7 days of steroid-free time has elapsed, and member continues to meet the above criteria.
VIII. Hematological		
Therapeutic Drug Class: ANTICOAGULANTS- Oral – <i>Effective 7/1/2025</i>		
No PA Required Dabigatran capsule ELIQUIS (apixaban) tablet, tablet pack Warfarin tablet XARELTO (rivaroxaban) ^{BNR} 10 mg, 15 mg, 20 mg tablet, dose pack	PA Required PRADAXA (dabigatran) capsule, pellet Rivaroxaban tablet Rivaroxaban oral suspension SAVAYSA (edoxaban) tablet XARELTO (rivaroxaban) 2.5 mg tablet XARELTO (rivaroxaban) oral suspension	SAVAYSA (edoxaban) may be approved if all the following criteria have been met: <ul style="list-style-type: none">• The member has failed therapy with two preferred agents. (Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction) AND• Member is not on dialysis AND• Member does not have CrCl > 95 mL/min AND• The member has a diagnosis of deep vein thrombosis (DVT), pulmonary embolism (PE) OR• The member has a diagnosis of non-valvular atrial fibrillation AND• The member does not have a mechanical prosthetic heart valve XARELTO 2.5mg (rivaroxaban) may be approved for members meeting all of the following criteria: <ul style="list-style-type: none">• Xarelto 2.5mg is being prescribed to reduce major CV events in members diagnosis of chronic coronary artery disease (CAD) or peripheral artery disease AND• Xarelto 2.5mg is being taken twice daily and in combination with aspirin 75-100mg daily AND• Member must not be receiving dual antiplatelet therapy, other non-aspirin antiplatelet therapy, or other oral anticoagulant AND• Member must not have had an ischemic, non-lacunar stroke within the past month AND• Member must not have had a hemorrhagic or lacunar stroke at any time XARELTO (rivaroxaban) oral suspension may be approved without prior authorization for members <18 years of age who require a rivaroxaban dose of less than 10 mg OR with prior authorization verifying the member is unable to use the solid oral dosage form. All other non-preferred oral agents require trial and failure of two preferred oral agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.

		Continuation of Care: Members with current prior authorization approval on file for a non-preferred <u>oral</u> anticoagulant medication may continue to receive approval for that medication
Therapeutic Drug Class: ANTICOAGULANTS- Parenteral – <i>Effective 7/1/2025</i>		
No PA Required Enoxaparin syringe Enoxaparin vial	PA Required ARIXTRA (fondaparinux) syringe Fondaparinux syringe FRAGMIN (dalteparin) vial, syringe LOVENOX (enoxaparin) syringe, vial	Non-preferred parenteral anticoagulants may be approved if member has trial and failure of one preferred parenteral agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction ARIXTRA (fondaparinux) may be approved if the following criteria have been met: <ul style="list-style-type: none">• Member is 18 years of age or older AND• Member has a CrCl > 30 ml/min AND• Member weighs > 50 kg AND• Member has a documented history of heparin induced-thrombocytopenia OR• Member has a contraindication to enoxaparin Members currently stabilized on fondaparinux (Arixtra) or dalteparin (Fragmin) may receive prior authorization approval to continue receiving that medication.
Therapeutic Drug Class: ANTI-PLATELETS – <i>Effective 4/8/2025</i>		
No PA Required Aspirin/dipyridamole ER capsule BRILINTA (ticagrelor) tablet ^{BNR} Cilostazol tablet Clopidogrel tablet Dipyridamole tablet Pentoxifylline ER tablet Prasugrel tablet	PA Required EFFIENT (prasugrel) tablet PLAVIX (clopidogrel) tablet Ticagrelor tablet	Zontivity (vorapaxar) may be approved for patients with a diagnosis of myocardial infarction or peripheral artery disease without a history of stroke, transient ischemic attack, intracranial bleeding, or active pathological bleeding. Patients must also be taking aspirin and/or clopidogrel concomitantly. Non-preferred products without criteria will be reviewed on a case-by-case basis.
Therapeutic Drug Class: COLONY STIMULATING FACTORS – <i>Effective 7/1/2025</i>		
PA Required for all agents in this class*		*Prior authorization for preferred agents may be approved if meeting the following criteria: <ul style="list-style-type: none">• Medication is being used for one of the following indications:<ul style="list-style-type: none">○ Patient with cancer receiving myelosuppressive chemotherapy –to reduce incidence of infection (febrile neutropenia) (Either the post nadir ANC is less than 10,000 cells/mm3 or the risk of neutropenia for the member is calculated to be greater than 20%)○ Acute Myeloid Leukemia (AML) patients receiving chemotherapy○ Bone Marrow Transplant (BMT)○ Peripheral Blood Progenitor Cell Collection and Therapy○ Hematopoietic Syndrome of Acute Radiation Syndrome
Preferred FULPHILA (pegfilgrastim-jmdb) syringe NEUPOGEN (filgrastim) vial, syringe	Non-Preferred FYLNETRA (pegfilgrastim-jmdb) syringe GRANIX (tbo-filgrastim) syringe, vial LEUKINE (sargramostim) vial	

	<div>NEULASTA (pegfilgrastim) kit, syringe</div> <div>NIVESTYM (filgrastim-aafi) syringe, vial</div> <div>NYVEPRIA (pegfilgrastim-apgf) syringe</div> <div>RELEUKO (filgrastim-ayow) syringe, vial</div> <div>RYZNEUTA (efbmalenograstim alfa-vuxw) syringe</div> <div>STIMUFEND (pegfilgrastim-fpgk) syringe</div> <div>UDENYCA (pegfilgrastim-cbqv) autoinjector, On-Body, syringe</div> <div>ZARXIO (filgrastim-sndz) syringe</div> <div>ZIEXTENZO (pegfilgrastim-bmez) syringe</div>	<div><div><div>○ Severe Chronic Neutropenia (Evidence of neutropenia infection exists or ANC is below 750 cells/mm3)</div></div></div> <div>Prior authorization for non-preferred agents may be approved if meeting the following criteria:</div> <div><div><div>● Medication is being used for one of the following indications:</div><div><div><div>○ Patient with cancer receiving myelosuppressive chemotherapy –to reduce incidence of infection (febrile neutropenia) (Either the post nadir ANC is less than 10,000 cells/mm3 or the risk of neutropenia for the member is calculated to be greater than 20%)</div><div>○ Acute Myeloid Leukemia (AML) patients receiving chemotherapy</div><div>○ Bone Marrow Transplant (BMT)</div><div>○ Peripheral Blood Progenitor Cell Collection and Therapy</div><div>○ Hematopoietic Syndrome of Acute Radiation Syndrome</div><div>○ Severe Chronic Neutropenia (Evidence of neutropenia infection exists or ANC is below 750 cells/mm3)</div></div></div></div><div>AND</div><div><div><div>● Member has history of trial and failure of Neupogen AND one other preferred agent. Failure is defined as a lack of efficacy with a 3-month trial, allergy, intolerable side effects, significant drug-drug interactions, or contraindication to therapy. Trial and failure of Neupogen will not be required if meeting one of the following:</div><div><div><div>○ Member has limited access to caregiver or support system for assistance with medication administration OR</div><div>○ Member has inadequate access to healthcare facility or home care interventions.</div></div></div></div></div></div>
Therapeutic Drug Class: ERYTHROPOIESIS STIMULATING AGENTS – <i>Effective 7/1/2025</i>		
PA Required for all agents in this class*		
<div>Preferred</div> <div>EPOGEN (epoetin alfa) vial</div> <div>RETACRIT (epoetin alfa-epbx) (<i>Pfizer only</i>) vial</div>	<div>Non-Preferred</div> <div>ARANESP (darbepoetin alfa) syringe, vial</div> <div>MIRCERA (methoxy peg-epoetin beta) syringe</div> <div>PROCRIT (epoetin alfa) vial</div> <div>RETACRIT (epoetin alfa-epbx) (<i>Vifor only</i>) vial</div>	<div>*Prior Authorization is required for all products and may be approved if meeting the following:</div> <div><div><div>● Medication is being administered in the member’s home or in a long-term care facility AND</div><div>● Member meets <u>one</u> of the following:</div><div><div><div>○ A diagnosis of cancer, currently receiving chemotherapy, with chemotherapy-induced anemia, and hemoglobin[†] of 10g/dL or lower OR</div><div>○ A diagnosis of chronic renal failure, and hemoglobin[†] below 10g/dL OR</div><div>○ A diagnosis of hepatitis C, currently taking ribavirin and failed response to a reduction of ribavirin dose, and hemoglobin[†] less than 10g/dL (or less than 11g/dL if symptomatic) OR</div><div>○ A diagnosis of HIV, currently taking zidovudine, hemoglobin[†] less than 10g/dL, and serum erythropoietin level of 500 mU/mL or less OR</div><div>○ Member is undergoing elective, noncardiac, nonvascular surgery and medication is given to reduce receipt of allogenic red blood cell transfusions, hemoglobin[†] is greater than 10g/dL, but less than or equal to 13g/dL and high risk for perioperative blood loss. Member is not willing or unable to donate autologous blood pre-operatively</div></div></div></div><div>AND</div><div><div><div>● For any non-preferred product, member has trialed and failed treatment with one preferred product. Failure is defined as lack of efficacy with a 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction.</div></div></div></div>

		†Hemoglobin results must be from the last 30 days.
<div>IX. Immunological</div>		
Therapeutic Drug Class: IMMUNE GLOBULINS – <i>Effective 1/1/2026</i>		
PA Required for all agents in this class*		Preferred agents may be approved for members meeting at least one of the approved conditions for immune globulin use listed below for prescribed doses not exceeding maximum (Table 1). Non-preferred agents may be approved for members meeting the following: <ul style="list-style-type: none">Member meets at least one of the approved conditions listed below ANDMember has history of trial and failure of two preferred agents (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions) ANDPrescribed dose does not exceed listed maximum (Table 1) Approved Conditions for Immune Globulin Use: <ul style="list-style-type: none">Primary Humoral Immunodeficiency disorders including:<ul style="list-style-type: none">Common Variable Immunodeficiency (CVID)Severe Combined Immunodeficiency (SCID)X-Linked AgammaglobulinemiaX-Linked with Hyperimmunoglobulin M (IgM) ImmunodeficiencyWiskott-Aldrich SyndromeMembers < 13 years of age with pediatric Human Immunodeficiency Virus (HIV) and CD-4 count > 200/mm3Neurological disorders including:<ul style="list-style-type: none">Guillain-Barré SyndromeRelapsing-Remitting Multiple SclerosisChronic Inflammatory Demyelinating PolyneuropathyMyasthenia GravisPolymyositis and DermatomyositisMultifocal Motor NeuropathyKawasaki SyndromeChronic Lymphocytic Leukemia (CLL)Autoimmune Neutropenia (AN) with absolute neutrophil count < 800 mm and history of recurrent bacterial infectionsAutoimmune Hemolytic Anemia (AHA)Liver or Intestinal TransplantImmune Thrombocytopenia Purpura (ITP) including:<ul style="list-style-type: none">Requiring preoperative therapy for undergoing elective splenectomy with platelet count < 20,000/mcLMembers with active bleeding & platelet count <30,000/mcLPregnant members with platelet counts <10,000/mcL in the third trimesterPregnant members with platelet count 10,000 to 30,000/mcL who are bleedingMultisystem Inflammatory Syndrome in Children (MIS-C)Measles post-exposure prophylaxis (PEP)
Preferred	Non-Preferred	
BIVIGAM 10% IV liquid	ALYGLO 10% IV liquid	
CUTAQUIG 16.5% SQ liquid	ASCENIV 10% IV liquid	
CUVITRU 20% SQ liquid	FLEBOGAMMA DIF 5%, 10% IV liquid	
GAMMAGARD 10% IV/SQ liquid	GAMMAGARD S/D vial	
GAMMAKED 10% IV/SQ liquid	GAMMAPLEX 5%, 10% IV liquid	
GAMUNEX-C 10% IV/SQ liquid	HIZENTRA 20% SQ syringe, vial	
PRIVIGEN 10% IV liquid	HYQVIA 10% SQ liquid	
<i>If immune globulin is being administered in a long-term care facility or in a member's home by a home healthcare provider, it should be billed as a pharmacy claim. All other claims must be submitted through the medical benefit.</i>	OCTAGAM 5%, 10% IV liquid	
	PANZYGA 10% IV liquid	
	XEMBIFY 20% IV liquid	

			Table 1: FDA-Approved Maximum Immune Globulin Dosing	
			Alyglo – IV admin	800mg/kg every 3 to 4 weeks
			Asceniv – IV admin	800 mg/kg every 3 to 4 weeks
			Bivigam – IV admin	800 mg/kg every 3 to 4 weeks
			Cutaquig – subcutaneous admin	See product labeling
			Cuvitru –subcutaneous admin	12 grams protein/site for up to four sites weekly (48grams/week)
			Flebogamma DIF – IV admin	600 mg/kg every 3 weeks
			Gammaplex 5% – IV admin	1 gram/kg for 2 consecutive days (ITP) 800 mg/kg every 3 to 4 weeks (PI)
			Gammagard liquid subcutaneous or IV admin	2.4 grams/kg/month (IV for MMN) 2 grams/kg over 2 to 5 consecutive days (IV for CIDP) 600 mg/kg every 3 weeks (IV for PI)
			Gammaked –subcutaneous or IV admin	2 grams/kg over 2 consecutive days (IV for ITP, CIDP) 600 mg/kg every 3 weeks (IV for PI)
			Gamunex-C –subcutaneous or IV admin	2 grams/kg over 2 to 5 consecutive days (IV for ITP, CIDP) 600 mg/kg every 3 weeks (IV for PI)
			Hizentra –subcutaneous admin	0.4 grams/kg per week over 2 consecutive days (CIDP)
			Octagam – IV admin	2 grams/kg over 2 to 5 consecutive days (ITP, DM) 600 mg/kg every 3 weeks (PI)
			Panzyga – IV admin	2 grams/kg over 2 consecutive days (ITP, CIDP) 600 mg/kg every 3 weeks (PI)
			Privigen – IV admin	2 grams/kg over 2 to 5 consecutive days (ITP, CIDP) 800 mg/kg every 3 weeks (PI)
Xembify – subcutaneous admin	150 mg/kg/day for 5 consecutive days (PI loading dose)			
CIDP=Chronic Inflammatory Demyelinating Polyneuropathy; DM=Dermatomyositis; ITP= Chronic Immune Thrombocytopenic				

		<div>Purpura; MMN=Multifocal Motor Neuropathy; PI=Primary Humoral Immunodeficiency</div>
		Members currently receiving a preferred or non-preferred immunoglobulin product may receive approval to continue therapy with that product at prescribed doses not exceeding maximum (Table 1).
Therapeutic Drug Class: NEWER GENERATION ANTIHISTAMINES – <i>Effective 1/1/2026</i>		
No PA Required Cetirizine (OTC) syrup/solution (OTC/RX), tablet Desloratadine tablet (RX) Levocetirizine tablet (RX/OTC) Loratadine tablet (OTC), syrup/solution (OTC)	PA Required Cetirizine (OTC) chewable tablet, softgel, UD cups solution CLARINEX (desloratadine) tablet Desloratadine ODT (RX) Fexofenadine tablet (OTC), suspension (OTC) Levocetirizine solution (RX) Loratadine chewable (OTC), ODT (OTC)	<p>Non-preferred single agent antihistamine products may be approved for members who have failed treatment with two preferred products in the last 6 months. For members with respiratory allergies, an additional trial of an intranasal corticosteroid will be required in the last 6 months.</p> <p>Failure is defined as lack of efficacy with a 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p>
Therapeutic Drug Class: ANTIHISTAMINE/DECONGESTANT COMBINATIONS – <i>Effective 1/1/2026</i>		
No PA Required Cetirizine-PSE ER (OTC) Loratadine-D (OTC) tablet	PA Required CLARINEX-D (desloratadine-D) Fexofenadine/PSE (OTC)	<p>Non-preferred antihistamine/decongestant combinations may be approved for members who have failed treatment with the preferred product in the last 6 months. For members with respiratory allergies, an additional trial of an intranasal corticosteroid will be required in the last 6 months.</p> <p>Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.</p>
Therapeutic Drug Class: INTRANASAL RHINITIS AGENTS – <i>Effective 1/1/2026</i>		
No PA Required Azelastine 137 mcg Budesonide (OTC) DYMISTA (azelastine/ fluticasone) ^{BNR} Fluticasone (RX) Ipratropium	PA Required Azelastine (Astepro) 0.15% Azelastine/Fluticasone BECONASE AQ (beclomethasone dipropionate) Flunisolide 0.025%	<p>Non-preferred products may be approved following trial and failure of treatment with three preferred products (failure is defined as lack of efficacy with a 2-week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>Non-preferred combination agents may be approved following trial of individual products with same active ingredients AND trial and failure of one additional preferred agent (failure is defined as lack of efficacy with 2-week trial, allergy, intolerable side effects or significant drug-drug interactions).</p>

Olopatadine Triamcinolone acetonide (OTC)	Fluticasone (OTC)	
	Mometasone	
	NASONEX (mometasone)	
	OMNARIS (ciclesonide)	
	PATANASE (olopatadine)	
	QNASL (beclomethasone)	
	RYALTRIS (olopatadine/mometasone)	
	XHANCE (fluticasone)	
	ZETONNA (ciclesonide)	
Therapeutic Drug Class: LEUKOTRIENE MODIFIERS – <i>Effective 1/1/2026</i>		
No PA Required Montelukast tablet, chewable	PA Required ACCOLATE (zafirlukast) tablet Montelukast granules SINGULAIR (montelukast) tablet, chewable, granules Zafirlukast tablet Zileuton ER tablet ZYFLO (zileuton) tablet	Non-preferred products may be approved if meeting the following criteria: <ul style="list-style-type: none">Member has trialed and failed treatment with one preferred product (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions) ANDMember has a diagnosis of asthma. Montelukast granules may be approved if a member has tried and failed montelukast chewable tablets AND has difficulty swallowing.
Therapeutic Drug Class: METHOTREXATE PRODUCTS – <i>Effective 1/1/2026</i>		
No PA Required Methotrexate tablet, vial	PA Required JYLAMVO (methotrexate) solution OTREXUP (methotrexate) auto-injector	OTREXUP, REDITREX or RASUVO may be approved if meeting the following criteria: <ul style="list-style-type: none">Member has diagnosis of severe, active rheumatoid arthritis OR active polyarticular juvenile idiopathic arthritis (pJIA) OR inflammatory bowel disease (IBD) ANDMember has trialed and failed preferred methotrexate tablet formulation (failure is defined as lack of efficacy, allergy, intolerable side effects, inability to take oral product formulation, or member has a diagnosis of pJIA and provider has determined that the subcutaneous formulation is necessary to optimize methotrexate therapy) ANDMember (or parent/caregiver) is unable to administer preferred methotrexate vial formulation due to limited functional ability (such as vision impairment, limited manual dexterity and/or limited hand strength).

	<div>RASUVO (methotrexate) auto-injector</div> <div>REDITREX (methotrexate) syringe</div> <div>TREXALL (methotrexate) tablet</div> <div>XATMEP (methotrexate) solution</div>	<p>TREXALL may be approved if meeting the following criteria:</p> <ul style="list-style-type: none">Member has trialed and failed preferred methotrexate tablet formulation. Failure is defined as allergy or intolerable side effects. <p>XATMEP may be approved for members who meet the following criteria:</p> <ul style="list-style-type: none">Member is < 18 years of ageMember has a diagnosis of acute lymphoblastic leukemia ORMember has a diagnosis of active polyarticular juvenile idiopathic arthritis (pJIA) and has had an insufficient therapeutic response to, or is intolerant to, an adequate trial of first-line therapy including full dose non-steroidal anti-inflammatory agents (NSAIDs) ANDMember has a documented swallowing difficulty due to young age and/or a medical condition and is unable to use the preferred methotrexate tablet formulation <p><i>Methotrexate can cause serious embryo-fetal harm when administered during pregnancy and it is contraindicated for use during pregnancy for the treatment of non-malignant diseases. Advise members of reproductive potential to use effective contraception during and after treatment with methotrexate, according to FDA product labeling.</i></p> <p>Members currently stabilized on a non-preferred methotrexate product may receive approval to continue that agent.</p>
Therapeutic Drug Class: MULTIPLE SCLEROSIS AGENTS – <i>Effective 6/5/2025</i>		
Disease Modifying Therapies		
<div>Preferred No PA Required (Unless indicated*)</div> <div>AVONEX (interferon beta 1a) pen, syringe</div> <div>BETASERON (interferon beta 1b) injection</div> <div>COPAXONE (glatiramer) 20mg injection ^{BNR}</div> <div>Dimethyl fumarate tablet, starter pack</div> <div>Fingolimod capsule</div> <div>Glatiramer 40mg injection</div> <div>*KESIMPTA (ofatumumab) pen ^{**2nd Line**}</div> <div>Teriflunomide tablet</div>	<div>Non-Preferred PA Required</div> <div>AUBAGIO (teriflunomide) tablet</div> <div>BAFIERTAM (monomethyl fumarate DR) capsule</div> <div>COPAXONE (glatiramer) 40mg injection</div> <div>EXTAVIA (interferon beta 1b) kit, vial</div> <div>GILENYA (fingolimod) capsule</div> <div>Glatiramer 20mg</div> <div>GLATOPA (glatiramer) injection</div> <div>MAVENCLAD (cladribine) tablet</div>	<p>*Kesimpta (ofatumumab) may be approved if member has trialed and failed treatment with one preferred agent (failure is defined as intolerable side effects, contraindication to therapy, drug-drug interaction, or lack of efficacy).</p> <p><u>Non-Preferred Products:</u></p> <p>Non-preferred products may be approved if meeting the following:</p> <ul style="list-style-type: none">Member has a diagnosis of a relapsing form of multiple sclerosis ANDMember has previous trial and failure with three preferred agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction ANDPrescribed dose does not exceed the maximum FDA-approved dose for the medication being ordered ANDIf indicated in the product labeling, a negative pre-treatment pregnancy test has been documented, ANDIf indicated in the product labeling, an ophthalmologic examination has been performed and documented prior to medication initiation, ANDThe request meets additional criteria listed for any of the following: <p>Mayzent (siponimod):</p> <ul style="list-style-type: none">Member has previous trial and failure of three preferred agents, one of which must be Gilenya (fingolimod). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. <p>Mavenclad (cladribine):</p>

	MAYZENT (siponimod) tablet, pack PLEGRIDY (peg-interferon beta 1a) pen, syringe PONVORY (ponesimod) tablet, pack REBIF (interferon beta 1a) syringe REBIF REDIDOSE (interferon beta 1a) pen TASCENSO ODT (fingolimod) tablet TECFIDERA (dimethyl fumarate) tablet, pack VUMERITY (diroximel DR) capsule ZEPOSIA (ozanimod) capsule, kit, starter pack	<ul style="list-style-type: none">Member has history of ≥ 1 relapse in the 12 months preceding initiation of therapy ANDMember has previous trial and failure of three other therapies for relapsing forms of multiple sclerosis (failure is defined as lack of efficacy with 3-month trial, allergy, intolerable side effects, or significant drug-drug interactions) <p>Vumerity (diroximel fumarate) or Bafiertam (monomethyl fumarate DR):</p> <ul style="list-style-type: none">Member has previous trial and failure of three preferred agents, one of which must be Tecfidera (dimethyl fumarate). Failure is defined as lack of efficacy, allergy, significant drug-drug interactions, intolerable side effects (if GI adverse events, must meet additional criteria below) ANDIf the requested medication is being prescribed due to GI adverse events with Tecfidera therapy (and no other reason for failure of Tecfidera is given), then the following additional criteria must be met:<ul style="list-style-type: none">Member has trialed a temporary dose reduction of Tecfidera (with maintenance dose being resumed within 4 weeks) ANDMember has trialed taking Tecfidera with food ANDGI adverse events remain significant despite maximized use of gastrointestinal symptomatic therapies (such as calcium carbonate, bismuth subsalicylate, PPIs, H2 blockers, anti-bloating/anti-constipation agents, anti-diarrheal, and centrally acting anti-emetics) ANDInitial authorization will be limited to 3 months. Continuation (12-month authorization) will require documentation of clinically significant reduction in GI adverse events. <p>Members currently stabilized on a preferred second line (Kesimpta) or non-preferred product (may receive approval to continue therapy with that agent).</p>
Symptom Management Therapies		
No PA Required Dalfampridine ER tablet	PA Required AMPYRA ER (dalfampridine) tablet	Non-preferred products may be approved with prescriber attestation that there is clinical rationale supporting why the preferred brand/generic equivalent product formulation is unable to be used. <u>Maximum Dose:</u> Ampyra (dalfampridine) 10mg twice daily
Therapeutic Drug Class: TARGETED IMMUNE MODULATORS – <i>Effective 1/1/2026</i> <i>Preferred agents:</i> Adalimumab-aacf syringe, aaty and adbm; ADBRY (tralokinumab-ldrm); AMJEVITA (adalimumab-atto), YUFLYMA (adalimumab-aaty) Cyltezo (adalimumab-adbm); DUPIXENT (dupilumab); ENBREL (etanercept); FASENRA (benralizumab) pen; OTEZLA (apremilast) tablet; KEVZARA (sarilumab); TALTZ (ixekizumab); TEZSPIRE (tezepelumab-ekko) pen; XELJANZ ER/IR (tofacitinib) tablet; XOLAIR (omalizumab) syringe; IMULDOSA (ustekinumab-slrf); STEQEYMA (ustekinumab-stba) syringe; SELARSDI (ustekinumab-AEKN) syringe		
Rheumatoid Arthritis, all other Arthritis (except psoriatic arthritis, see below), and Ankylosing Spondylitis		
Preferred No PA Required (If diagnosis met) (*Must meet eligibility criteria)	Non-Preferred PA Required	First line preferred agents (preferred adalimumab products, ENBREL, and XELJANZ) may receive approval for use for FDA-labeled indications.

Adalimumab-aacf syringe	ABRILADA (adalimumab-afzb) pen, syringe	*TALTZ (ixekizumab) may receive approval for use for FDA-labeled indications following trial and failure‡ of a preferred adalimumab product or ENBREL.
Adalimumab-aaty pen	ACTEMRA (tocilizumab) syringe, Actpen	*KEVZARA (sarilumab) may receive approval for use for FDA-labeled indications following trial and failure‡ of: <ul style="list-style-type: none">A preferred adalimumab product or ENBREL ANDXELJANZ.
Adalimumab-adbm pen (IJ Kit)	Adalimumab-aacf pen	
AMJEVITA (adalimumab-atto) auto-injector, syringe	Adalimumab-adaz pen, syringe	*TYENNE (tocilizumab-aazg) may receive approval for use for FDA-labeled indications following trial and failure‡ of: <ul style="list-style-type: none">A preferred adalimumab product or ENBREL ANDXELJANZ.
CYLTEZO (adalimumab-adbm) pen, syringe	Adalimumab-adbm syringe, Crohns pen IJ Kit, PS-UV pen IJ kit	
ENBREL (etanercept)	Adalimumab-fkjp pen, syringe	<u>Quantity Limits:</u> XELJANZ IR is limited to 2 tablets per day or 60 tablets for a 30-day supply XELJANZ XR is limited to 1 tablet per day or 30 tablets for a 30-day supply
*KEVZARA (sarilumab) pen, syringe	Adalimumab-ryvk auto-injector	
*TALTZ (ixekizumab) 80 mg syringe, autoinjector	Adalimumab-aaty (2 pack) 20 mg, 40 mg syringe (Celltrion manufacturer)	<u>Non-Preferred Agents:</u>
*TYENNE (tocilizumab-aazg) pen, syringe	BIMZELX (bimekizumab-bkzx) pen	COSENTYX (secukinumab) may receive approval if meeting the following criteria: <ul style="list-style-type: none">The request meets general non-preferred criteria listed below ORThe requested drug is prescribed for treatment of enthesitis-related arthritis and meets the following:<ul style="list-style-type: none">Member is ≥ 4 years of age and weighs ≥ 15 kg ANDMember has had trialed and failed‡ NSAID therapy and either ENBREL or a preferred adalimumab product.
XELJANZ IR (tofacitinib) tablet	CIMZIA (certolizumab pegol) syringe, vial	HUMIRA brand and non-preferred adalimumab agents may receive approval if meeting the following: <ul style="list-style-type: none">The request meets one of the following:<ul style="list-style-type: none">The prescribed agent is a preferred adalimumab product ORIf the prescribed agent is brand Humira or a non-preferred adalimumab product, then the member has trialed and failed at least one preferred adalimumab product. Failure is defined as lack of efficacy or intolerable side effects with the preferred adalimumab product.
XELJANZ XR (tofacitinib ER) tablet	COSENTYX (secukinumab) syringe, pen-injector	AND <ul style="list-style-type: none">The general non-preferred criteria listed below are met.
YUFLYMA (adalimumab-aaty) auto-injector, syringe	HADLIMA (adalimumab-bwwd) Pushtouch, syringe	KINERET (anakinra) may receive approval for: <ul style="list-style-type: none">Treatment of systemic juvenile idiopathic arthritis (sJIA) or Adult-Onset Still’s Disease (AOSD) ORTreatment of rheumatoid arthritis following trial and failure‡ of<ul style="list-style-type: none">A preferred adalimumab product or ENBREL ANDXELJANZ
	HULIO (adalimumab-fkjp) pen, syringe	
	HUMIRA (adalimumab)	
	HYRIMOZ (adalimumab-adaz) pen, syringe	
	IDACIO (adalimumab-aacf) pen, syringe	ILARIS (canakinumab) may receive approval if meeting the following: <ul style="list-style-type: none">Medication is being prescribed for systemic juvenile idiopathic arthritis (sJIA) or Adult-Onset Still’s Disease (AOSD) ANDThe request meets general non-preferred drug criteria listed below.
	ILARIS (canakinumab) vial	
	KINERET (anakinra) syringe	

	<div>OLUMIANT (baricitinib) tablet</div> <div>ORENCIA (abatacept) clickject, syringe</div> <div>RINVOQ (upadacitinib), solution, tablet</div> <div>SIMLANDI (adalimumab-ryvk) auto-injector</div> <div>SIMPONI (golimumab) pen, syringe</div> <div>SKYRIZI (risankizumab-rzaa) OnBody, SC pen, syringe</div> <div>XELJANZ (tofacitinib) solution</div> <div>YUSIMRY (adalimumab-aqvh) pen</div> <div><i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i></div>	<div>Quantity Limit: 300mg (2mL) every 4 weeks</div> <div>XELJANZ (tofacitinib) oral solution may be approved with verification that the member cannot swallow a tofacitinib tablet</div> <div>All other non-preferred agents may receive approval for FDA-labeled indications following trial and failure‡ of the following preferred agents when FDA-indicated or having strong evidence supporting use for the prescribed indication from clinically recognized guideline compendia:<ul style="list-style-type: none">Adalimumab or ENBREL ANDXELJANZ ANDTYENNE, KEVZARA, or TALTZ</div> <div>Non-preferred agents that are being prescribed per FDA labeling to treat non-radiographic axial spondyloarthritis (nr-axSpA) will require trial and failure‡ of preferred agents that are FDA-labeled for treating an axial spondyloarthritis condition, including ankylosing spondylitis (AS) or nr-axSpA.</div> <div><u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent that does not have a preferred biosimilar may receive approval for continuation of therapy with the prescribed agent.</div> <div>‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction. Note that trial and failure of preferred TNF inhibitors will not be required when prescribed to treat polyarticular juvenile idiopathic arthritis (pJIA) in members with documented clinical features of lupus.</div>
Psoriatic Arthritis		
<div>Preferred No PA Required (If diagnosis met) (*Must meet eligibility criteria)</div> <div>Adalimumab-aacf syringe</div> <div>Adalimumab-aaty pen</div> <div>Adalimumab-adbm pen (IJ Kit)</div> <div>AMJEVITA (adalimumab-atto) auto-injector, syringe</div> <div>CYLTEZO (adalimumab-adbm) pen, syringe</div> <div>ENBREL (etanercept)</div> <div>*IMULDOSA (ustekinumab-SRLF) syringe</div> <div>*OTEZLA (apremilast) tablet</div>	<div>Non-Preferred PA Required</div> <div>ABRILADA (adalimumab-afzb) pen, syringe</div> <div>Adalimumab-aacf pen</div> <div>Adalimumab-adaz pen, syringe</div> <div>Adalimumab-adbm syringe, Crohns pen IJ Kit, PS-UV pen IJ kit</div> <div>Adalimumab-fkjp pen, syringe</div> <div>Adalimumab-ryvk auto-injector</div> <div>Adalimumab-aaty (2 pack) 20 mg, 40 mg syringe (Celltrion manufacturer)</div>	<div>First line preferred agents (preferred adalimumab products, ENBREL, XELJANZ) may receive approval for psoriatic arthritis indication.</div> <div>*OTEZLA (apremilast) may receive approval for psoriatic arthritis indication following trial and failure‡ of:<ul style="list-style-type: none">A preferred adalimumab product or ENBREL ANDXELJANZ</div> <div>*TALTZ (ixekizumab) may receive approval for psoriatic arthritis indication following trial and failure‡ of:<ul style="list-style-type: none">A preferred adalimumab product or ENBREL ANDXELJANZ</div> <div>*USTEKINUMAB preferred products (IMULDOSA, SELARSDI, STEQEYMA) may receive approval for psoriatic arthritis indication following trial and failure‡ of:<ul style="list-style-type: none">A preferred adalimumab product or ENBREL ANDXELJANZ.</div> <div><u>Quantity Limits:</u> XELJANZ IR is limited to 2 tablets per day or 60 tablets for a 30-day supply XELJANZ XR is limited to 1 tablet per day or 30 tablets for a 30-day supply</div>

*SELARSDI (ustekinumab-AEKN) syringe	BIMZELX (bimekizumab-bkzx) pen	<u>Non-Preferred Agents:</u>
*STEQEYMA (ustekinumab-stba) syringe	CIMZIA (certolizumab pegol) syringe, vial	COSENTYX (secukinumab) may receive approval for psoriatic arthritis indication for members ≥ 2 years of age and weighing ≥ 15 kg following trial and failure‡ of:
*TALTZ (ixekizumab) 80 mg syringe	COSENTYX (secukinumab) syringe, pen-injector	<ul style="list-style-type: none">• A preferred adalimumab product or ENBREL AND• XELJANZ AND• TALTZ or OTEZLA or a preferred ustekinumab product
XELJANZ IR (tofacitinib) tablet	HADLIMA (adalimumab-bwwd) Pushtouch, syringe	HUMIRA brand and non-preferred adalimumab agents may receive approval if meeting the following:
XELJANZ XR (tofacitinib ER) tablet	HULIO (adalimumab-fkjp) pen, syringe	<ul style="list-style-type: none">• The request meets one of the following:<ul style="list-style-type: none">○ The prescribed agent is a preferred adalimumab product OR○ If the prescribed agent is brand Humira or a non-preferred adalimumab product, then the member has trialed and failed at least one preferred adalimumab product. Failure is defined as lack of efficacy or intolerable side effects with the preferred adalimumab product.
YUFLYMA (adalimumab-aaty) auto-injector, syringe	HUMIRA (adalimumab)	AND
	HYRIMOZ (adalimumab-adaz) pen, syringe	<ul style="list-style-type: none">• The general non-preferred criteria listed below are met.
	IDACIO (adalimumab-aacf) pen, syringe	STELARA brand and non-preferred ustekinumab agents may receive approval if meeting the following:
	ORENCIA (abatacept) syringe, clickject	<ul style="list-style-type: none">• The request meets one of the following:<ul style="list-style-type: none">○ The prescribed agent is a preferred ustekinumab product OR○ If the prescribed agent is brand Stelara or a non-preferred ustekinumab product, then the member has trialed and failed at least one preferred ustekinumab product. Failure is defined as lack of efficacy or intolerable side effects with the preferred ustekinumab product.
	OTULFI (ustekinumab-aaуз) syringe	AND
	PYZCHIVA (ustekinumab-ttwe) syringe	<ul style="list-style-type: none">• The general non-preferred criteria listed below are met.
	RINVOQ (upadacitinib) tablet	XELJANZ (tofacitinib) XR approval will require verification of the clinically relevant reason for use of the XELJANZ XR formulation versus the XELJANZ IR formulation, in addition to meeting non-preferred criteria listed below.
	RINVOQ LQ (upadacitinib) solution	All other non-preferred agents may receive approval for psoriatic arthritis following trial and failure‡ of:
	SIMLANDI (adalimumab-ryvk) auto-injector	<ul style="list-style-type: none">• A preferred adalimumab product or ENBREL AND• Two other preferred products (XELJANZ, TALTZ, OTEZLA, ustekinumab)
	SIMPONI (golimumab) pen, syringe	‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction.
	SKYRIZI (risankizumab-rzaa) OnBody, pen, syringe	<u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent may receive approval for continuation of therapy with the prescribed agent.
	STELARA (ustekinumab) syringe	

	<div>TREMFYA (guselkumab) pen, injector, syringe</div> <div>Ustekinumab (generic Stelara, TTWE, AEKN) syringe, vial</div> <div>WEZLANA (ustekinumab-auub) syringe, vial</div> <div>XELJANZ (tofacitinib) solution</div> <div>YESINTEK (ustekinumab-kfce) syringe, vial</div> <div>YUSIMRY (adalimumab-aqvh) pen</div> <div><i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i></div>	
Plaque Psoriasis		
<div>Preferred No PA Required (If diagnosis met) (*Must meet eligibility criteria)</div> <div>Adalimumab-aacf syringe</div> <div>Adalimumab-aaty pen</div> <div>Adalimumab-adbm pen (IJ Kit)</div> <div>AMJEVITA (adalimumab-atto) auto-injector, syringe</div> <div>CYLTEZO (adalimumab-adbm) pen, syringe</div> <div>ENBREL (etanercept)</div> <div>*IMULDOSA (ustekinumab-SRLF) syringe</div> <div>*OTEZLA (apremilast) tablet</div> <div>*SELARSDI (ustekinumab-AEKN) syringe</div> <div>*STEQEYMA (ustekinumab-stba) syringe</div> <div>*TALTZ (ixekizumab) 80 mg syringe</div>	<div>Non-Preferred PA Required</div> <div>ABRILADA (adalimumab-afzb) pen, syringe</div> <div>Adalimumab-aacf pen</div> <div>Adalimumab-adaz pen, syringe</div> <div>Adalimumab-adbm syringe, Crohns pen IJ Kit, PS-UV pen IJ kit</div> <div>Adalimumab-fkjp pen, syringe</div> <div>Adalimumab-ryvk auto-injector</div> <div>Adalimumab-aaty (2 pack) 20 mg, 40 mg syringe (Celltrion manufacturer)</div> <div>BIMZELX (bimekizumab-bkzx) pen</div> <div>CIMZIA (certolizumab pegol) syringe, vial</div>	<div>First line preferred agents (preferred adalimumab products, ENBREL) may receive approval for plaque psoriasis indication.</div> <div>*Second line preferred agents (TALTZ, OTEZLA, preferred ustekinumab products) may receive approval for plaque psoriasis indication following trial and failure‡ of a preferred adalimumab product OR ENBREL.</div> <div><u>Non-Preferred Agents:</u></div> <div>HUMIRA brand and non-preferred adalimumab agents may receive approval if meeting the following:<ul style="list-style-type: none">The request meets one of the following:<ul style="list-style-type: none">The prescribed agent is a preferred adalimumab product ORIf the prescribed agent is brand Humira or a non-preferred adalimumab product, then the member has trialed and failed at least one preferred adalimumab product. Failure is defined as lack of efficacy or intolerable side effects with the preferred adalimumab product.</div> <div>AND<ul style="list-style-type: none">The general non-preferred criteria listed below are met.</div> <div>STELARA brand and non-preferred ustekinumab agents may receive approval if meeting the following:<ul style="list-style-type: none">The request meets one of the following:<ul style="list-style-type: none">The prescribed agent is a preferred ustekinumab product ORIf the prescribed agent is brand Stelara or a non-preferred ustekinumab product, then the member has trialed and failed at least one preferred ustekinumab product. Failure is defined as lack of efficacy or intolerable side effects with the preferred ustekinumab product.</div>

YUFLYMA (adalimumab-aaty) auto-injector, syringe	COSENTYX (secukinumab) syringe, pen-injector	AND <ul style="list-style-type: none">The general non-preferred criteria listed below are met. <p>All other non-preferred agents may receive approval for plaque psoriasis indication following trial and failure‡ of one indicated first line agent (a preferred adalimumab product, ENBREL) AND two second line agents (TALTZ, OTEZLA, or a preferred ustekinumab product).</p> <p>‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p><u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent that does not have a preferred biosimilar may receive approval for continuation of therapy with the prescribed agent.</p>
	HADLIMA (adalimumab-bwwd) Pushtouch, syringe	
	HULIO (adalimumab-fkjp) pen, syringe	
	HYRIMOZ (adalimumab-adaz) pen, syringe	
	HUMIRA (adalimumab)	
	IDACIO (adalimumab-aacf) pen, syringe	
	IMULDOSA (ustekinumab-SRLF) syringe, vial	
	OTULFI (ustekinumab-aaуз) syringe	
	PYZCHIVA (ustekinumab-ttwe) syringe	
	SELARSDI (ustekinumab-AEKN) syringe	
	SILIQ (brodalumab) syringe	
	SIMLANDI (adalimumab-ryvk) auto-injector	
	SKYRIZI (risankizumab-rzaa) OnBody, pen, syringe	
	SOTYKTU (ducravacitinib) oral tablet	
	STELARA (ustekinumab) syringe	

	STEQEYMA (ustekinumab-stba) syringe	
	TALTZ (ixekizumab) 20mg, 40mg syringe	

	<p>TREMFYA (guselkumab) injector, syringe</p> <p>Ustekinumab (generic Stelara, TTWE, AEKN) syringe, vial</p> <p>WEZLANA (ustekinumab-auub) syringe, vial</p> <p>YESINTEK (ustekinumab-kfce) syringe, vial</p> <p>YUSIMRY (adalimumab-aqvh) pen</p> <p><i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i></p>	
Crohn’s Disease and Ulcerative Colitis		
<p>Preferred No PA Required (If diagnosis met) (*Must meet eligibility criteria)</p> <p>Adalimumab-aacf syringe</p> <p>Adalimumab-aaty pen</p> <p>Adalimumab-adbm pen (IJ Kit)</p> <p>AMJEVITA (adalimumab-atto) auto-injector, syringe</p> <p>CYLTEZO (adalimumab-adbm) pen, syringe</p> <p>*IMULDOSA (ustekinumab-SRLF) syringe</p> <p>*SELARSDI (ustekinumab-AEKN) syringe</p> <p>*STEQEYMA (ustekinumab-stba) syringe</p> <p>*XELJANZ IR (tofacitinib) tablet</p> <p>*XELJANZ XR (tofacitinib ER) tablet</p> <p>YUFLYMA (adalimumab-aaty) auto-injector, syringe</p>	<p>Non-Preferred PA Required</p> <p>ABRILADA (adalimumab-afzb) pen, syringe</p> <p>Adalimumab-aacf pen</p> <p>Adalimumab-adaz pen, syringe</p> <p>Adalimumab-adbm syringe, Crohns pen IJ Kit, PS-UV pen IJ kit</p> <p>Adalimumab-fkjp pen, syringe</p> <p>Adalimumab-ryvk auto-injector</p> <p>Adalimumab-aaty (2 pack) 20 mg, 40 mg syringe (Celltrion manufacturer)</p> <p>CIMZIA (certolizumab pegol) syringe, vial</p> <p>COSENTYX (secukinumab) syringe, pen-injector</p>	<p>First and second line preferred agents (preferred adalimumab products, preferred ustekinumab products, XELJANZ) may receive approval for Crohn’s disease and ulcerative colitis indications.</p> <p><u>Quantity Limits:</u> XELJANZ IR is limited to 2 tablets per day or 60 tablets for a 30-day supply XELJANZ XR is limited to 1 tablet per day or 30 tablets for a 30-day supply</p> <p><u>Non-Preferred Agents:</u> HUMIRA brand and non-preferred adalimumab agents may receive approval if meeting the following:</p> <ul style="list-style-type: none">• The request meets one of the following:<ul style="list-style-type: none">○ The prescribed agent is a preferred adalimumab product OR○ If the prescribed agent is brand Humira or a non-preferred adalimumab product, then the member has trialed and failed at least one preferred adalimumab product. Failure is defined as lack of efficacy or intolerable side effects with the preferred adalimumab product. <p>AND</p> <ul style="list-style-type: none">• The general non-preferred criteria listed below are met. <p>STELARA brand and non-preferred ustekinumab agents may receive approval if meeting the following:</p> <ul style="list-style-type: none">• The request meets one of the following:<ul style="list-style-type: none">○ The prescribed agent is a preferred ustekinumab product OR○ If the prescribed agent is brand Stelara or a non-preferred ustekinumab product, then the member has trialed and failed at least one preferred ustekinumab product. Failure is defined as lack of efficacy or intolerable side effects with the preferred ustekinumab product. <p>AND</p> <ul style="list-style-type: none">• The general non-preferred criteria listed below are met.

	<div>ENTYVIO (vedolizumab) pen</div> <div>HADLIMA (adalimumab-bwwd) Pushtouch, syringe</div> <div>HULIO (adalimumab-fkjp) syringe</div> <div>HUMIRA (adalimumab)</div> <div>HYRIMOZ (adalimumab-adaz) pen, syringe</div> <div>IDACIO (adalimumab-aacf) pen, syringe</div> <div>OLUMIANT (baricitinib) tablet</div> <div>OMVOH (mirikizumab-mrkz) pen</div> <div>OTULFI (ustekinumab-aauz) syringe</div> <div>PYZCHIVA (ustekinumab-ttwe) syringe</div> <div>RINVOQ (upadacitinib) tablet</div> <div>RINVOQ LQ (upadacitinib) solution</div> <div>SIMLANDI (adalimumab-ryvk) auto-injector</div> <div>SIMPONI (golimumab) pen, syringe</div> <div>SKYRIZI (risankizumab-rzaa) OnBody, pen, syringe</div> <div>STELARA (ustekinumab) syringe, vial</div> <div>Ustekinumab (generic Stelara, TTWE, AEKN) syringe, vial</div> <div>VELSIPITY (etrasimod) tablet</div>	<p>All other non-preferred agents may receive approval for FDA-labeled indications if meeting the following:</p> <ul style="list-style-type: none">• The requested medication is being prescribed for treating moderately-to-severely active Crohn’s disease or moderately-to-severely active Ulcerative Colitis in alignment with indicated use outlined in FDA-approved product labeling AND• The requested medication meets FDA-labeled indicated age for prescribed use AND• For treatment of moderately-to-severely active Crohn’s disease, member has trial and failure‡ of one preferred adalimumab product OR for treatment of moderately-to-severely active ulcerative colitis, member has trial and failure‡ of one preferred adalimumab product AND one preferred ustekinumab product• For treatment of moderately-to-severely active ulcerative colitis, member has trial and failure‡ of:<ul style="list-style-type: none">○ One preferred adalimumab product or XELJANZ AND○ One preferred ustekinumab product. <p><u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent that does not have a preferred biosimilar may receive approval for continuation of therapy with the prescribed agent.</p> <p>‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction. Note that trial and failure of Xeljanz (tofacitinib) will not be required when prescribed for ulcerative colitis for members ≥ 50 years of age that have an additional CV risk factor.</p>
--	---	---

	<div>WEZLANA (ustekinumab-auub) syringe, vial</div> <div>XELJANZ (tofacitinib) solution</div> <div>YESINTEK (ustekinumab-kfce) syringe, vial</div> <div>YUSIMRY (adalimumab-aqvh) pen</div> <div>ZYMFENTRA (infliximab-dyyb) pen kit, syringe kit</div> <div><i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i></div>	
Asthma		
<div>Preferred PA Required (*Must meet eligibility criteria)</div> <div>*DUPIXENT (dupilumab) pen, syringe</div> <div>*FASENRA (benralizumab) pen, syringe</div> <div>*TEZSPIRE (tezepelumab-ekko) pen</div> <div>*XOLAIR (omalizumab) syringe, autoinjector</div>	<div>Non-Preferred PA Required</div> <div>NUCALA (mepolizumab) auto-injector, syringe</div> <div><i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i></div>	<div>*Preferred products (Dupixent, Fasenra, Tezspire, Xolair) may receive approval if meeting the following:</div> <div>DUPIXENT (dupilumab):</div> <div><ul style="list-style-type: none">Member is 6 years of age or older ANDMember has an FDA-labeled indicated use for treating one of the following:<ul style="list-style-type: none">Moderate to severe asthma (on medium to high dose inhaled corticosteroid and a long-acting beta agonist) with eosinophilic phenotype based on a blood eosinophil level of $\geq 150/\text{mcL}$ OROral corticosteroid dependent asthmaAND<ul style="list-style-type: none">Member’s asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies ANDMedication is being prescribed as add-on therapy to existing asthma regimen.<div>Quantity Limit: 2 syringes every 28 days after initial 14 days of therapy (first dose is twice the regular scheduled dose)</div></div> <div>FASENRA (benralizumab):</div> <div><ul style="list-style-type: none">Member is ≥ 6 years of age ANDMember has an FDA-labeled indicated use for treating severe asthma with an eosinophilic phenotype based on a blood eosinophil level of $\geq 150/\text{mcL}$ ANDMember’s asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies ANDThe requested medication is being prescribed as add-on therapy to existing asthma regimen.<div>Quantity Limit: One 30 mg unit dose pack every 28 days for the first 3 doses and then every 8 weeks thereafter</div></div> <div>TEZSPIRE (tezepelumab-ekko):</div> <div><ul style="list-style-type: none">Member is ≥ 12 years of age ANDMember has a diagnosis of severe asthma AND</div>

		<ul style="list-style-type: none">Member’s asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies ANDThe requested medication is being prescribed as add-on therapy to existing asthma regimen. <p><u>Quantity Limit:</u> Four 210 mg unit dose packs every 28 days</p> <p>XOLAIR (omalizumab) may receive approval if meeting the following based on prescribed indication:</p> <ul style="list-style-type: none">Member is ≥ 6 years of age ANDMember has an FDA-labeled indicated use for treating asthma ANDMember has a positive skin test or in vitro reactivity to a perennial inhaled allergen or has a pre-treatment IgE serum concentration ≥ 30 IU/mL ANDMember’s asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies ANDThe requested medication is being prescribed as add-on therapy to existing asthma regimen. <p><u>Non-Preferred Agents:</u></p> <p>Non-preferred FDA-indicated biologic agents for asthma may receive approval if meeting the following:</p> <ul style="list-style-type: none">The requested medication is being prescribed for treating asthma in alignment with indicated use outlined in FDA-approved product labeling (including asthma type and severity) ANDIf prescribed for use for asthma with eosinophilic phenotype, member has a blood eosinophil count ≥ 150 cells/mcL ANDThe requested medication meets FDA-labeled indicated age for prescribed use ANDMember’s asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies ANDThe requested medication is being prescribed as add-on therapy to existing asthma regimen ANDMember has trialed and failed‡ two preferred agents. <p><u>Quantity Limits:</u> Non-preferred medications will be subject to quantity limitations in alignment with FDA-approved dosing per product package labeling. Nucala (mepolizumab) is limited to 100mg every 4 weeks (members ≥ 12 years of age) or 40mg every 4 weeks (members 6-11 years of age).</p> <p>‡Failure is defined as a lack of efficacy, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions.</p> <p><u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent that does not have a preferred biosimilar may receive approval for continuation of therapy with the prescribed agent.</p>
Atopic Dermatitis		
Preferred	Non-Preferred PA Required	*Preferred products (Adbry and Dupixent) may receive approval if meeting the following:

<p>(*Must meet eligibility criteria)</p> <p>*ADBRY (tralokinumab-ldrm) syringe, autoinjector</p> <p>*DUPIXENT (dupilumab) pen, syringe</p>	<p>CIBINQO (abrocitinib) tablet</p> <p>RINVOQ (upadacitinib) tablet</p> <p><i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i></p>	<p>ADBRY (tralokinumab-ldrm):</p> <ul style="list-style-type: none">• The requested drug is being prescribed for moderate-to-severe atopic dermatitis AND• Member has trialed and failed‡ the following agents:<ul style="list-style-type: none">○ One medium potency to very-high potency topical corticosteroid (such as mometasone furoate, betamethasone dipropionate) AND○ One topical calcineurin inhibitor (such as pimecrolimus or tacrolimus) <p><u>Maximum Dose:</u> 600 mg/2 weeks</p> <p><u>Quantity Limit:</u> Four 150 mg/mL prefilled syringes/2 weeks</p> <p>DUPIXENT (dupilumab):</p> <ul style="list-style-type: none">• Member has a diagnosis of moderate to severe atopic dermatitis AND• Member has trialed and failed‡ the following agents:<ul style="list-style-type: none">○ One medium potency to very-high potency topical corticosteroid [such as mometasone furoate, betamethasone dipropionate, or fluocinonide (see PDL for list of preferred products) AND○ One topical calcineurin inhibitor (such as pimecrolimus or tacrolimus) <p><u>Quantity Limit:</u> 2 syringes every 28 days after initial 14 days of therapy (first dose is twice the regular scheduled dose)</p> <p><u>Non-Preferred Agents:</u></p> <p>Non-preferred agents indicated for the treatment of atopic dermatitis may receive approval if meeting the following:</p> <ul style="list-style-type: none">• Member has a diagnosis of moderate to severe chronic atopic dermatitis AND• Member has trialed and failed‡ all of the following agents:<ul style="list-style-type: none">○ One medium potency to very-high potency topical corticosteroid (such as mometasone furoate, betamethasone dipropionate, or fluocinonide) AND○ One topical calcineurin inhibitor (such as pimecrolimus and tacrolimus) AND○ Opzelura (ruxolitinib) topical cream• Member has trialed and failed‡ therapy with two preferred agents (ADBRY and DUPIXENT) for the prescribed indication• The medication is being prescribed by or in consultation with a dermatologist, allergist, immunologist, or rheumatologist <p><u>Approval:</u> One year</p> <p>‡Failure is defined as a lack of efficacy, allergy, intolerable side effects, contraindication , or significant drug-drug interaction.</p> <p><u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent that does not have a preferred biosimilar may receive approval for continuation of therapy with the prescribed agent.</p>
Other indications		

Preferred (If diagnosis met, No PA required) (Must meet eligibility criteria*)	Non-Preferred PA Required	<u>Preferred Agents:</u>
CYLTEZO (adalimumab-adbm) pen, syringe	ACTEMRA (tocilizumab) syringe, Actpen	*DUPIXENT (dupilumab) may receive approval if meeting the following based on prescribed indication:
*DUPIXENT (dupilumab) pen, syringe	ARCALYST (rilonacept) injection	<u>Bullous Pemphigoid</u>
ENBREL (etanercept)	CIMZIA (certolizumab pegol) syringe	<ul style="list-style-type: none">Member is 18 years of age or older ANDMember is diagnosed with bullous pemphigoid ANDMember has trialed and failed‡ one of the following therapies:<ul style="list-style-type: none">High-potency topical corticosteroidOral prednisoneDoxycycline
*FASENRA (benralizumab) pen, syringe	COSENTYX (secukinumab) syringe, pen-injector	<u>Chronic Spontaneous Urticaria</u>
HUMIRA (adalimumab)	ILARIS (canakinumab) vial	<ul style="list-style-type: none">Member is 12 years of age or older ANDMember is diagnosed with chronic spontaneous urticaria ANDMember is symptomatic despite H1 antihistamine treatment ANDMember has tried and failed‡ at least three of the following<ul style="list-style-type: none">High-dose second generation H1 antihistamineH2 antihistamineFirst-generation antihistamineLeukotriene receptor antagonistHydroxyzine or doxepin
*KEVZARA (sarilumab)	KINERET (anakinra) syringe	<u>Chronic Obstructive Pulmonary Disease</u>
*OTEZLA (apremilast) tablet	NUCALA (mepolizumab) auto-injector, syringe	<ul style="list-style-type: none">Member is ≥ 18 years of age ANDMedication is being prescribed by or in consultation with a pulmonologist or allergist ANDRequested medication is being prescribed as an add-on maintenance treatment for inadequately controlled chronic obstructive pulmonary disease (COPD) ANDMember’s COPD is an eosinophilic phenotype based on a blood eosinophil level of ≥ 150 cells/mcL ANDMember is receiving, and will continue, standard maintenance triple therapy for COPD (inhaled corticosteroid, long-acting muscarinic agent, long-acting beta agonist) as recommended by the current Global Initiative for Chronic Obstructive Lung Disease (GOLD) guidelines ANDMember has experienced at least 2 moderate OR 1 severe COPD exacerbation during the past 12 months
*TYENNE (tocilizumab-aazg)	OLUMIANT (baricitinib) tablet	<u>Chronic Rhinosinusitis with Nasal Polyposis</u>
XELJANZ IR (tofacitinib) tablet	<i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i>	<ul style="list-style-type: none">Member is ≥ 12 years of age ANDMedication is being prescribed as an add-on maintenance treatment for inadequately controlled chronic rhinosinusitis with nasal polyposis (CRSwNP) ANDMember has trialed and failed‡ therapy with at least two intranasal corticosteroid regimens
XELJANZ XR (tofacitinib ER) tablet		<u>Eosinophilic Esophagitis (EoE):</u>
*XOLAIR (omalizumab) syringe, autoinjector		<ul style="list-style-type: none">Member is ≥ 1 year of age ANDMember weighs at least 15 kg ANDMember has a diagnosis of eosinophilic esophagitis (EoE) with ≥ 15 intraepithelial eosinophils per high-power field (eos/hpf), with or without a history of esophageal dilations AND
YUFLYMA (adalimumab-aaty) auto-injector		

		<ul style="list-style-type: none">• Member is following appropriate dietary therapy interventions AND• Medication is being prescribed by or in consultation with a gastroenterologist, allergist or immunologist AND• Member has trialed and failed‡ one of the following treatment options for EoE:<ul style="list-style-type: none">○ Proton pump inhibitor trial of at least eight weeks in duration if reflux is a contributing factor OR○ Minimum four-week trial of local therapy with a corticosteroid medication <p><u>Prurigo Nodularis:</u></p> <ul style="list-style-type: none">• Member is ≥ 18 years of age AND• Medication is being prescribed as treatment for prurigo nodularis AND• Member has trialed and failed‡ therapy with at least two corticosteroid regimens (topical or intralesional injection). <p>*FASENRA (benralizumab) may be approved for the treatment of adult patients with eosinophilic granulomatosis with polyangiitis (EGPA).</p> <p>*KEVZARA (sarilumab) treatment of adult patients with polymyalgia rheumatica who have had an inadequate response to corticosteroids or who cannot tolerate corticosteroid taper.</p> <p>*OTEZLA (apremilast) treatment of adult patients with oral ulcers associated with Behçet's Disease.</p> <p>*TYENNE (tocilizumab-aazg) may receive approval for use for FDA-label indications following trial and failure‡ of a preferred adalimumab product or ENBREL</p> <p>*XOLAIR (omalizumab) may receive approval if meeting the following based on prescribed indication:</p> <p><u>Chronic Rhinosinusitis with Nasal Polyps:</u></p> <ul style="list-style-type: none">• Member is 18 years of age or older AND• Medication is being prescribed as add-on maintenance treatment of chronic rhinosinusitis with nasal polyps (CRSwNP) in adult patients 18 years of age and older with inadequate response to nasal corticosteroids AND• Member has tried and failed‡ therapy with at least two intranasal corticosteroid regimens <p><u>Chronic Spontaneous Urticaria:</u></p> <ul style="list-style-type: none">• Member is 12 years of age or older AND• Member is diagnosed with chronic idiopathic urticaria AND• Member is symptomatic despite H1 antihistamine treatment AND• Member has tried and failed‡ at least three of the following:<ul style="list-style-type: none">○ High-dose second generation H1 antihistamine○ H2 antihistamine○ First-generation antihistamine○ Leukotriene receptor antagonist○ Hydroxyzine or doxepin (must include) <p>AND</p>
--	--	---

		<ul style="list-style-type: none">• Prescriber attests that the need for continued therapy will be periodically reassessed (as the appropriate duration of Xolair therapy for CIU has currently not been evaluated). <p><u>IgE-Mediated Food Allergy:</u></p> <ul style="list-style-type: none">• Medication is being prescribed for reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods in adult and pediatric patients aged 1 year and older with IgE-mediated food allergy. <p>All other preferred agents may receive approval for use for FDA-labeled indications.</p> <p><u>Non-Preferred Agents:</u></p> <p>ARCALYST (rilonacept) may receive approval if meeting the following:</p> <ul style="list-style-type: none">• Medication is being prescribed for one of the following autoinflammatory periodic fever syndromes (approval for all other indications is subject to meeting non-preferred criteria listed below):<ul style="list-style-type: none">○ Cryopyrin-associated Autoinflammatory Syndrome (CAPS), including:<ul style="list-style-type: none">▪ Familial Cold Autoinflammatory Syndrome (FCAS)▪ Muckle-Wells Syndrome (MWS)○ Maintenance of remission of Deficiency of Interleukin-1 Receptor Antagonist (DIRA) in adults and pediatric patients weighing at least 10 kg○ Treatment of recurrent pericarditis and reduction in risk of recurrence in adults and children ≥ 12 years of age<p>AND</p><ul style="list-style-type: none">• Member has trialed and failed‡ colchicine AND• Initial approval will be given for 12 weeks and authorization approval for continuation will be provided based on clinical response.<p>ILARIS (canakinumab) may receive approval if meeting the following:</p><ul style="list-style-type: none">• Medication is being prescribed for one of the following (approval for all other indications is subject to meeting non-preferred criteria listed below):<ul style="list-style-type: none">○ Familial Mediterranean Fever (FMF)○ Hyperimmunoglobulinemia D syndrome (HIDS)○ Mevalonate Kinase Deficiency (MKD)○ Neonatal onset multisystem inflammatory disease (NOMID)○ TNF Receptor Associated Periodic Syndrome (TRAPS)○ Cryopyrin-associated Autoinflammatory Syndrome (including Familial Cold Autoinflammatory Syndrome and Muckle-Wells Syndrome)○ Symptomatic treatment of adult patients with gout flares in whom NSAIDs and colchicine are contraindicated, are not tolerated, or do not provide an adequate response, and in whom repeated courses of corticosteroids are not appropriate (limited to four 150mg doses per one year approval)<p>AND</p><ul style="list-style-type: none">• Member has trialed and failed‡ colchicine.
--	--	--

		<ul style="list-style-type: none">Quantity Limits:<ul style="list-style-type: none">Cryopyrin-associated periodic syndrome: 600mg (4mL) every 8 weeksAll other indications: 300mg (2mL) every 4 weeks <p>KINERET (anakinra) may receive approval if meeting the following:</p> <ul style="list-style-type: none">Medication is being prescribed for one of the following indications (approval for all other indications is subject to meeting non-preferred criteria below):<ul style="list-style-type: none">Neonatal onset multisystem inflammatory disease (NOMID).Familial Mediterranean Fever (FMF) <p>AND</p> <ul style="list-style-type: none">Member has trialed and failed‡ colchicine. <p>NUCALA (mepolizumab) may receive approval if meeting the following based on prescribed indication (for any FDA-labeled indications in this subclass category that are not listed, approval is subject to meeting non-preferred criteria listed below):</p> <p><u>Maintenance Treatment of COPD:</u></p> <ul style="list-style-type: none">Member is 18 years of age or older ANDRequested medication is being prescribed as an add-on maintenance treatment for inadequately controlled chronic obstructive pulmonary disease (COPD) ANDMember’s COPD is an eosinophilic phenotype based on a blood eosinophil level of ≥300 cells/mcL ANDMedication is being prescribed by or in consultation with a pulmonologist or allergist ANDMember is receiving, and will continue, standard maintenance triple therapy for COPD (long-acting beta agonist, long-acting muscarinic agent, inhaled corticosteroid) ANDMember has experienced at least 2 moderate COPD exacerbations OR 1 severe exacerbation during the past 12 months ANDMember has trialed and failed‡ therapy with Dupixent (dupilumab). <p><u>Chronic Rhinosinusitis with Nasal Polyps:</u></p> <ul style="list-style-type: none">Member is 18 years of age or older ANDMedication is being prescribed as an add-on maintenance treatment in adult patients with inadequately controlled chronic rhinosinusitis with nasal polyposis (CRSwNP) ANDMember has a baseline bilateral endoscopic nasal polyps score (NPS; scale 0-8) AND nasal congestion/obstruction score (NC; scale 0-3) averaged over 28-day period ANDMember has trialed and failed‡ therapy with three intranasal corticosteroids (see PDL Class) ANDMedication is being prescribed by or in consultation with a rheumatologist, allergist, ear/nose/throat specialist or pulmonologist ANDInitial authorization will be for 24 weeks, for additional 12-month approval member must meet the following criteria:<ul style="list-style-type: none">NC and NPS scores are provided and show a 20% reduction in symptoms from baseline ANDMember continues to use primary therapies such as intranasal corticosteroids. <p><u>Eosinophilic Granulomatosis with polyangiitis (EGPA):</u></p>
--	--	---

		<ul style="list-style-type: none">• Member is 18 years of age or older AND• Member has been diagnosed with relapsing or refractory EGPA at least 6 months prior to request as demonstrated by ALL the following:<ul style="list-style-type: none">○ Member has a diagnosis of asthma AND○ Member has a blood eosinophil count of greater than or equal to 1000 cells/mcL or a blood eosinophil level of 10%AND• Member has the presence of two of the following EGPA characteristics:<ul style="list-style-type: none">○ Histopathological evidence of eosinophilic vasculitis, perivascular eosinophilic infiltration, or eosinophil-rich granulomatous inflammation○ Neuropathy○ Pulmonary infiltrates○ Sinonasal abnormality○ Cardiomyopathy○ Glomerulonephritis○ Alveolar hemorrhage○ Palpable purpura○ Antineutrophil cytoplasmic antibody (ANCA) positiveAND• Member has trialed and failed‡ Fasenra (benralizumab) AND• Dose of NUCALA (mepolizumab) 300 mg once every 4 weeks is being prescribed. <p><u>Hypereosinophilic Syndrome (HES):</u></p> <ul style="list-style-type: none">• Member is 12 years of age or older AND• Member has a diagnosis for HES for at least 6 months that is nonhematologic secondary HES AND• Member has a blood eosinophil count of greater than or equal to 1000 cells/mcL AND• Member has a history of two or more HES flares (defined as worsening clinical symptoms or blood eosinophil counts requiring an increase in therapy) AND <ul style="list-style-type: none">• Member has been on stable dose of HES therapy for at least 4 weeks, at time of request, including at least one of the following:<ul style="list-style-type: none">○ Oral corticosteroids○ Immunosuppressive therapy○ Cytotoxic therapyAND• Dose of 300 mg once every 4 weeks is being prescribed. <p>All other non-preferred agent indications may receive approval for FDA-labeled use following trial and failure‡ of all preferred agents that are FDA-indicated or have strong evidence supporting use for the prescribed indication from clinically recognized guideline compendia (only one preferred adalimumab product trial required).</p> <p>‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction.</p>
--	--	---

		<p><u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent that does not have a preferred biosimilar will be subject to meeting reauthorization criteria above when listed for the prescribed indication, or if reauthorization criteria are not listed for the prescribed indication, may receive approval for continuation of therapy.</p> <p><u>Note:</u> Prior authorization requests for agents prescribed solely for treating alopecia areata will not be approved.</p>
<h2>X. Miscellaneous</h2>		
Therapeutic Drug Class: EPINEPHRINE PRODUCTS – <i>Effective 1/1/2026</i>		
No PA Required AUVI-Q (epinephrine) auto-injector Epinephrine 0.15mg/0.15ml, 0.3mg/0.3ml auto-injector EPIPEN 0.3 mg/0.3 ml (epinephrine) auto-injector EPIPEN JR 0.15 mg/0.15 ml, (epinephrine) auto-injector NEFFY Spray	PA Required SYMJEPI 0.15mg/0.3ml, 0.3mg/0.3ml (epinephrine) syringe	<p>Non-preferred products may be approved if the member has failed treatment with one of the preferred products. Failure is defined as allergy to ingredients in product or intolerable side effects.</p> <p>Quantity limit: 4 single-dose units per year unless used / damaged / lost</p>
Therapeutic Drug Class: NEWER HEREDITARY ANGIOEDEMA PRODUCTS – <i>Effective 1/1/2026</i>		
PA Required for all agents in this class		<u>Medications Indicated for Routine Prophylaxis:</u>
Preferred <u>Prophylaxis:</u> HAEGARDA (C1 esterase inhibitor) vial ORLADEYO (berotralstat) oral capsule TAKHZYRO (lanadelumab-flyo) syringe, vial <u>Treatment:</u> BERINERT (C1 esterase inhibitor) kit, vial Icatibant syringe (generic FIRAZYR)	Non-Preferred <u>Prophylaxis:</u> ANDEMBRY (garadacimab-gxii) autoinjector CINRYZE (C1 esterase inhibitor) kit <u>Treatment:</u> EKTERLY (sebetralstat) tablet FIRAZYR (icatibant acetate) syringe BNR RUCONEST (C1 estera se inhibitor, recomb) vial	<p>Members are restricted to coverage of one medication for <u>routine prophylaxis</u> at one time. Prior authorization approval will be for one year.</p> <p>Preferred products for routine prophylaxis (Haegarda, Orladeyo, Takhzyro) may be approved if the following criteria are met:</p> <ul style="list-style-type: none">Member has one of the following diagnoses:<ul style="list-style-type: none">Type I HAE (Hereditary Angioedema with deficient C1-inhibitor) ORType II HAE (Hereditary Angioedema with dysfunctional C1-inhibitor) confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, C1-INH level) ORDiagnosis of HAE with normal C1-inhibitor and based on clinical presentation <p>AND</p> <ul style="list-style-type: none">Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema ANDThe request meets one of the following:<ul style="list-style-type: none">The requested product is being used for short-term prophylaxis to undergo a surgical procedure or major dental work ORThe requested product is being used for long-term prophylaxis and member meets one of the following:<ul style="list-style-type: none">History of ≥ 1 attack per month resulting in documented ED admission or hospitalization OR

- History of laryngeal attacks OR
- History of ≥ 2 attacks per month involving the face, throat, or abdomen

AND

- Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications **AND**
- The request meets minimum age and maximum dose limits listed in Table 1 **AND**
- The following criteria are met when listed for the requested medication:
 - For Haegarda (C1 esterase inhibitor), prescriber acknowledges that the member will receive information and/or counseling regarding the information from the FDA-labeled package insert outlining transmission of infectious agents with a medication made from human blood.
 - For Orladeyo (berotralstat): Appropriate drug interaction interventions will be made for members using concomitant medications that may require dose adjustments (such as cyclosporine, fentanyl, pimozide, digoxin)
 - For Takhzyro (lanadelumab-flyo), prescriber acknowledges that though the recommended starting dose is 300 mg every 2 weeks, a dosing interval of 300 mg every 4 weeks is also effective and may be considered if the patient is well-controlled (attack free) for more than 6 months.

Non-preferred products for routine prophylaxis may be approved if the following criteria are met:

- The request meets all criteria listed for preferred products above **AND**
- The member has trialed and failed at least two preferred agents indicated for routine prophylaxis. Failure is defined as lack of efficacy, allergy, intolerable side effect, or a significant drug-drug interaction.

Product Name	Minimum Age	Maximum Dose
CINRYZE (C1 esterase inhibitor-human)	6 years	2,000 units IV every 3 or 4 days
HAEGARDA (C1 esterase inhibitor-human)	6 years	60 units/kg twice weekly
ORLADEYO (berotralstat)	2 years	150 mg once daily
TAKHZYRO (lanadelumab-flyo)	2 years	300 mg every 2 weeks

Medications Indicated for Treatment of Acute Attacks:

Members are restricted to coverage of one medication for treatment of acute attacks at one time. Prior authorization approval will be for one year.

Preferred products for treatment of acute attacks (Berinert, Icatibant) may be approved if the following criteria are met:

- Member has one of the following diagnoses:
 - Type I HAE (Hereditary Angioedema with deficient C1-inhibitor) OR

		<ul style="list-style-type: none">○ Type II HAE (Hereditary Angioedema with dysfunctional C1-inhibitor) confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, C1-INH level) OR○ A diagnosis of HAE with normal C1-inhibitor based on clinical presentation <p>AND</p> <ul style="list-style-type: none">• Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND• Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications AND• The request meets minimum age and maximum dose limits listed in Table 2 AND• For Berinert (C1 esterase inhibitor): Prescriber acknowledges that the member will receive information and/or counseling regarding the information from the FDA-labeled package insert outlining transmission of infectious agents with a medication made from human blood. <p>Non-preferred products for treatment of acute attacks may be approved if the following criteria are met:</p> <ul style="list-style-type: none">• The request meets all criteria listed for preferred products above AND• The member has trialed and failed at least two preferred agents indicated for treatment of acute attacks. Failure is defined as lack of efficacy, allergy, intolerable side effect, or a significant drug-drug interaction. <p><u>Quantity limit:</u> EKTERLY (sebetralstat) limited to four 300 mg tablets (1,200 mg) per 30 days unless used, damaged, or lost.</p> <p><u>Continuation of therapy:</u> Members with previous PA approval on file for Ruconest (C1 esterase inhibitor recombinant) may receive approval for continuation of therapy.</p> <table><tr><th colspan="3">Table 2: FDA-approved Minimum Age and Maximum Dose</th></tr><tr><th>Product Name</th><th>Minimum Age</th><th>Maximum Dose</th></tr><tr><td>BERINERT (C1 esterase inhibitor)</td><td>5 years</td><td>20 units/kg</td></tr><tr><td>EKTERLY (sebetralstat)</td><td>12 years</td><td>1,200 mg/24 hours</td></tr><tr><td>FIRAZYR (icatibant acetate)</td><td>18 years</td><td>30 mg</td></tr><tr><td>RUCONEST (C1 esterase inhibitor recombinant)</td><td>13 years</td><td>4,200 Units</td></tr></table> <p>All other non-preferred agents may be approved if the member has trialed and failed at least two preferred agents with the same indicated role in therapy as the prescribed medication (prophylaxis or treatment). Failure is defined as lack of efficacy, allergy, intolerable side effects, or a significant drug-drug interaction.</p>	Table 2: FDA-approved Minimum Age and Maximum Dose			Product Name	Minimum Age	Maximum Dose	BERINERT (C1 esterase inhibitor)	5 years	20 units/kg	EKTERLY (sebetralstat)	12 years	1,200 mg/24 hours	FIRAZYR (icatibant acetate)	18 years	30 mg	RUCONEST (C1 esterase inhibitor recombinant)	13 years	4,200 Units
Table 2: FDA-approved Minimum Age and Maximum Dose																				
Product Name	Minimum Age	Maximum Dose																		
BERINERT (C1 esterase inhibitor)	5 years	20 units/kg																		
EKTERLY (sebetralstat)	12 years	1,200 mg/24 hours																		
FIRAZYR (icatibant acetate)	18 years	30 mg																		
RUCONEST (C1 esterase inhibitor recombinant)	13 years	4,200 Units																		
Therapeutic Drug Class: PHOSPHATE BINDERS – <i>Effective 10/1/2025</i>																				
No PA Required Calcium acetate capsule	PA Required AURYXIA (ferric citrate) tablet	Prior authorization for non-preferred products in this class may be approved if member meets all the following criteria: <ul style="list-style-type: none">• Member has diagnosis of end stage renal disease AND• Member has elevated serum phosphorus [> 4.5 mg/dL or > 1.46 mmol/L] AND• Provider attests to member avoidance of high phosphate containing foods from diet AND																		

PHOSLYRA (calcium acetate) solution	Calcium acetate tablet	<ul style="list-style-type: none">Member has trialed and failed‡ one preferred agent (lanthanum products require trial and failure‡ of a preferred sevelamer product). <p>Auryxia (ferric citrate) may be approved if the member meets all the following criteria:</p> <ul style="list-style-type: none">Member is diagnosed with end-stage renal disease, receiving dialysis, and has elevated serum phosphate (> 4.5 mg/dL or > 1.46 mmol/L). ANDProvider attests to counseling member regarding avoiding high phosphate containing foods from diet ANDMember has trialed and failed‡ three preferred agents with different mechanisms of action prescribed for hyperphosphatemia in end stage renal disease <p>OR</p> <ul style="list-style-type: none">Member is diagnosed with chronic kidney disease with iron deficiency anemia and is not receiving dialysis ANDMember has tried and failed‡ at least two different iron supplement product formulations (OTC or RX) <p>Velphoro (sucroferric oxyhydroxide tablet, chewable) may be approved if the member meets all of the following criteria:</p> <ul style="list-style-type: none">Member is diagnosed with chronic kidney disease and receiving dialysis and has elevated serum phosphate (> 4.5 mg/dL or > 1.46 mmol/L). ANDProvider attests to counseling member regarding avoiding high phosphate containing foods from diet ANDMember has trialed and failed‡ two preferred agents, one of which must be a preferred sevelamer product <p>Maximum Dose: Velphoro 3000mg daily</p> <p>Members currently stabilized on a non-preferred lanthanum product may receive approval to continue therapy with that product.</p> <p>‡Failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p><i>Note: Medications administered in a dialysis unit or clinic are billed through the Health First Colorado medical benefit or Medicare with members with dual eligibility.</i></p>
Sevelamer carbonate tablet, powder pack	CALPHRON (calcium acetate) tablet	
	Ferric citrate tablet	
	FOSRENOL (lanthanum carbonate) chewable tablet, powder pack	
	Lanthanum carbonate chewable tablet	
	REVELA (sevelamer carbonate) powder pack, tablet	
	Sevelamer HCl tablet	
	VELPHORO (sucroferric oxide) chewable tablet	
	XPHOZAH (tenapanor) tablet	
Therapeutic Drug Class: PRENATAL VITAMINS / MINERALS – <i>Effective 10/1/2025</i>		
<p>Preferred *Must meet eligibility criteria</p> <p>COMPLETE NATAL DHA pack</p> <p>M-NATAL PLUS tablet</p> <p>NESTABS tablets</p> <p>PRENATAL VITAMIN PLUS LOW IRON tablet (<i>Patrin Pharma only</i>)</p> <p>SE-NATAL 19 chewable tablet^{BNR}</p> <p>TARON-C DHA capsule</p>	<p>Non-Preferred PA Required</p> <p>All other rebateable prescription products are non-preferred</p>	<p>*Preferred and non-preferred prenatal vitamin products are a benefit for members from 11-60 years of age who are pregnant, lactating, or trying to become pregnant.</p> <p>Prior authorization for non-preferred agents may be approved if member fails 7-day trial with four preferred agents. Failure is defined as allergy, intolerable side effects, or significant drug-drug interaction.</p>

THRIVITE RX tablet		
TRINATAL RX 1 tablet		
VITAFOL gummies		
WESNATAL DHA COMPLETE tablet		
WESTAB PLUS tablet		

<h2>XI. Ophthalmic</h2>

Therapeutic Drug Class: OPHTHALMIC, ALLERGY – <i>Effective 4/1/2025</i>
--

No PA Required	PA Required	Non-preferred products may be approved following trial and failure of therapy with two preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions).
ALREX ^{BNR} (loteprednol) 0.2% Azelastine 0.05% Cromolyn 4% Ketotifen 0.025% (OTC) LASTACRAFT (alcaftadine) 0.25% (OTC) Olopatadine 0.1%, 0.2% (OTC) (generic Pataday Once/Twice Daily)	ALAWAY (ketotifen) 0.025% (OTC) ALOCRIL (nedocromil) 2% ALOMIDE (lodoxamide) 0.1% Bepotastine 1.5% BEPREVE (bepotastine) 1.5% Epinastine 0.05% Loteprednol 0.2% Olopatadine 0.1%, 0.2% (RX) PATADAY ONCE DAILY (olopatadine) 0.2% (OTC)	

	PATADAY TWICE DAILY (olopatadine) 0.1% (OTC) PATADAY XS ONCE DAILY (olopatadine) 0.7% (OTC) ZADITOR (ketotifen) 0.025% (OTC) ZERViate (cetirizine) 0.24%	
Therapeutic Drug Class: OPHTHALMIC, IMMUNOMODULATORS – <i>Effective 4/1/2025</i>		
No PA Required RESTASIS ^{BNR} (cyclosporine 0.05%) vials	PA Required CEQUA (cyclosporine) 0.09% solution Cyclosporine 0.05% vials MIEBO (Perfluorohexyloctane/PF) RESTASIS MULTIDOSE (cyclosporine) 0.05% TYRVAYA (varenicline) nasal spray VERKAZIA (cyclosporin emulsion) VEVYE (cyclosporine) 0.1% XIIDRA (lifitegrast) 5% solution	Non-preferred products may be approved for members meeting all of the following criteria: <ul style="list-style-type: none">• Member is 18 years and older AND• Member has a diagnosis of chronic dry eye AND• Member has failed a 3-month trial of one preferred product. Failure is defined as a lack of efficacy, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions AND• Prescriber is an ophthalmologist, optometrist or rheumatologist <u>Maximum Dose/Quantity:</u> 60 single use containers for 30 days 5.5 mL/20 days for Restasis Multi-Dose and Vevye 3mL/30 days for Miebo Verkazia (cyclosporine ophthalmic emulsion) may be approved if the following criteria are met: <ul style="list-style-type: none">• Member is ≥ 4 years of age AND• Verkazia is being used for the treatment of vernal keratoconjunctivitis (VKC) AND• Member has trialed and failed therapy with three agents from the following pharmacologic categories: preferred dual-acting mast cell stabilizer/antihistamine from the Ophthalmics-Allergy PDL class, oral antihistamine, preferred topical ophthalmic corticosteroid from the Ophthalmics-Anti-inflammatories PDL class. Failure is defined as lack of efficacy with 2-week trial, allergy, contraindication to therapy, intolerable side effects, or significant drug-drug interaction• <u>Quantity limit:</u> 120 single-dose 0.3 mL vials/15 days
Therapeutic Drug Class: OPHTHALMIC, ANTI-INFLAMMATORIES – <i>Effective 12/2/2025</i>		
NSAIDs		
No PA Required Diclofenac 0.1% Flurbiprofen 0.03%	PA Required ACULAR (ketorolac) 0.5%, LS 0.4% ACUVAIL (ketorolac/PF) 0.45%	Durezol (difluprednate) may be approved if meeting the following criteria:

Ketorolac 0.5%, Ketorolac LS 0.4%	Bromfenac 0.07%, 0.075%, 0.09%	<ul style="list-style-type: none">Member has a diagnosis of severe intermediate uveitis, severe panuveitis, or severe uveitis with the complication of uveitic macular edema AND has trialed and failed prednisolone acetate 1% (failure is defined as lack of efficacy, allergy, contraindication to therapy, intolerable side effects, or significant drug-drug interaction) ORMembers with a diagnosis other than those listed above require trial and failure of three preferred agents (failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction).
NEVANAC (nepafenac) 0.1%	BROMSITE (bromfenac) 0.075% ILEVRO (nepafenac) 0.03% PROLENSA (bromfenac) 0.07%	
Corticosteroids		Eysuvis (loteprednol etabonate) may be approved if meeting all of the following: <ul style="list-style-type: none">Member is ≥ 18 years of age ANDEysuvis (loteprednol etabonate) is being used for short-term treatment (up to two weeks) of the signs and symptoms of dry eye disease ANDMember has failed treatment with one preferred product in the Ophthalmic Immunomodulator therapeutic class. Failure is defined as lack of efficacy with a 3-month trial, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction) ANDMember does not have any of the following conditions:Viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella ORMycobacterial infection of the eye and fungal diseases of ocular structures<u>Quantity limit</u>: one bottle/15 days Lotemax SM (loteprednol etabonate) or Inveltys (loteprednol etabonate) may be approved if meeting all of the following: <ul style="list-style-type: none">Member is ≥ 18 years of age ANDLotemax SM or Inveltys (loteprednol etabonate) is being used for the treatment of post-operative inflammation and pain following ocular surgery ANDMember has trialed and failed therapy with two preferred loteprednol formulations (failure is defined as lack of efficacy with 2-week trial, allergy, contraindication to therapy, intolerable side effects, or significant drug-drug interaction) ANDMember has trialed and failed therapy with two preferred agents that do not contain loteprednol (failure is defined as lack of efficacy with 2-week trial, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction) ANDMember does not have any of the following conditions:<ul style="list-style-type: none">Viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella ORMycobacterial infection of the eye and fungal diseases of ocular structures All other non-preferred products may be approved with trial and failure of three preferred agents (failure is defined as lack of efficacy with 2-week trial, allergy, contraindication, intolerable side effects, or significant drug-drug interaction).
No PA Required	PA Required	
FLAREX (fluorometholone) 0.1%	Dexamethasone 0.1%	
Fluorometholone 0.1% drops	Difluprednate 0.05%	
FML FORTE (fluorometholone) 0.25% drops	DUREZOL (difluprednate) 0.05%	
LOTEMAX (loteprednol) 0.5% drops, gel	EYSUVIS (loteprednol) 0.25%	
LOTEMAX (loteprednol) 0.5% ointment	FML LIQUIFILM (fluorometholone) 0.1% drop	
Loteprednol 0.5% drops, 0.5% gel	FML S.O.P (fluorometholone) 0.1% ointment	
MAXIDEX (dexamethasone) 0.1%	INVELTYS (loteprednol) 1%	
PRED MILD (prednisolone) 0.12%	LOTEMAX SM (loteprednol) 0.38% gel	
Prednisolone acetate 1%	PRED FORTE (prednisolone) 1%	
	Prednisolone sodium phosphate 1%	

Therapeutic Drug Class: OPHTHALMIC, GLAUCOMA – <i>Effective 4/1/2025</i>		
Beta-blockers		<p>Non-preferred products may be approved following trial and failure of therapy with three preferred products, including one trial with a preferred product having the same general mechanism (such as prostaglandin analogue, alpha2-adrenergic agonist, beta-blocking agent, or carbonic anhydrase inhibitor). Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions.</p> <p>Non-preferred combination products may be approved following trial and failure of therapy with one preferred combination product AND trial and failure of individual products with the same active ingredients as the combination product being requested (if available) to establish tolerance. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions.</p> <p>Preservative free products may be approved with provider documentation of adverse effect to preservative-containing product.</p>
No PA Required	PA Required	
Carteolol 1%	Betaxolol 0.5%	
Levobunolol 0.5%	BETIMOL (timolol) 0.25%, 0.5%	
Timolol (generic Timoptic) 0.25%, 0.5%	BETOPIC-S (betaxolol) 0.25%	
	ISTALOL (timolol) 0.5%	
	Timolol (generic Istalol) 0.5% drops	
	Timolol GFS 0.25%, 0.5%	
	Timolol/PF (generic Timoptic Ocudose) 0.25%, 0.5%	
	TIMOPTIC, TIMOPTIC OCUDOSE (timolol) 0.25%, 0.5%	
	TIMOPTIC-XE (timolol GFS) 0.25%, 0.5%	
Carbonic anhydrase inhibitors		
No PA Required	PA Required	
Brinzolamide 1%	AZOPT (brinzolamide) 1%	
Dorzolamide 2%		
Prostaglandin analogue		
No PA Required	PA Required	
Latanoprost 0.005%	Bimatoprost 0.03%	
LUMIGAN ^{BNR} (bimatoprost) 0.01%	IYUZEH (latanoprost/PF) 0.005%	
TRAVATAN Z ^{BNR} (travoprost) 0.004%	Tafluprost 0.0015%	

	Tafluprost PF 0.0015% Travoprost 0.004% VYZULTA (latanoprostene) 0.024% XALATAN (latanoprost) 0.005% XELPROS (latanoprost) 0.005% ZIOPTAN (tafluprost PF) 0.0015%	
Alpha-2 adrenergic agonists		
No PA Required	PA Required	
ALPHAGAN P ^{BNR} 0.1%, 0.15% (brimonidine) Brimonidine 0.2%	Apraclonidine 0.5% Brimonidine 0.1%, 0.15% IOPIDINE (apraclonidine) 0.5%, 1%	
Other ophthalmic, glaucoma and combinations		
No PA Required	PA Required	
COMBIGAN ^{BNR} 0.2%-0.5% (brimonidine/timolol) Dorzolamide/Timolol 2%-0.5% RHOPRESSA (netarsudil) 0.02% ROCKLATAN (netarsudil/latanoprost) 0.02%-0.005%	Brimonidine/Timolol 0.2%-0.5% COSOPT/COSOPT PF (dorzolamide/timolol) 2%-0.5% Dorzolamide/Timolol PF 2%-0.5% PHOSPHOLINE IODIDE (echothiophate) 0.125% Pilocarpine 1%, 1.25%, 2%, 4% SIMBRINZA (brinzolamide/brimonidine) 1%- 0.2% VUITY (pilocarpine) 1.25%	

XII. Renal/Genitourinary		
Therapeutic Drug Class: BENIGN PROSTATIC HYPERPLASIA (BPH) AGENTS – <i>Effective 10/1/2025</i>		
No PA Required Alfuzosin ER tablet Doxazosin tablet Dutasteride capsule Finasteride tablet Tamsulosin capsule Terazosin capsule	PA Required AVODART (dutasteride) softgel CARDURA (doxazosin) tablet CARDURA XL (doxazosin ER) tablet *CIALIS (tadalafil) 2.5 mg, 5 mg tablet Dutasteride/tamsulosin capsule Finasteride/tadalafil capsule FLOMAX (tamsulosin) capsule PROSCAR (finasteride) tablet RAPAFLO (silodosin) capsule Silodosin capsule *Tadalafil 2.5 mg, 5 mg tablet Tezruly (terazosin) solution	<p>*CIALIS (tadalafil) may be approved if meeting the following criteria:</p> <ul style="list-style-type: none">• Member has a documented diagnosis of BPH AND• Member has trialed and failed each of the following:• Finasteride. Failure is defined as lack of efficacy with a 3-month trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction AND• Either a nonselective alpha blocker or tamsulosin. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction AND• Documentation of BPH diagnosis will require BOTH of the following:<ul style="list-style-type: none">○ AUA Prostate Symptom Score ≥ 8 AND○ Results of a digital rectal examAND• Cialis (tadalafil) is not being prescribed for use for continuing alpha blocker therapy, as use of tadalafil in this population is not recommended due to the potential for hypotension. <p><u>Maximum Dose:</u> Doses exceeding Cialis (tadalafil) 5mg per day will not be approved.</p> <p>Prior authorization for all other non-preferred products may be approved if meeting the following criteria:</p> <ul style="list-style-type: none">• Member has tried and failed‡ three preferred agents AND• For combinations agents, member has tried and failed‡ each of the individual agents within the combination agent and one other preferred agent. <p>‡Failure is defined as lack of efficacy with 8-week trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.</p>
Therapeutic Drug Class: ANTI-HYPERURICEMICS – <i>Effective 10/1/2025</i>		
No PA Required Allopurinol 100 mg, 300 mg tablets Colchicine tablet	PA Required Allopurinol 200 mg tablets Colchicine capsule	<p>Non-preferred xanthine oxidase inhibitor products (allopurinol or febuxostat formulations) may be approved following trial and failure of preferred allopurinol. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. If member has tested positive for the HLA-B*58:01 allele, it is not recommended that they trial allopurinol. A positive result on this genetic test will count as a failure of allopurinol.</p>

Febuxostat tablet	COLCRYS (colchicine) tablet	Prior authorization for all other non-preferred agents (non-xanthine oxidase inhibitors) may be approved after trial and failure of two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. GLOPERBA (colchicine) oral solution may be approved for members who require individual doses <0.6 mg OR for members who are unable to use a solid oral dosage form. Colchicine tablet quantity limits: <ul style="list-style-type: none">Chronic hyperuricemia/gout prophylaxis: 60 tablets per 30 daysFamilial Mediterranean Fever: 120 tablets per 30 days
Probenecid tablet	GLOPERBA (colchicine) oral solution	
Probenecid/Colchicine tablet	MITIGARE (colchicine) capsule ULORIC (febuxostat) tablet	

Therapeutic Drug Class: OVERACTIVE BLADDER AGENTS – <i>Effective 10/1/2025</i>

No PA Required	PA Required	Non-preferred products may be approved for members who have failed treatment with two preferred products. Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.
Fesoterodine ER tablet	Darifenacin ER tablet	
MYRBETRIQ (mirabegron) tablet ^{BNR}	DETROL (tolterodine) tablet	
Oxybutynin IR, ER tablets, syrup	DETROL LA (tolterodine) ER capsule	
Solifenacin tablet	Flavoxate tablet	
Tolterodine tablet, ER capsule	GEMTESA (vibegron) tablet	
Trospium ER tablet	Mirabegron tablet	
	MYRBETRIQ (mirabegron) suspension	
	Oxybutynin 2.5 mg tablet	
	OXYTROL (oxybutynin patch)	
	TOVIAZ (Fesoterodine ER) tablet	
	Trospium ER capsule	
	VESICARE (solifenacin) tablet	
	VESICARE LS (solifenacin) suspension	

XIII. RESPIRATORY		
Therapeutic Drug Class: RESPIRATORY AGENTS – <i>Effective 1/1/2026</i>		
Inhaled Anticholinergics		
<p>Preferred No PA Required (Unless indicated*)</p> <p><u>Solutions</u> Ipratropium solution</p> <p><u>Short-Acting Inhalation Devices</u> ATROVENT HFA (ipratropium)</p> <p><u>Long-Acting Inhalation Devices</u></p> <p>SPIRIVA Handihaler^{BNR} (tiotropium)</p> <p>*SPIRIVA RESPIMAT (tiotropium)</p>	<p>Non-Preferred PA Required</p> <p><u>Solutions</u> YUPELRI (revefenacin) solution</p> <p><u>Short-Acting Inhalation Devices</u></p> <p><u>Long-Acting Inhalation Devices</u></p> <p>INCRUSE ELLIPTA (umeclidinium)</p> <p>Tiotropium DPI</p> <p>TUDORZA PRESSAIR (aclidinium)</p>	<p>*SPIRIVA RESPIMAT (tiotropium) 1.25 mcg may be approved for members ≥ 6 years of age with a diagnosis of asthma (qualifying diagnosis verified by AutoPA). SPIRIVA RESPIMAT is intended to be used by members whose asthma is not controlled with regular use of a combination medium-dose inhaled corticosteroid and long-acting beta agonist (LABA).</p> <p>*SPIRIVA RESPIMAT (tiotropium) 2.5 mcg may be approved for members with a diagnosis of COPD who have trialed and failed SPIRIVA HANDIHALER. Failure is defined as intolerable side effects or inability to use dry powder inhaler (DPI) formulation.</p> <p>LONHALA MAGNAIR (glycopyrrolate) may be approved for members ≥ 18 years of age with a diagnosis of COPD including chronic bronchitis and emphysema who have trialed and failed‡ treatment with two preferred anticholinergic agents.</p> <p>Non-preferred single agent anticholinergic agents may be approved for members with a diagnosis of COPD including chronic bronchitis and/or emphysema who have trialed and failed‡ treatment with two preferred agents, one of which must be SPIRIVA HANDIHALER.</p> <p>‡Failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p>
Inhaled Anticholinergic Combinations		
<p>No PA Required</p> <p><u>Solutions</u> Ipratropium/Albuterol solution</p> <p><u>Short-Acting Inhalation Devices</u> COMBIVENT RESPIMAT (albuterol/ipratropium)</p> <p><u>Long-Acting Inhalation Devices</u> ANORO ELLIPTA (umeclidinium/vilanterol) ^{BNR}</p>	<p>PA Required</p> <p><u>Solutions</u></p> <p><u>Short-Acting Inhalation Devices</u></p> <p><u>Long-Acting Inhalation Devices</u> BEVESPI AEROSPHERE (glycopyrrolate /formoterol fumarate)</p> <p>BREZTRI AEROSPHERE (budesonide/glycopyrrolate/ formoterol)</p> <p>DUAKLIR PRESSAIR (aclidinium/formoterol)</p>	<p>BREZTRI AEROSPHERE (budesonide/glycopyrrolate/formoterol) may be approved for members ≥ 18 years of age with a diagnosis of COPD who have trialed and failed‡ treatment with two preferred anticholinergic-containing agents.</p> <p>DUAKLIR PRESSAIR (aclidinium/formoterol) may be approved for members ≥ 18 years of age with a diagnosis of COPD who have trialed and failed‡ treatment with two preferred anticholinergic-containing agents.</p> <p>All other non-preferred inhaled anticholinergic combination agents may be approved for members with a diagnosis of COPD including chronic bronchitis and/or emphysema who have trialed and failed‡ treatment with two preferred inhaled anticholinergic combination agents OR three preferred inhaled anticholinergic-containing agents (single ingredient or combination).</p> <p>Members who are currently stabilized on Bevespi Aerosphere may receive approval to continue therapy with that product.</p> <p>‡Failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p>

	STIOLTO RESPIMAT (tiotropium/olodaterol) Umeclidinium/Vilanterol	
Inhaled Beta2 Agonists (short acting)		
No PA Required <u>Solutions</u> Albuterol solution, for nebulizer <u>Inhalers</u> VENTOLIN ^{BNR} HFA (albuterol)	PA Required <u>Solutions</u> Levalbuterol solution <u>Inhalers</u> AIRSUPRA (budesonide/albuterol) Albuterol HFA Levalbuterol HFA PROAIR RESPICLICK (albuterol) XOPENEX (levalbuterol) Inhaler	Non-preferred short acting beta-2 agonists may be approved for members who have failed treatment with one preferred agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. MDI formulation quantity limits: 2 inhalers / 30 days <u>Airsupra minimum age:</u> 18 years old
Inhaled Beta2 Agonists (long acting)		
Preferred <u>Solutions</u> <u>Inhalers</u> SEREVENT DISKUS (salmeterol) inhaler	Non-Preferred PA Required <u>Solutions</u> Arformoterol solution BROVANA (arformoterol) solution Formoterol solution PERFOROMIST (formoterol) solution <u>Inhalers</u> STRIVERDI RESPIMAT (olodaterol)	Non-preferred agents may be approved for members with moderate to severe COPD, AND members must have failed a trial of Serevent. Failure is defined as lack of efficacy with a 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction. For treatment of members with diagnosis of asthma needing add-on therapy, please refer to preferred agents in combination Long-Acting Beta Agonist/Inhaled Corticosteroid therapeutic class.
Inhaled Corticosteroids		
No PA Required <u>Solutions</u> Budesonide nebules <u>Inhalers</u> ARNUITY ELLIPTA ^{BNR} (fluticasone furoate)	PA Required <u>Solutions</u> PULMICORT (budesonide) respules <u>Inhalers</u> ALVESCO (ciclesonide) inhaler	Non-preferred inhaled corticosteroids may be approved in members with asthma who have failed an adequate trial of two preferred agents. An adequate trial is defined as at least 6 weeks. (Failure is defined as: lack of efficacy with a 6-week trial, allergy, contraindication to, intolerable side effects, or significant drug-drug interactions, or dexterity/coordination limitations (per provider notes) that significantly impact appropriate use of a specific dosage form.)

ASMANEX HFA (mometasone furoate) inhaler	Fluticasone Ellipta	*FLUTICASONE PROPIONATE HFA is available to members without prior authorization for: <ul style="list-style-type: none">Members with a diagnosis of eosinophilic esophagitis (EoE) ORMembers ≤ 12 years of age. <u>Maximum Dose:</u> Pulmicort (budesonide) nebulizer suspension: 2mg/day <u>Quantity Limits:</u> Pulmicort flexhaler: 2 inhalers / 30 days
ASMANEX Twisthaler (mometasone)	Fluticasone propionate diskus	
PULMICORT FLEXHALER (budesonide)	*Fluticasone propionate HFA	
QVAR REDHALER (beclomethasone)		
Inhaled Corticosteroid Combinations		
No PA Required (*Must meet eligibility criteria) ADVAIR DISKUS ^{BNR} (fluticasone/salmeterol) ADVAIR HFA ^{BNR} (fluticasone/salmeterol) AIRDUO RESPICLICK ^{BNR} (fluticasone/salmeterol) DULERA (mometasone/formoterol) SYMBICORT ^{BNR} (budesonide/formoterol) inhaler *TRELEGY ELLIPTA (fluticasone furoate/ umeclidinium/vilanterol)	PA Required BREO ELLIPTA (vilanterol/fluticasone furoate) Budesonide/formoterol (generic Symbicort) Fluticasone/salmeterol (generic Airduo/Advair Diskus) Fluticasone/salmeterol HFA (generic Advair HFA) Fluticasone/vilanterol (generic Breo Ellipta) WIXELA INHUB (fluticasone/salmeterol)	*TRELEGY ELLIPTA (fluticasone furoate/umeclidinium/vilanterol) may be approved if meeting the following: <ul style="list-style-type: none">The member has trialed and failed‡ 6 weeks of continuous therapy with a long-acting beta agonist (LABA) used in combination with a long-acting muscarinic antagonist (LAMA) ORThe member has documented eosinophils ≥ 300 cells/μL and has trialed and failed‡ 6 weeks of continuous therapy with one of the following:<ul style="list-style-type: none">A product containing a long-acting beta agonist (LABA) ORA product containing a long-acting muscarinic antagonist (LAMA). Non-preferred inhaled corticosteroid combinations may be approved for members meeting both of the following criteria: <ul style="list-style-type: none">Member has a qualifying diagnosis of asthma or severe COPD ANDMember has trailed and failed‡ two preferred agents ‡Failure is defined as lack of efficacy with a 6-week trial, allergy, intolerable side effects, significant drug-drug interactions, or dexterity/coordination limitations (per provider notes) that significantly impact appropriate use of a specific dosage form.
Phosphodiesterase Inhibitors (PDEIs)		
No PA Required Roflumilast tablet	PA Required DALIRESP (roflumilast) tablet OHTUVAYRE (ensifentrine) suspension	Requests for use of the non-preferred brand product formulation may be approved if meeting criteria outlined in the Appendix P “Generic Mandate” section.