



COLORADO
Department of Health Care
Policy & Financing



Colorado Department of Health Care Policy and Financing Preferred Drug List (PDL) Effective October 1, 2025

Prior Authorization Forms: Available online at <https://hcpf.colorado.gov/pharmacy-resources>

Prior Authorization (PA) Requests: Colorado Pharmacy Call Center Phone Number: 800-424-5725 | Fax Number: 800-424-5881

Electronic Prior Authorization (ePA): Electronic Prior Authorization Requests are supported by CoverMyMeds and may be submitted via Electronic Health Record (EHR) systems or through the CoverMyMeds provider portal.

The PDL applies to Medicaid fee-for-service members. It does not apply to members enrolled in Rocky Mountain Health HMO or Denver Health Medicaid Choice.

Initiation of pharmaceutical product subject to Prior Authorization: Please note that starting the requested drug, including a non-preferred drug, prior to a PA request being reviewed and approved, through either inpatient use, by using office “samples,” or by any other means, does not necessitate Medicaid approval of the PA request.

Health First Colorado, at section 25.5-5-501, C.R.S., requires the generic of a brand name drug be prescribed if the generic is therapeutically equivalent to the brand name drug. Exceptions to this rule are: 1) If the brand name drug is more cost effective than the generic as determined by the Department, 2) If the patient has been stabilized on a brand name drug and the prescriber believes that transition to a generic would disrupt care, and 3) If the drug is being used for treatment of mental illness, cancer, epilepsy, or human immunodeficiency virus and acquired immune deficiency syndrome.

Please see the [Brand Favored Product List](#) for a list of medications where the brand name drug is more cost effective than the generic drug.

A provider may request a step therapy exception for the treatment of a serious or complex medical condition pursuant to section 25.5-4-428, C.R.S. Serious or complex medical condition means the following medical conditions: serious mental illness, cancer, epilepsy, multiple sclerosis, or human immunodeficiency virus (HIV)/ acquired immune deficiency syndrome (AIDS), or a condition requiring medical treatment to avoid death, hospitalization, or a worsening or advancing of disease progression resulting in significant harm or disability. The step therapy exception request form is available by visiting <https://hcpf.colorado.gov/pharmacy-resources>

Brand Name Required = BNR, Prior Authorization = PA, AutoPA = authorization can be automated at the point-of-sale transaction if criteria are met
Preferred drug list applies only to prescription (RX) products, unless specified.

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-preferred products will be approved for one year unless otherwise stated.)
I. Analgesics		
Therapeutic Drug Class: NON-OPIOID ANALGESIA AGENTS - Oral – Effective 4/1/2025		
No PA Required	PA Required	JOURNAVX (suzetrigine) may be approved if the following criteria are met: • Member is ≥ 18 years of age AND • Member is being prescribed suzetrigine for up to 14 days of treatment for moderate-to- severe acute pain AND • Prescriber attests that the member's pain is unable to be managed with an NSAID, acetaminophen, or other non-opioid analgesic AND • Journavx (suzetrigine) is not being prescribed to treat chronic pain AND • The medication is not being prescribed to treat pain associated with migraine AND • Member does not have severe hepatic impairment (Child-Pugh Class C) AND • Member has been counseled to avoid food or drink containing grapefruit during treatment with Journavx (suzetrigine) AND • Member is not concurrently taking a strong CYP3A inhibitor (such as ketoconazole, itraconazole, posaconazole, ritonavir, indinavir, saquinavir, clarithromycin, fluvoxamine) AND • Member is not concurrently taking a strong or moderate CYP3A inducer (such as carbamazepine, phenytoin, rifampin, efavirenz, rifabutin, St. John's Wort) · Members using hormonal contraceptives containing progestins other than levonorgestrel and norethindrone have been counseled regarding alternative or additional contraception, if appropriate, per product labeling. <u>Duration of Approval:</u> 3 months <u>Dosing Limit:</u> One 14-day course per approval on file <u>Quantity limit:</u> 29 tablets/14 days All other non-preferred oral non-opioid analgesic agents may be approved if member meets all of the following criteria: • Member has trialed and failed duloxetine (20mg, 30mg, or 60mg) AND has trialed and failed gabapentin OR pregabalin capsule (Failure is defined as lack of efficacy with 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction) Prior authorization will be required for Lyrica (pregabalin) capsule dosages > 600mg per day (maximum of 3 capsules daily) and gabapentin dosages > 3600mg per day.
Therapeutic Drug Class: NON-OPIOID ANALGESIA AGENTS - Topical – Effective 4/1/2025		
No PA Required	PA Required	Non-preferred topical products require a trial/failure with an adequate 8-week trial of gabapentin AND pregabalin AND duloxetine AND a preferred lidocaine 5% patch.
Duloxetine 20 mg, 30 mg, 60 mg capsule	CYMBALTA (duloxetine) capsule	
Gabapentin capsule, tablet, solution	DRIZALMA (duloxetine DR) sprinkle capsules	
Pregabalin capsule	Duloxetine 40 mg capsule	
SAVELLA (milnacipran) tablet, titration pack	GRALISE (gabapentin ER) tablet	
	Gabapentin ER tablet	
	HORIZANT (gabapentin ER) tablet	
	JOURNAVX (suzetrigine) tablet	
	LYRICA (pregabalin) capsule, solution, CR tablet	
	NEURONTIN (gabapentin) capsule, solution, tablet	
	Pregabalin solution, ER tablet	
Lidocaine patch	Lidocaine patch (Puretek)	

LIDODERM (lidocaine) patch	ZTLIDO (lidocaine) topical system	Failure is defined as lack of efficacy with an 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction. Lidocaine 5% patch (<i>Puretek manufacturer only</i>) may be approved if the following criteria are met: - Member is ≥ 18 years of age AND - Member has had an adequate 8-week trial and failure of: gabapentin AND pregabalin AND duloxetine AND a preferred lidocaine 5% patch. Failure is defined as lack of efficacy with an 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction AND - Prescriber has provided a justification of clinical necessity indicating that an alternative generic lidocaine 5% patch formulation cannot be used.
Therapeutic Drug Class: NON-STEROIDAL ANTI-INFLAMMATORIES (NSAIDs) - Oral – Effective 4/1/2025		
No PA Required	PA Required	
Celecoxib capsule	ARTHROTEC (diclofenac sodium/ misoprostol) tablet	DUEXIS (ibuprofen/famotidine) or VIMOVO (naproxen/esomeprazole) may be approved if the member meets the following criteria: - Trial and failure[‡] of all preferred NSAIDs at maximally tolerated doses AND - Trial and failure[‡] of three preferred proton pump inhibitors in combination with NSAID within the last 6 months AND - Has a documented history of gastrointestinal bleeding Diclofenac potassium 25 mg immediate-release tablets may be approved if the following criteria are met: - Member is ≥ 18 years of age AND - Member does not have any of the following medical conditions: - History of recent coronary artery bypass graft (CABG) surgery - History of myocardial infarction - Severe heart failure - Advanced renal disease - History of gastrointestinal bleeding AND - Member has trial and failure[‡] of four preferred oral NSAIDs at maximally tolerated doses ELYXYB (celecoxib) oral solution may be approved if the following criteria are met: - Member is ≥ 18 years of age AND - Requested medication is being prescribed for acute treatment of migraine (with or without aura) AND - Member does <u>not</u> have any of the following medical conditions: - History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDs - History of recent coronary artery bypass graft (CABG) surgery
Diclofenac potassium 50 mg tablet	CELEBREX (celecoxib) capsule	
Diclofenac sodium EC/DR tablet	COMBOGESIC (Ibuprofen/Acetaminophen) tablet	
Ibuprofen suspension, tablet (RX)	DAYPRO (oxaprozin) caplet	
Indomethacin capsule, ER capsule	Diclofenac potassium capsule, powder pack	
Ketorolac tablet*	Diclofenac potassium 25 mg tablet	
Meloxicam tablet	Diclofenac sodium ER/SR tablet	
Nabumetone tablet	Diclofenac sodium/misoprostol tablet	
Naproxen DR/ER, tablet (RX)	Diffunisal tablet	
Naproxen suspension	DUEXIS (ibuprofen/famotidine) tablet	
Sulindac tablet	ELYXYB (celecoxib) solution	
	Etodolac capsule; IR, ER tablet	
	FELDENE (piroxicam) capsule	
	Fenoprofen capsule, tablet	

	Flurbiprofen tablet Ibuprofen/famotidine tablet Ibuprofen 300 mg tablet Ketoprofen IR, ER capsule LOFENA (diclofenac) tablet Meclofenamate capsule Mefenamic acid capsule Meloxicam submicronized capsule, suspension NALFON (fenoprofen) capsule, tablet NAPRELAN (naproxen CR) tablet Naproxen sodium CR, ER, IR tablet Naproxen/esomeprazole DR tablet Oxaprozin tablet Piroxicam capsule RELAFEN DS (nabumetone) tablet Tolmetin tablet VIMOVO (naproxen/esomeprazole) DR tablet	○ History of allergic-type reactions to sulfonamides ○ Severe heart failure ○ History of myocardial infarction ○ History of gastrointestinal bleeding ○ Advanced renal disease ○ Pregnancy past 30 weeks gestation AND ● Member is unable to take an alternative NSAID in a solid oral dosage form AND ● Member has tried and failed[‡] one preferred NSAID oral liquid AND ● Member is unable to use celecoxib capsules, opened and sprinkled into applesauce or other soft food [‡]Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions. <u>Maximum dose:</u> 120 mg/day All other non-preferred oral agents may be approved following trial and failure[‡] of four preferred agents. [‡]Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions. *Ketorolac tablets quantity limit: 5-day supply per 30 days and 20 tablets per 30 days
Therapeutic Drug Class: NON-STEROIDAL ANTI-INFLAMMATORIES (NSAIDS) - Non-Oral – Effective 4/1/2025		
No PA Required	PA Required	SPRIX (ketorolac) may be approved if meeting the following criteria: ● Member is unable to tolerate, swallow or absorb oral NSAID formulations OR ● Member has trialed and failed three preferred oral or topical NSAID agents (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions) ● Quantity limit: 5-single day nasal spray bottles per 30 days All other non-preferred topical agents may be approved for members who have trialed and failed one preferred agent. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction.
Diclofenac 1.5% topical solution Diclofenac sodium 1% gel (OTC/Rx)	Diclofenac 1.3% topical patch, 2% pump FLECTOR (diclofenac) 1.3% topical patch Ketorolac nasal spray LICART (diclofenac) 1.3% topical patch	

	PENNSAID (diclofenac solution) 2% pump, 2% solution packet	Diclofenac topical patch quantity limit: 2 patches per day Diclofenac 3% gel (generic Solaraze) prior authorization criteria can be found in the Antineoplastic agents, topical, section of the PDL.
--	--	--

Opioid Utilization Policy (long-acting and short-acting opioids):

It is highly encouraged that the healthcare team utilize the Prescription Drug Monitoring Program (PDMP) to aid in ensuring safe and efficacious therapy for members using controlled substances.

Total Morphine Milligram Equivalent Policy Effective 10/1/17:

- The maximum allowable morphine milligram equivalent (MME) is 200 MME. Prescriptions for short-acting (SA) and long-acting (LA) opioids are cumulatively included in this calculation. The prescription that exceeds the cumulative MME limit of 200 MME for a member will require prior authorization and may require a provider-to-provider telephone consultation with the pain management physician (free of charge and provided by Health First Colorado).
- Prior authorization will be granted to allow for tapering
- Prior authorization for 1 year will be granted for diagnosis of sickle cell anemia
- Prior authorization for 1 year will be granted for admission to or diagnosis of hospice or end of life care
- Prior authorization for 1 year will be granted for pain associated with cancer

MME calculation is conducted using conversion factors from the following link: <https://pharmacypmp.az.gov/resources/mme-calculator>

Only one long-acting opioid agent (including different strengths) and one short-acting opioid agent (including different strengths) will be considered for a prior authorization.

Medicaid provides guidance on the treatment of pain, including tapering, on our webpage under the heading Pain Management Resources and Opioid Use at: <https://www.colorado.gov/pacific/hcpf/pain-management-resources-and-opioid-use>

Opioid Naïve Policy Effective 8/1/17 (*Update effective 04/01/23 in Italics*):

Members who have not filled a prescription for an opioid within the past 180 days will be identified as “opioid treatment naïve” and have the following limitations placed on the initial prescription(s):

- The prescription is limited to short-acting opioid agents *or Butrans (buprenorphine)*. Use of other long-acting opioid agents will require prior authorization approval for members identified as opioid treatment naïve.
- The days’ supply of the first, second, and third prescription for an opioid will be limited to 7 days, the quantity will be limited to 8 dosage forms per day (tablets, capsules), maximum #56 tablets/capsules for a 7-day supply
- The fourth prescription for an opioid will require prior authorization, filling further opioid prescriptions may require a clinical pharmacist review or provider to provider telephone consultation with a pain management physician (free of charge and provided by Health First Colorado).
- If a member has had an opioid prescription filled within the past 180 days, then this policy would not apply to that member and other opioid policies would apply as applicable.

Dental Prescriptions Opioid Policy Effective 11/15/18 (implemented in the claims system 01/07/19):

Members who receive an opioid prescribed by a dental provider will be subject to day supply limits and quantity per day limits for short acting opioids.

- The prescription is limited to short-acting opioid agents only. Use of long-acting opioid agents and short acting fentanyl agents will require prior authorization approval for members’ prescriptions written by a dental provider.
- The days’ supply of the first, second, and third prescription for an opioid will be limited to 4 days, the quantity will be limited to 6 dosage forms per day (tablets, capsules), maximum #24 tablets/capsules for a 4-day supply
- The fourth prescription for an opioid will require prior authorization. A prior authorization for the fourth fill may be approved for up to a 7-day supply and the quantity will be limited to 8 dosage forms per day (#56 tablets/capsules) for members with any of the following diagnoses/undergoing any of the following procedures:

- Traumatic oro-facial tissue injury with major mandibular/maxillary surgical procedures
- Severe cellulitis of facial planes
- Severely impacted teeth with facial space infection necessitating surgical management
- Other potential exemptions that exceed the first 3 fill limits (day supply and quantity) may be evaluated with a provider-to-provider telephone consult with a pain management specialist (free of charge and provided by Health First Colorado)

If a member has had an opioid prescription prescribed by a non-dental provider, then this policy would not apply to that member and other opioid policies would apply as applicable. Dental prescriptions do not impact the opioid treatment naïve policy, but the prescriptions will be counted towards the Morphine Milligram Equivalent (MME) daily dose.

Opioid and Benzodiazepine Combination Effective 9/15/19:

Prior authorization will be required for members receiving long-term therapy with an opioid medication who are newly started on a benzodiazepine medication **OR** for members receiving long-term therapy with a benzodiazepine medication who are newly started on an opioid medication. Prior authorization may be approved if meeting the following:

- The member discontinued or is no longer taking either the opioid or benzodiazepine medication and will not be using these in combination **OR**
- The member will not be taking the prescribed opioid and benzodiazepine medications at the same time based on prescribed dosing interval (such as prn administration) for the regimen **AND** the prescriber attests that the member has received appropriate counseling* regarding the risks associated with combining opioid and benzodiazepine medications including increased risk for sedation, respiratory depression, overdose, and overdose-related death and counseling regarding the FDA Boxed Warning for combining these medications **OR**
- The prescriber has evaluated the regimen and attests that it is appropriate for the member to continue use of the concomitant opioid and benzodiazepine medication regimen as prescribed **AND** the prescriber attests that the member has received appropriate counseling* regarding the risks associated with combining opioid and benzodiazepine medications including increased risk for sedation, respiratory depression, overdose, and overdose-related death and counseling regarding the FDA Boxed Warning for combining these medications **OR**
- Prior authorization may be approved for members receiving palliative or hospice care **OR**
- For benzodiazepine prior authorizations, approval may be granted if the benzodiazepine is being prescribed for seizure disorder or convulsions.

**If counseling has not been provided, the prescriber attests that a reasonable effort will be made to contact the member or the member's pharmacy to ensure that counseling is provided.*

Opioid and Quetiapine Combination Effective 9/15/19:

Pharmacy claims for members receiving opioid and quetiapine medications in combination will require entry of point-of-sale DUR service codes (Reason for Service, Professional Service, Result of Service) for override of drug-drug interaction (DD) related to risk of increased sedation from concomitant use of this drug combination.

Opioid and Buprenorphine-Containing substance use disorder (SUD) Product Combination Effective 06/01/21:

Opioid claims submitted for members currently receiving buprenorphine-containing SUD medications will require entry of point-of-sale DUR service codes (Reason for Service, Professional Service, Result of Service) for override of drug-drug interaction (DD) with use of this drug combination.

Therapeutic Drug Class: OPIOIDS, Short Acting – Effective 4/1/2025

Preferred No PA Required* (If criteria and quantity limit are met)	Non-Preferred PA Required	
*Acetaminophen/codeine tablets	Acetaminophen / codeine elixir	*Preferred codeine and tramadol products do not require prior authorization for adult members (18 years of age or greater) if meeting all other opioid policy criteria. Preferred codeine or tramadol products prescribed for members < 18 years of age must meet the following criteria: ● Preferred tramadol and tramadol-containing products may be approved for members < 18 years of age if meeting the following: ○ Member is 12 years to 17 years of age AND
Hydrocodone/acetaminophen solution, tablet	ASCOMP WITH CODEINE (codeine/butalbital/aspirin/cafeine)	

Hydromorphone tablet	*Butalbital/caffeine/acetaminophen/codeine capsule	- Tramadol is NOT being prescribed for post-surgical pain following tonsil or adenoid procedure AND - Member's BMI-for-age is not > 95th percentile per CDC guidelines AND - Member does not have obstructive sleep apnea or severe lung disease OR - For members < 12 years of age with complex conditions or life-limiting illness who are receiving care under a pediatric specialist, tramadol and tramadol-containing products may be approved on a case-by-case basis
Morphine IR solution, tablet	Butalbital/caffeine/aspirin/codeine capsule	
Oxycodone solution, tablet	Butalbital compound/codeine	
Oxycodone/acetaminophen tablet	Butorphanol tartrate (nasal) spray	Preferred Codeine and codeine-containing products will receive prior authorization approval for members meeting the following criteria may be approved for members < 18 years of age if meeting the following: - Member is 12 years to 17 years of age AND - Codeine is NOT being prescribed for post-surgical pain following tonsil or adenoid procedure AND - Member's BMI-for-age is not > 95th percentile per CDC guidelines AND - Member does not have obstructive sleep apnea or severe lung disease AND - Member is not pregnant, or breastfeeding AND - Renal function is not impaired (GFR > 50 ml/min) AND - Member is not receiving strong inhibitors of CYP3A4 (such as erythromycin, clarithromycin, itraconazole, ketoconazole, posaconazole, fluconazole [$\geq 200\text{mg}$ daily], voriconazole, delavirdine, and milk thistle) AND - Member meets <u>one</u> of the following: - Member has trialed codeine or codeine-containing products in the past with no history of allergy or adverse drug reaction to codeine - Member has not trialed codeine or codeine-containing products in the past and the prescriber acknowledges reading the following statement: "Approximately 1-2% of the population metabolizes codeine in a manner that exposes them to a much higher potential for toxicity. Another notable proportion of the population may not clinically respond to codeine. We ask that you please have close follow-up with members newly starting codeine and codeine-containing products to monitor for safety and efficacy."
*Tramadol 25mg, 50mg	Carisoprodol/aspirin/codeine	
*Tramadol/acetaminophen tablet	Codeine tablet	
	Dihydrocodeine/acetaminophen/caffeine tablet	
	DILAUDID (hydromorphone) solution, tablet	
	FIORICET/CODEINE (codeine/butalbital/acetaminophen/caffeine) capsule	
	Hydrocodone/ibuprofen tablet	
	Hydromorphone solution	
	Levorphanol tablet	
	Meperidine solution, tablet	
	Morphine concentrated solution, oral syringe	
	NALOCET (oxycodone/acetaminophen) tablet	Non-preferred tramadol products may be approved following trial and failure of generic tramadol 50mg tablet AND generic tramadol/acetaminophen tablet.
	Oxycodone capsule, syringe, concentrated solution	All other non-preferred short-acting opioid products may be approved following trial and failure of three preferred products. Failure is defined as allergy [‡] , lack of efficacy, intolerable side effects, or significant drug-drug interaction.
	Oxycodone/acetaminophen solution	
	Oxycodone/acetaminophen tablet (generic PROLATE)	[‡] Allergy: hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, and angioedema
	Oxymorphone tablet	Quantity Limits: Short-acting opioids will be limited to a total of 120 tablets per 30 days (4/day) per member for members who are not included in the opioid treatment naive policy.
	Pentazocine/naloxone tablet	Exceptions will be made for members with a diagnosis of a terminal illness (hospice or palliative care) or sickle cell anemia.
	PERCOCET (oxycodone/acetaminophen) tablet	

	ROXICODONE (oxycodone) tablet ROXYBOND (oxycodone) tablet SEGLENTIS (tramadol/celecoxib) tablet Tramadol 100mg tablet Tramadol solution	- For members who are receiving more than 120 tablets currently and who do not have a qualifying exemption diagnosis, a 6-month prior authorization can be granted via the prior authorization process for providers to taper members. - Please note that if more than one agent is used, the combined total utilization may not exceed 120 units in 30 days. There may be allowed certain exceptions to this limit for acute situations (for example: post-operative surgery, fractures, shingles, car accident). <u>Maximum Doses:</u> Tramadol: 400mg/day Codeine: 360mg/day Butorphanol intranasal: 10ml per 30 days (four 2.5ml 10mg/ml package units per 30 days)
Therapeutic Drug Class: FENTANYL PREPARATIONS (buccal, transmucosal, sublingual) – Effective 4/1/2025		
	PA Required ACTIQ (fentanyl citrate) lozenge Fentanyl citrate lozenge, buccal tablet FENTORA (fentanyl citrate) buccal tablet	Fentanyl buccal, intranasal, transmucosal, and sublingual products: Prior authorization approval may be granted for members experiencing breakthrough cancer pain and those that have already received and are tolerant to opioid drugs for the cancer pain AND are currently being treated with a long-acting opioid drug. The prior authorization may be granted for up to 4 doses per day. For patients in hospice or palliative care, prior authorization will be automatically granted regardless of the number of doses prescribed.
Therapeutic Drug Class: OPIOIDS, Long Acting – Effective 4/1/2025		
Preferred No PA Required (unless indicated by * criteria) BELBUCA (buprenorphine) buccal film BUTRANS ^{BNR} (buprenorphine) transdermal patch *Fentanyl 12mcg, 25mcg, 50mcg, 75mcg, 100mcg transdermal patch Morphine ER (generic MS Contin) tablet Tramadol ER (generic Ultram ER) tablet	Non-Preferred PA Required **OXYCONTIN (oxycodone ER) tablet Buprenorphine transdermal patch CONZIP (tramadol ER) capsule Fentanyl 37mcg, 62mcg, 87mcg transdermal patch Hydrocodone ER capsule, tablet Hydromorphone ER tablet HYSINGLA (hydrocodone ER) tablet Methadone (all forms)	*Belbuca (buprenorphine) buccal film may be approved for members who have trialed and failed‡ treatment with Butrans (buprenorphine) patch at a dose of 20 mcg/hr OR with prescriber confirmation that the maximum dose of Butrans 20 mcg/hr will not provide adequate analgesia. <u>Quantity limit:</u> 60 films/30 days. Oxycontin (oxycodone ER) may be approved for members who have trialed and failed‡ treatment with TWO preferred agents. All other non-preferred products may be approved for members who have trialed and failed‡ three preferred products. ‡Failure is defined as lack of efficacy with 14-day trial, allergy (hives, maculopapular rash, erythema multiforme, pustular rash, intolerable application site skin reactions, severe hypotension, bronchospasm, and angioedema), intolerable side effects, or significant drug-drug interaction.

	Morphine ER capsule MS CONTIN (morphine ER) tablet Oxycodone ER tablet Oxymorphone ER tablet Tramadol ER capsule	<u>Methadone:</u> Members may receive 30-day approval when prescribed for neonatal abstinence syndrome without requiring trial and failure of preferred agents or opioid prescriber consultation. <u>Methadone Continuation:</u> Members who have been receiving methadone for pain indications do not have to meet non-preferred criteria. All new starts for methadone will require prior authorization under the non-preferred criteria listed above. <i>If a prescriber would like to discuss strategies for tapering off methadone or transitioning to other pain management therapies for a Health First Colorado member, consultation with the Health First Colorado pain management physician is available free of charge by contacting the pharmacy call center helpdesk and requesting an opioid prescriber consult.</i> <u>Reauthorization:</u> Reauthorization for a non-preferred agent may be approved if the following criteria are met: • Provider attests to continued benefit outweighing risk of opioid medication use AND • Member met original prior authorization criteria for this drug class at time of original authorization <u>Quantity/Dosing Limits:</u> • Oxycontin and Hydrocodone ER (generic Zohydro ER) will only be approved for twice daily dosing. • Hysingla will only be approved for once daily dosing. • Fentanyl patches will require a PA for doses of more than 15 patches/30 days (if using one strength) or 30 patches for 30 days (if using two strengths). For fentanyl patch strengths of 37mcg/hr, 62mcg/hr, and 87mcg/hr, member must trial and fail two preferred strengths of separate patches that will provide the desired dose (such as 12mcg/hr + 50mcg/hr = 62mcg/hr).
Therapeutic Drug Class: BUPRENORPHINE, Injectable – Effective 7/1/2025		
Preferred No PA Required (*Must meet eligibility criteria) Brixadi Weekly/Monthly (buprenorphine) syringe Sublocade (buprenorphine) syringe	Non-Preferred PA Required	Preferred agents may be approved if the following criteria are met: • The requested medication is being dispensed directly to the healthcare professional (medication should not be dispensed directly to the member) AND • Provider attests to member's enrollment in a complete treatment program, including counseling and psychosocial support AND • Member has a documented diagnosis of moderate to severe opioid use disorder AND • For members newly started on therapy who are not currently using a transmucosal buprenorphine-containing product, prescriber attests that

		transmucosal buprenorphine induction therapy will be initiated in accordance with product labeling. <u>Maximum dose:</u> - Brixadi (buprenorphine) injection: 128 mg/month - Sublocade (buprenorphine) injection: 600 mg/month during 1st month of induction therapy; 300 mg/month maintenance dose thereafter
II. Anti-Infectives		
Therapeutic Drug Class: ANTIBIOTICS, Inhaled – <i>Effective 1/1/2025</i>		
Preferred No PA Required (*Must meet eligibility criteria) Tobramycin inhalation solution (generic TOBI) *CAYSTON (aztreonam) inhalation solution	Non-Preferred PA Required ARIKAYCE (amikacin liposomal) inhalation vial BETHKIS (tobramycin) inhalation ampule KITABIS (tobramycin) nebulizer pak TOBI (tobramycin) inhalation solution TOBI PODHALER (tobramycin) inhalation capsule Tobramycin inhalation ampule (generic Bethkis) Tobramycin nebulizer pak (generic Kitabis)	*CAYSTON (aztreonam) inhalation solution may be approved if the following criteria are met: - Member has a history of trial and failure of preferred tobramycin solution for inhalation (failure is defined as lack of efficacy with a 4-week trial, intolerable side effects, or significant drug-drug interactions) OR provider attests that member cannot use preferred tobramycin solution for inhalation due to documented allergy or contraindication to therapy AND - The member has known colonization of <i>Pseudomonas aeruginosa</i> in the lungs AND - The member has been prescribed an inhaled beta agonist to use prior to nebulization of Cayston (aztreonam). ARIKAYCE (amikacin) may be approved if the following criteria are met: - Member has refractory mycobacterium avium complex (MAC) lung disease with limited or no alternative treatment options available AND - Member has trialed and failed 6 months of therapy with a 3-drug regimen that includes a macrolide (failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions). All other non-preferred inhaled antibiotic agents may be approved if the following criteria are met: - The member has a diagnosis of cystic fibrosis with known colonization of <i>Pseudomonas aeruginosa</i> in the lungs AND - Member has history of trial and failure of preferred tobramycin solution for inhalation (failure is defined as lack of efficacy with a 4-week trial, contraindication to therapy, allergy, intolerable side effects or significant drug-drug interactions).
Table 1: Minimum Age, Maximum Dose, and Quantity Limitations		

		<table><tr><th>Drug Name</th><th>Minimum Age</th><th>Maximum Dose</th><th>Quantity Limit (Based on day supply limitation for pack size dispensed)</th></tr><tr><td>ARIKAYCE (amikacin)</td><td>≥ 18 years</td><td>590 mg once daily</td><td>Not applicable</td></tr><tr><td>BETHKIS (tobramycin)</td><td>Age ≥ 6 years</td><td>300 mg twice daily</td><td>28-day supply per 56-day period</td></tr><tr><td>CAYSTON (aztreonam)</td><td>≥ 7 years</td><td>75 mg three times daily</td><td>28-day supply per 56-day period</td></tr><tr><td>KITABIS PAK (tobramycin)</td><td>Age ≥ 6 years</td><td>300 mg twice daily</td><td>28-day supply per 56-day period</td></tr><tr><td>TOBI[†] (tobramycin)</td><td>Age ≥ 6 years</td><td>300 mg twice daily</td><td>28-day supply per 56-day period</td></tr><tr><td>TOBI PODHALER (tobramycin)</td><td>Age ≥ 6 years</td><td>112 mg twice daily</td><td>28-day supply per 56-day period</td></tr><tr><td colspan="4">[†] Limitations apply to brand product formulation only</td></tr></table> Members currently stabilized on any inhaled antibiotic agent in this class may receive approval to continue that agent.	Drug Name	Minimum Age	Maximum Dose	Quantity Limit (Based on day supply limitation for pack size dispensed)	ARIKAYCE (amikacin)	≥ 18 years	590 mg once daily	Not applicable	BETHKIS (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period	CAYSTON (aztreonam)	≥ 7 years	75 mg three times daily	28-day supply per 56-day period	KITABIS PAK (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period	TOBI [†] (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period	TOBI PODHALER (tobramycin)	Age ≥ 6 years	112 mg twice daily	28-day supply per 56-day period	[†] Limitations apply to brand product formulation only			
Drug Name	Minimum Age	Maximum Dose	Quantity Limit (Based on day supply limitation for pack size dispensed)																															
ARIKAYCE (amikacin)	≥ 18 years	590 mg once daily	Not applicable																															
BETHKIS (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period																															
CAYSTON (aztreonam)	≥ 7 years	75 mg three times daily	28-day supply per 56-day period																															
KITABIS PAK (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period																															
TOBI [†] (tobramycin)	Age ≥ 6 years	300 mg twice daily	28-day supply per 56-day period																															
TOBI PODHALER (tobramycin)	Age ≥ 6 years	112 mg twice daily	28-day supply per 56-day period																															
[†] Limitations apply to brand product formulation only																																		
Therapeutic Drug Class: ANTI-HERPETIC AGENTS - Oral – Effective 1/1/2025																																		
No PA Required Acyclovir tablet, capsule *Acyclovir suspension (<i>members under 18 years or cannot swallow a solid dosage form</i>) Famciclovir tablet Valacyclovir tablet	PA Required Acyclovir suspension (<i>all other members</i>) VALTREX (valacyclovir) tablet	Non-preferred products may be approved for members who have failed an adequate trial with two preferred products with different active ingredients. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction. Sitavig (acyclovir) buccal tablet may be approved for diagnosis of recurrent herpes labialis (cold sores) if member meets non-preferred criteria listed above AND has failed trial with oral acyclovir suspension. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction. *Acyclovir suspension does not require prior authorization for members < 18 years of age and may be approved for members ≥ 18 years of age who cannot swallow an oral dosage form. <table><tr><th colspan="3">Maximum Dose Table</th></tr><tr><th></th><th>Adult</th><th>Pediatric</th></tr><tr><td>Acyclovir</td><td>4,000 mg/day</td><td>3,200 mg/day</td></tr><tr><td>Famciclovir</td><td>2,000 mg/day</td><td></td></tr></table>	Maximum Dose Table				Adult	Pediatric	Acyclovir	4,000 mg/day	3,200 mg/day	Famciclovir	2,000 mg/day																					
Maximum Dose Table																																		
	Adult	Pediatric																																
Acyclovir	4,000 mg/day	3,200 mg/day																																
Famciclovir	2,000 mg/day																																	

		Valacyclovir	4,000 mg/day	Age 2-11 years: 3,000 mg/day Age ≥ 12 years: 4,000 mg/day	
Therapeutic Drug Class: ANTI-HERPETIC AGENTS - Topical – <i>Effective 1/1/2025</i>					
No PA Required Acyclovir cream (<i>Teva only</i>) Acyclovir ointment DENAVIR ^{BNR} (penciclovir) cream	PA Required Acyclovir cream (<i>all other manufacturers</i>) Penciclovir cream XERESE (acyclovir/ hydrocortisone) cream ZOVIRAX (acyclovir) cream, ointment	Non-Preferred Zovirax and acyclovir ointment/cream formulations may be approved for members who have failed an adequate trial with the preferred topical acyclovir ointment/cream product (diagnosis, dose and duration) as deemed by approved compendium. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) Xerese (acyclovir/hydrocortisone) prior authorization may be approved for members that meet the following criteria: • Documented diagnosis of recurrent herpes labialis AND • Member is immunocompetent AND • Member has failed treatment of at least 10 days with acyclovir (Failure is defined as significant drug-drug interaction, lack of efficacy, contraindication to or intolerable side effects) AND • Member has failed treatment of at least one day with famciclovir 1500 mg OR valacyclovir 2 grams twice daily (Failure is defined as significant drug-drug interaction, lack of efficacy, contraindication to or intolerable side effects)			
Therapeutic Drug Class: FLUOROQUINOLONES – Oral – <i>Effective 1/1/2025</i>					
Preferred No PA Required (*if meeting eligibility criteria) *CIPRO (ciprofloxacin) oral suspension ^{BNR} Ciprofloxacin tablet Levofloxacin tablet Moxifloxacin tablet	Non-Preferred PA Required BAXDELA (delafloxacin) tablet CIPRO (ciprofloxacin) tablet Ciprofloxacin oral suspension Levofloxacin oral solution Ofloxacin tablet	*CIPRO suspension does not require prior authorization for members < 18 years of age and may be approved for members ≥ 18 years of age Non-preferred products may be approved for members who have failed an adequate trial (7 days) with at least one preferred product. (Failure is defined as: lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction). Levofloxacin solution may be approved for members with prescriber attestation that member: • is unable to take Cipro (ciprofloxacin) crushed tablet or suspension OR • is < 5 years of age and being treated for pneumonia OR • has failed† an adequate trial (7 days) of ciprofloxacin suspension †Failure is defined as lack of efficacy, allergy, intolerable side effects, significant drug-drug interaction, or contraindication to therapy.			
Therapeutic Drug Class: HEPATITIS C VIRUS TREATMENTS – <i>Effective 1/1/2025</i>					
Direct Acting Antivirals (DAAs)					
Preferred No PA Required for initial treatment (*must meet eligibility criteria)	Non-Preferred PA Required EPCLUSA 400 mg-100 mg (sofosbuvir/velpatasvir) tablet	Pharmacy claims for preferred products prescribed for initial treatment will be eligible for up to a 90-day supply fill allowing for the appropriate days’ duration for completing the initial treatment regimen (with no PA required). Subsequent fills will require prior authorization meeting re-treatment criteria below.			

EPCLUSA (sofosbuvir/velpatasvir) 200 mg -50 mg, 150 mg-37.5 mg tablet, pellet pack HARVONI (ledipasvir/sofosbuvir) 45mg-200mg tablet, pellet pack Ledipasvir/Sofosbuvir 90 mg-400 mg tablet (<i>Asegua only</i>) MAVYRET (glecaprevir/pibrentasvir) tablet, pellet pack Sofosbuvir/Velpatasvir 400mg-100mg (<i>Asegua only</i>) *VOSEVI tablet (sofosbuvir/velpatasvir/voxilaprevir)	HARVONI 90 mg-400 mg (ledipasvir/sofosbuvir) tablet SOVALDI (sofosbuvir) tablet, pellet packet ZEPATIER (elbasvir/grazoprevir) tablet	*Second line preferred agents (Vosevi) may be approved for members 18 years of age or older with chronic HCV infection who are non-cirrhotic or have compensated cirrhosis (Child-Pugh A) AND meet the following criteria: - GT 1-6 and has previously failed treatment with a regimen containing an NS5A inhibitor (such as ledipasvir, daclatasvir, or ombitasvir) OR - GT 1a or 3 and has previously failed treatment with a regimen containing sofosbuvir without an NS5A inhibitor AND - Request meets the applicable criteria below for re-treatment. Re-treatment: All requests for HCV re-treatment for members who have failed therapy with a DAA will be reviewed on a case-by-case basis. Additional information may be requested for re-treatment requests including: - Assessment of member readiness for re-treatment - Previous regimen medications and dates treated - Genotype of previous HCV infection - Any information regarding adherence to previously trialed regimen(s) and current chronic medications - Adverse effects experienced from previous treatment regimen - Concomitant therapies during previous treatment regimen - Vosevi regimens will require verification that member has been tested for evidence of active hepatitis B virus (HBV) infection and for evidence of prior HBV infection prior to initiating treatment. Non-preferred agents may be approved if documentation is provided indicating an acceptable rationale for not prescribing a preferred treatment regimen (acceptable rationale may include patient-specific medical contraindications to a preferred treatment or cases where a member has initiated treatment on a non-preferred drug and needs to complete therapy). Members currently receiving treatment with a non-preferred agent will receive approval to finish their treatment regimen, provided required documentation is sent via normal prior authorization request process.
Ribavirin Products		
No PA Required Ribavirin capsule Ribavirin tablet		Preferred products are eligible for up to a 90-day supply fill. Non-preferred ribavirin products require prior authorizations which will be evaluated on a case-by-case basis.
Therapeutic Drug Class: HUMAN IMMUNODEFICIENCY VIRUS (HIV) TREATMENTS, ORAL – <i>Effective 1/1/2025</i>		

Oral products indicated for HIV pre-exposure prophylaxis (PrEP) or post-exposure prophylaxis (PEP) are eligible for coverage with a written prescription by an enrolled pharmacist. Additional information regarding pharmacist enrollment can be found at https://hcpf.colorado.gov/pharm-serv.		
Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs)		
No PA Required EDURANT (rilpivirine) tablet Efavirenz capsule, tablet Etravirine tablet INTELENCE (etravirine) tablet Nevirapine suspension, IR tablet, ER tablet PIFELTRO (doravirine) tablet		All products are preferred and do not require prior authorization.
Nucleoside/Nucleotide Reverse Transcriptase Inhibitors (NRTIs)		
No PA Required Abacavir solution, tablet Didanosine DR capsule Emtricitabine capsule EMTRIVA (emtricitabine) capsule, solution EPIVIR (lamivudine) solution, tablet Lamivudine solution, tablet RETROVIR (zidovudine) capsule, syrup Stavudine capsule Tenofovir disoproxil fumarate (TDF) tablet VIREAD (TDF) oral powder, tablet ZIAGEN (abacavir) solution, tablet Zidovudine capsule, syrup, tablet		All products are preferred and do not require prior authorization.
Protease Inhibitors (PIs)		
No PA Required		All products are preferred and do not require prior authorization.

APTIVUS (tipranavir) capsule Atazanavir capsule Darunavir tablet Fosamprenavir tablet LEXIVA (fosamprenavir) suspension, tablet NORVIR (ritonavir) powder packet, tablet PREZISTA (darunavir) suspension, tablet REYATAZ (atazanavir) capsule, powder pack Ritonavir tablet VIRACEPT (nelfinavir) tablet		
Other Agents		
No PA Required ISENTRESS (raltegravir) chewable, powder pack, tablet ISENTRESS HD (raltegravir) tablet Maraviroc tablet RUKOBIA (fostemsavir tromethamine ER) tablet SELZENTRY (maraviroc) solution, tablet SUNLENCA (lenacapavir) tablet TIVICAY (dolutegravir) tablet TIVICAY PD (dolutegravir) tablet for suspension TYBOST (cobicistat) tablet VOCABRIA (cabotegravir) tablet		All products are preferred and do not require prior authorization.

YEZTUGO (lenacapavir)		
Combination Agents		
No PA Required Abacavir/Lamivudine tablet ATRIPLA (efavirenz/Emtricitabine/TDF) tablet BIKTARVY (bictegravir/emtricitabine/TAF) tablet CIMDUO (lamivudine/TDF) tablet COMBIVIR (lamivudine/zidovudine) tablet COMPLERA (emtricitabine/rilpivirine/TDF) tablet DELSTRIGO (doravirine/lamivudine/TDF) tablet DESCOVY (emtricitabine/TAF) tablet DOVATO (dolutegravir/lamivudine) tablet Efavirenz/Emtricitabine/TDF tablet Efavirenz/Lamivudine/TDF tablet Emtricitabine/rilpivirine/TDF tablet Emtricitabine/TDF tablet EPZICOM (abacavir/lamivudine) tablet EVOTAZ (atazanavir/cobicistat) tablet GENVOYA (elvitegravir/cobicistat/emtricitabine/TAF) tablet JULUCA (dolutegravir/rilpivirine) tablet KALETRA (lopinavir/ritonavir) solution, tablet Lamivudine/Zidovudine tablet		All products are preferred and do not require prior authorization.

Lopinavir/Ritonavir solution, tablet ODEFSEY (emtricitabine/rilpivirine/TAF) tablet PREZCOBIX (darunavir/cobicistat) tablet STRIBILD (elvitegravir/cobicistat/emtricitabine/TDF) tablet SYMFI/SYMFI LO (efavirenz/lamivudine/TDF) tablet SYMTUZA (darunavir/cobicistat/emtricitabine/TAF) tablet TRIUMEQ (abacavir/dolutegravir/ lamivudine) tablet TRIUMEQ PD (abacavir/dolutegravir) tablet for suspension TRIZIVIR (abacavir/lamivudine/zidovudine) tablet *TRUVADA (emtricitabine/TDF) tablet			
Therapeutic Drug Class: TETRACYCLINES – <i>Effective 7/1/2025</i>			
No PA Required	PA Required		
Doxycycline hyclate capsules	Demeclocycline tablet		Prior authorization for non-preferred tetracycline agents may be approved if member has trialed/failed a preferred doxycycline product AND preferred minocycline. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. Prior authorization for liquid oral tetracycline formulations may be approved if member is unable to take a solid oral dosage form. Nuzyra (omadacycline) prior authorization may be approved if member meets all of the following criteria: the above “non-preferred” prior authorization criteria and the following: - Member has trialed and failed[†] therapy with a preferred doxycycline product and preferred minocycline OR clinical rationale is provided describing why these medications cannot be trialed (including resistance and sensitivity) AND - Member has diagnosis of either Community Acquired Bacterial Pneumonia (CABP) or Acute Bacterial Skin and Skin Structure Infection (ABSSI) or
Doxycycline hyclate tablets	DORYX (doxycycline DR) tablet		
Doxycycline monohydrate 50mg, 100mg capsule	Doxycycline hyclate DR tablet		
Doxycycline monohydrate tablets	Doxycycline monohydrate 75mg, 150mg capsule		
Minocycline capsules	Doxycycline monohydrate suspension		
	Minocycline IR, ER tablet		
	MINOLIRA (minocycline ER) tablet		
	MORGIDOX (doxycycline/skin cleanser) kit		

	NUZYRA (omadacycline) tablet SOLODYN ER (minocycline ER) tablet Tetracycline capsule XIMINO (minocycline ER) capsule	clinical rationale and supporting literature describing/supporting intended use AND one of the following: ○ If member diagnosis is ABSSSI, member must have trial and failure[†] of sulfamethoxazole/trimethoprim product in addition to preferred tetracyclines OR ○ If member diagnosis is CABP, member must have trial and failure[†] of either a beta-lactam antibiotic (amoxicillin/clavulanic acid) or a macrolide (azithromycin) AND • Maximum duration of use is 14 days [†]Failure is defined as lack of efficacy with 7-day trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.
--	---	---

III. Cardiovascular

Therapeutic Drug Class: **ALPHA-BLOCKERS** – *Effective 7/1/2025*

No PA Required	PA Required	
Prazosin capsule	MINIPRESS (prazosin) capsule	Non-preferred products may be approved following trial and failure of one preferred product (failure is defined as lack of efficacy with 4-week trial, allergy or intolerable side effects).

Therapeutic Drug Class: **BETA-BLOCKERS** – *Effective 7/1/2025*

Beta-Blockers, Single Agent

No PA Required (*Must meet eligibility criteria)	PA Required	
	Betaxolol tablet	*HEMANGEOL (propranolol) oral solution may be approved for members between 5 weeks and 1 year of age with proliferating infantile hemangioma requiring systemic therapy. Maximum dose: 1.7 mg/kg twice daily
Acebutolol capsule	BYSTOLIC (nebivolol) tablet	
Atenolol tablet	COREG (carvedilol) tablet	Non-preferred products may be approved following trial and failure with two preferred products (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Bisoprolol tablet	COREG CR (carvedilol ER) capsule	
Carvedilol IR tablet	Carvedilol ER capsule	INNOPRAN XL (propranolol ER) capsule brand product formulation may be approved if meeting the following: • Request meets non-preferred criteria listed above AND • Member has trialed and failed therapy with a generic propranolol ER capsule formulation OR prescriber provides clinical rationale supporting why generic propranolol ER capsule product formulations cannot be trialed. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions.
*HEMANGEOL (propranolol) solution	INDERAL LA/XL (propranolol ER) capsule	
Labetalol tablet	INNOPRAN XL (propranolol ER) capsule	KASPARGO SPRINKLE (metoprolol succinate) extended-release capsule may be approved for members ≥ 6 years of age who are unable to take a solid oral dosage form. Maximum dose: 200mg/day (adult); 50mg/day (pediatric)
Metoprolol tartrate tablet	KASPARGO (metoprolol succinate) sprinkle capsule	
Metoprolol succinate ER tablet	LOPRESSOR (metoprolol tartrate) tablet	
Nadolol tablet	Pindolol tablet	

Nebivolol tablet	TENORMIN (atenolol) tablet	Members currently stabilized on timolol oral tablet non-preferred products may receive approval to continue on that product.																																																																						
Propranolol IR tablet, solution	Timolol tablet	Members currently stabilized on the non-preferred Bystolic (nebivolol) tablets may receive approval to continue on that product.																																																																						
Propranolol ER capsule	TOPROL XL (metoprolol succinate) tablet	Members currently stabilized on the non-preferred carvedilol ER capsules may receive approval to continue on that product.																																																																						
<table><tr><th colspan="5">Table 1: Receptor Selectivity and Other Properties of Preferred Beta Blockers</th></tr><tr><th></th><th>β₁</th><th>β₂</th><th>Alpha-1 receptor antagonist</th><th>Intrinsic sympathomimetic activity (ISA)</th></tr><tr><td>Acebutolol</td><td>X</td><td></td><td></td><td>X</td></tr><tr><td>Atenolol</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Betaxolol</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Bisoprolol</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Carvedilol</td><td>X</td><td>X</td><td>X</td><td></td></tr><tr><td>Labetalol</td><td>X</td><td>X</td><td>X</td><td></td></tr><tr><td>Metoprolol succinate</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Metoprolol tartrate</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Nadolol</td><td>X</td><td>X</td><td></td><td></td></tr><tr><td>Nebivolol</td><td>X</td><td></td><td></td><td></td></tr><tr><td>Pindolol</td><td>X</td><td>X</td><td></td><td>X</td></tr><tr><td>Propranolol</td><td>X</td><td>X</td><td></td><td></td></tr></table>			Table 1: Receptor Selectivity and Other Properties of Preferred Beta Blockers						β ₁	β ₂	Alpha-1 receptor antagonist	Intrinsic sympathomimetic activity (ISA)	Acebutolol	X			X	Atenolol	X				Betaxolol	X				Bisoprolol	X				Carvedilol	X	X	X		Labetalol	X	X	X		Metoprolol succinate	X				Metoprolol tartrate	X				Nadolol	X	X			Nebivolol	X				Pindolol	X	X		X	Propranolol	X	X		
Table 1: Receptor Selectivity and Other Properties of Preferred Beta Blockers																																																																								
	β ₁	β ₂	Alpha-1 receptor antagonist	Intrinsic sympathomimetic activity (ISA)																																																																				
Acebutolol	X			X																																																																				
Atenolol	X																																																																							
Betaxolol	X																																																																							
Bisoprolol	X																																																																							
Carvedilol	X	X	X																																																																					
Labetalol	X	X	X																																																																					
Metoprolol succinate	X																																																																							
Metoprolol tartrate	X																																																																							
Nadolol	X	X																																																																						
Nebivolol	X																																																																							
Pindolol	X	X		X																																																																				
Propranolol	X	X																																																																						
Beta-Blockers, Anti-Arrhythmics																																																																								
No PA Required	PA Required																																																																							
Sotalol tablet	BETAPACE/AF (sotalol) tablet SOTYLIZE (sotalol) solution	SOTYLIZE (sotalol) oral solution may be approved for members 3 days to < 5 years of age. For members ≥ 5 years of age, SOTYLIZE (sotalol) oral solution may be approved for members who are unable to take a solid oral dosage form OR members that have trialed and failed therapy with one preferred product. (Failure is defined as allergy or intolerable side effects.) Maximum dose: 320 mg/day																																																																						
Beta-Blockers, Combinations																																																																								
No PA Required	PA Required																																																																							

Atenolol/Chlorthalidone tablet	TENORETIC (atenolol/chlorthalidone) tablet	Non-preferred products may be approved following trial and failure with two preferred products (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Bisoprolol/HCTZ tablet	ZIAC (bisoprolol/HCTZ) tablet	
Metoprolol/HCTZ tablet		
Therapeutic Drug Class: CALCIUM CHANNEL-BLOCKERS – <i>Effective 7/1/2025</i>		
Dihydropyridines (DHPs)		
No PA Required	PA Required	Non-preferred products may be approved following trial and failure of two preferred agents. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions. Nimodipine oral capsule may be approved for adult members (≥ 18 years of age) with subarachnoid hemorrhage NYMALIZE (nimodipine) oral syringe may be approved for adult members (≥ 18 years of age) with subarachnoid hemorrhage who also have a feeding tube or have difficulty swallowing solid dosage forms. Maximum dose: 360 mg/day for 21 days (6 syringes/day or 126 syringes/21 days) KATERZIA (amlodipine) suspension may be approved if meeting the following: • The member has a feeding tube or confirmed difficulty swallowing solid oral dosage forms OR cannot obtain the required dose through crushed amlodipine tablets AND • For members < 6 years of age, the prescriber confirms that the member has already been receiving the medication following initiation in a hospital or other clinical setting
Amlodipine tablet	ADALAT CC (nifedipine ER) tablet	
Felodipine ER tablet	NORLIQVA (amlodipine) suspension	
Nifedipine ER tablet	KATERZIA (amlodipine) suspension	
Nifedipine IR capsule	Isradipine capsule	
	Levamlodipine tablet	
	Nicardipine capsule	
	Nimodipine capsule	
	Nisoldipine ER tablet	
	NORVASC (amlodipine) tablet	
	NYMALIZE (nimodipine) solution, oral syringe	
	PROCARDIA XL (nifedipine ER) tablet	
	SULAR (nisoldipine ER) tablet	
Non-Dihydropyridines (Non-DHPs)		
No PA Required	PA Required	Non-preferred products may be approved following trial and failure of three preferred agents. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions.
Diltiazem IR tablet	CARDIZEM (diltiazem) tablet	
Diltiazem CD/ER capsule	CARDIZEM CD/LA (diltiazem CD/ER) capsule, tablet	
Verapamil IR, ER tablet	Diltiazem ER/LA tablet	
Verapamil ER 120 mg, 180 mg, 240 mg capsule	TIAZAC ER (diltiazem ER) capsule	

	Verapamil ER 360 mg capsule Verapamil PM ER 100 mg, 200 mg, 300 mg capsule VERELAN/PM (verapamil ER) pellet capsule	
Therapeutic Drug Class: ANGIOTENSIN MODIFIERS – <i>Effective 7/1/2025</i>		
Angiotensin-converting enzyme inhibitors (ACE Inh)		
No PA Required	PA Required	Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products may be approved for members who have trialed and failed treatment with three preferred products (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction). Enalapril solution may be approved without trial and failure of three preferred agents for members who are unable to take a solid oral dosage form. QBRELIS (lisinopril) solution may be approved for members 6 years of age or older who are unable to take a solid oral dosage form and have trialed and failed Epaned (enalapril) solution. Failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction.
Benazepril tablet	ACCUPRIL (quinapril) tablet	
Enalapril tablet	ALTACE (ramipril) capsule	
Fosinopril tablet	Captopril tablet	
Lisinopril tablet	Enalapril solution	
Quinapril tablet	EPANED (enalapril) solution	
Ramipril tablet	LOTENSIN (benazepril) tablet	
	Moexipril tablet	
	Perindopril tablet	
	PRINIVIL (lisinopril) tablet	
	QBRELIS (lisinopril) solution	
	Trandolapril tablet	
	VASOTEC (enalapril) tablet	
	ZESTRIL (lisinopril) tablet	
ACE Inhibitor Combinations		
No PA Required	PA Required	Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products may be approved for members who have trialed and failed treatment with three preferred products (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction).
Amlodipine/Benazepril capsule	ACCURETIC (quinapril/HCTZ) tablet	
Benazepril/HCTZ tablet	Captopril/HCTZ tablet	

Enalapril/HCTZ tablet	Fosinopril/HCTZ tablet	
Lisinopril/HCTZ tablet	LOTENSIN HCT (benazepril/HCTZ) tablet	
Quinapril/HCTZ tablet	LOTREL (amlodipine/benazepril) capsule	
	VASERETIC (enalapril/HCTZ) tablet	
	ZESTORETIC (lisinopril/HCTZ) tablet	
Angiotensin II receptor blockers (ARBs)		
No PA Required	PA Required	Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products may be approved for members who have trialed and failed treatment with three preferred products (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction).
Irbesartan tablet	ARBLI (losartan) oral suspension	
Losartan tablet	ATACAND (candesartan) tablet	
Olmesartan tablet	AVAPRO (irbesartan) tablet	
Telmisartan tablet	BENICAR (olmesartan) tablet	
Valsartan tablet	Candesartan tablet	
	COZAAR (losartan) tablet	
	DIOVAN (valsartan) tablet	
	EDARBI (azilsartan) tablet	
	Eprosartan tablet	
	MICARDIS (telmisartan) tablet	
	Valsartan solution	
ARB Combinations		
Preferred No PA Required (Unless indicated*)	Non-Preferred PA Required	Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products may be approved for members who have trialed and failed treatment with three preferred products (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction).
*ENTRESTO (sacubitril/valsartan) tablet ^{BNR}	ATACAND HCT (candesartan/HCTZ) tablet	
	AVALIDE (irbesartan/HCTZ) tablet	
Irbesartan/HCTZ tablet	AZOR (olmesartan/amlodipine) tablet	
Losartan/HCTZ tablet	BENICAR HCT (olmesartan/HCTZ) tablet	
		*ENTRESTO (sacubitril/valsartan) may be approved for members if the following criteria are met:

Olmesartan/Amlodipine tablet	Candesartan/HCTZ tablet	- Member is 1 to 17 years of age and has a diagnosis of symptomatic heart failure with systemic left ventricular systolic dysfunction (LVSD) and/or has chronic heart failure with a below-normal left ventricular ejection fraction (LVEF) OR - Member is ≥ 18 years of age and has a diagnosis of chronic heart failure. - Diagnosis will be verified through automated verification (AutoPA) of the appropriate corresponding ICD-10 diagnosis codes related to the indicated use of the medication.
Olmesartan/HCTZ tablet	DIOVAN HCT (valsartan/HCTZ) tablet	
Telmisartan/HCTZ tablet	EDARBYCLOR (azilsartan/chlorthalidone) tablet	
Valsartan/Amlodipine tablet	ENTRESTO (sacubitril/valsartan) sprinkles	
Valsartan/HCTZ tablet	EXFORGE (valsartan/amlodipine) tablet	
	EXFORGE HCT (valsartan/amlodipine/HCTZ) tablet	
	HYZAAR (losartan/HCTZ) tablet	
	MICARDIS HCT (telmisartan/HCTZ) tablet	
	Olmesartan/amlodipine/HCTZ tablet	
	Sacubitril/valsartan tablet	
	Telmisartan/amlodipine tablet	
	TRIBENZOR (olmesartan/amlodipine/HCTZ) tablet	
	Valsartan/Amlodipine/HCTZ tablet	
Renin Inhibitors & Renin Inhibitor Combinations		
	PA Required	
	Aliskiren tablet	Non-preferred renin inhibitors and renin inhibitor combination products may be approved for members who have failed treatment with three preferred products from the angiotensin modifier class (failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction).
	TEKTURNA (aliskiren) tablet	
	TEKTURNA HCT (aliskiren/HCTZ) tablet	
Renin inhibitors and combinations will not be approved in patients with diabetes. Renin inhibitors are contraindicated when used in combination with an ACE inhibitor, ACE inhibitor combination, ARB, or ARB combination.		
Therapeutic Drug Class: PULMONARY ARTERIAL HYPERTENSION THERAPIES – Effective 7/1/2025		
Phosphodiesterase Inhibitors		

Preferred *Must meet eligibility criteria	Non-Preferred PA Required	*Eligibility criteria for preferred products:
*Sildenafil oral suspension, tablet *Tadalafil 20mg tablet	ADCIRCA (tadalafil) tablet ALYQ (tadalafil) tablet LIQREV (sildenafil) suspension REVATIO (sildenafil) suspension, tablet TADLIQ suspension	Preferred sildenafil and tadalafil tablet formulations may be approved for a diagnosis of pulmonary hypertension or right-sided heart failure. Sildenafil suspension may be approved for a diagnosis of pulmonary hypertension for members < 5 years of age who cannot take a solid oral dosage form. Non-preferred oral tablet products may be approved if meeting the following: • Member has a diagnosis of pulmonary hypertension AND • Member has trialed and failed treatment with preferred sildenafil tablet AND preferred tadalafil tablet. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction. Members who have been previously stabilized on a non-preferred product may receive approval to continue on the medication. Non-preferred oral liquid products may be approved if meeting the following: • Member has a diagnosis of pulmonary hypertension AND • Request meets one of the following: ○ Member has trialed and failed treatment with one preferred oral liquid formulation (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction) OR ○ Prescriber verifies that the member is unable to take a solid oral dosage form and that there is clinical necessity for use of a regimen with a less frequent dosing interval.

Endothelin Receptor Antagonists		
Preferred *Must meet eligibility criteria	Non-Preferred PA Required *Must meet eligibility criteria	*Eligibility Criteria for all agents in the class
*Ambrisentan tablet *Bosentan 62.5mg, 125mg tablet	Bosentan tablet for suspension LETAIRIS (ambrisentan) tablet OPSUMIT (macitentan) tablet OPSYNVI (macitentan/tadalafil) tablet	Approval may be granted for a diagnosis of pulmonary hypertension. Member and prescriber should be enrolled in applicable REMS program for prescribed medication. Non-preferred agents may be approved for members who have trialed and failed two preferred agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. TRACLEER (bosentan) tablet for suspension may be approved if meeting one of the following: • The member cannot swallow a solid oral dosage form OR • The request meets eligibility criteria and non-preferred criteria listed above.

	TRACLEER (bosentan) 32mg tablet for suspension TRACLEER (bosentan) 62.5mg, 125mg tablet	Members who have been previously stabilized on a non-preferred product may receive approval to continue the medication.
Prostacyclin Analogues and Receptor Agonists		
Preferred (*Must meet eligibility criteria) *FLOLAN (epoprostenol) vial *ORENITRAM (treprostinil ER) tablet, titration kit *REMODULIN (treprostinil) vial *VENTAVIS (iloprost) inhalation solution	Non-Preferred PA Required Epoprostenol vial Treprostinil vial TYVASO (treprostinil) inhaler, inhalation solution UPTRAVI (selexipag) tablet, dose pack, vial VELETRI (epoprostenol) vial YUTREPIA (treprostinil) capsule for inhalation	*Eligibility Criteria for all agents in the class Approval will be granted for a diagnosis of pulmonary hypertension. Non-preferred products may be approved for members who have failed treatment with a Preferred Product. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, contraindication to IV therapy or significant drug-drug interaction). Members who have been previously stabilized on a non-preferred product may receive approval to continue on the medication.
Guanylate Cyclase (sGC) Stimulator		
	Non-Preferred PA Required ADEMPAS (riociguat) tablet	ADEMPAS (riociguat) may be approved for members who meet the following criteria: - For members of childbearing potential: - Member is not pregnant and is able to receive monthly pregnancy tests while taking ADEMPAS and one month after stopping therapy AND - Member and their partners are utilizing one of the following contraceptive methods during treatment and for one month after stopping treatment (IUD, contraceptive implants, tubal sterilization, a hormone method with a barrier method, two barrier methods, vasectomy with a hormone method, or vasectomy with a barrier method) AND - Member has a CrCl ≥ 15 mL/min and is not on dialysis AND - Member does not have severe liver impairment (Child Pugh C) AND - Member has a diagnosis of persistent/recurrent chronic thromboembolic pulmonary hypertension (CTEPH) (WHO Group 4) after surgical treatment or has inoperable CTEPH OR - Member has a diagnosis of pulmonary hypertension and has failed treatment with a preferred product for pulmonary hypertension. (Failure is defined as a lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction).
Therapeutic Drug Class: LIPOTROPICS – <i>Effective 7/1/2025</i>		
Bile Acid Sequestrants		
No PA Required Colesevelam tablet	PA Required Colesevelam packet	Non-preferred bile acid sequestrants may be approved if the member has failed treatment with 2 preferred products in the last 12 months (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).

Colestipol tablet Cholestyramine packet, light packet, powder	COLESTID (colestipol) tablet, granules Colestipol granules QUESTRAN (cholestyramine/sugar) packet, powder QUESTRAN LIGHT (cholestyramine/ aspartame) packet, powder WELCHOL (colesevelam) packet, tablet	Non-preferred lipotropic agents with a preferred product with same strength, dosage form, and active ingredient may be approved with adequate trial and/or failure of the preferred product with the same ingredient (such as preferred ezetimibe and Zetia) and 2 additional agents. (Failure is defined as: lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Fibrates		
No PA Required Fenofibric acid DR (generic Trilipix) capsule Fenofibrate capsule, tablet (generic Lofibra/Tricor) Gemfibrozil tablet	PA Required ANTARA (fenofibrate) capsule Fenofibric acid tablet Fenofibrate capsule (generic Antara/Fenoglide/Lipofen) FENOGLIDE (fenofibrate) tablet LIPOFEN (fenofibrate) capsule LOPID (gemfibrozil) tablet TRICOR (fenofibrate nano) tablet TRILIPIX (fenofibric acid) capsule	Non-preferred fibrates may be approved if the member has failed treatment with generic gemfibrozil or generic fenofibrate and niacin ER in the last 12 months (failure is defined as lack of efficacy with 4-week trial of each drug, allergy, intolerable side effects or significant drug-drug interactions). Non-preferred lipotropic agents with a preferred product with same strength, dosage form, and active ingredient may be approved with adequate trial and/or failure of the preferred product with the same ingredient (such as preferred ezetimibe and Zetia) and 2 additional agents. (Failure is defined as: lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Other Lipotropics		
No PA Required (*Must meet eligibility criteria) Ezetimibe tablet Niacin ER tablet *Omega-3 ethyl esters capsule (generic Lovaza)	PA Required Icosapent ethyl capsule LOVAZA (omega-3 ethyl esters) capsule NEXLETOL (bempedoic acid) tablet	Non-preferred lipotropic agents with a preferred product with same strength, dosage form, and active ingredient may be approved with adequate trial and/or failure of the preferred product with the same ingredient and 2 additional agents. (Failure is defined as: lack of efficacy with 4-week trial, contraindication, allergy, intolerable side effects or significant drug-drug interactions). *Omega-3 ethyl esters (generic Lovaza) may be approved for members who have a baseline triglyceride level $\geq$ 500 mg/dL

	NEXLIZET (bempedoic acid/ezetimibe) tablet ZETIA (ezetimibe) tablet	Lovaza (brand name) may be approved if meeting the following: • Member has a baseline triglyceride level ≥ 500 mg/dl AND • Member has failed an adequate trial of omega-3 Ethyl Esters AND an adequate trial of gemfibrozil or fenofibrate (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions) Nexletol (bempedoic acid) or Nexlizet (bempedoic acid/ezetimibe) may be approved if meeting the following criteria: • Member is ≥ 18 years of age AND • Member is not pregnant AND • Member is not receiving concurrent simvastatin > 20 mg daily or pravastatin > 40 mg daily AND • Member has a diagnosis of either heterozygous familial hypercholesterolemia or established atherosclerotic cardiovascular disease (see definition below), AND <table border="1"><tr><td>Conditions Which Define Clinical Atherosclerotic Cardiovascular Disease</td></tr><tr><td> • Acute Coronary Syndrome • History of Myocardial Infarction • Stable or Unstable Angina • Coronary or other Arterial Revascularization • Stroke • Transient Ischemic Attack • Peripheral Arterial Disease of Atherosclerotic Origin </td></tr></table> • Member is concurrently adherent ($> 80\%$ of the past 180 days) on a maximally tolerated dose of a high intensity statin therapy (atorvastatin ≥ 40 mg daily OR rosuvastatin ≥ 20 mg daily [as a single-entity or as a combination product]) AND ezetimibe (as a single-entity or as a combination product) concomitantly for ≥ 8 continuous weeks), AND • If intolerant to a statin due to side effects, member must have a one month documented trial with at least two other maximally dosed statins in addition to ezetimibe. For members with a past or current incidence of rhabdomyolysis, a one-month trial and failure of a statin is not required, AND • Member has a treated LDL > 70 mg/dL for a clinical history of ASCVD OR LDL > 100 mg/dL if familial hypercholesterolemia <u>Initial Approval:</u> 1 year <u>Reauthorization:</u> Reauthorization may be approved for 1 year with provider attestation of medication safety and efficacy during the initial treatment period	Conditions Which Define Clinical Atherosclerotic Cardiovascular Disease	• Acute Coronary Syndrome • History of Myocardial Infarction • Stable or Unstable Angina • Coronary or other Arterial Revascularization • Stroke • Transient Ischemic Attack • Peripheral Arterial Disease of Atherosclerotic Origin
Conditions Which Define Clinical Atherosclerotic Cardiovascular Disease				
• Acute Coronary Syndrome • History of Myocardial Infarction • Stable or Unstable Angina • Coronary or other Arterial Revascularization • Stroke • Transient Ischemic Attack • Peripheral Arterial Disease of Atherosclerotic Origin				
Therapeutic Drug Class: STATINS – Effective 7/1/2025				
No PA Required Atorvastatin tablet Lovastatin tablet	PA Required ALTOPREV (lovastatin ER) tablet ATORVALIQ (atorvastatin) suspension	Non-preferred products may be approved following trial and failure of treatment with two preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions). For members who are unable to take a		

Pravastatin tablet	CRESTOR (rosuvastatin) tablet	solid oral dosage form, non-preferred liquid product formulations may be approved without requiring trial and failure of preferred products. Age Limitations: Altoprev (lovastatin ER) will not be approved for members < 10 years of age. Fluvastatin will not be approved for members < 10 years of age. Livalo (pitavastatin) will not be approved for members < 8 years of age.
Rosuvastatin tablet	EZALLOR (rosuvastatin) sprinkle capsule	
Simvastatin tablet	FLOLIPID (simvastatin) suspension	
	Fluvastatin capsule, ER tablet	
	LESCOL XL (fluvastatin ER) tablet	
	LIPITOR (atorvastatin) tablet	
	LIVALO (pitavastatin) tablet	
	Pitavastatin tablet	
	ZOCOR (simvastatin) tablet	
	ZYPITAMAG (pitavastatin) tablet	
Therapeutic Drug Class: STATIN COMBINATIONS – Effective 7/1/2025		
No PA Required	PA Required	Non-preferred Statin combinations may be approved following trial and failure of treatment with two preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions). <u>Age Limitations:</u> Vytorin and generic ezetimibe/simvastatin will not be approved for members < 18 years of age. Caduet and generic amlodipine/atorvastatin will not be approved for members < 10 years of age.
Simvastatin/Ezetimibe tablet	Atorvastatin/Amlodipine tablet	
	CADUET (atorvastatin/amlodipine) tablet	
	VYTORIN (simvastatin/ezetimibe) tablet	
Therapeutic Drug Class: Movement Disorders – Effective 7/1/2025		
No PA Required (*Must meet eligibility criteria)	PA Required	*Eligibility Criteria for all agents in the class
*Austedo (deutetrabenazine) tablet	Xenazine (tetrabenazine) tablet	• Member is ≥18 years of age AND • Member has been diagnosed with tardive dyskinesia or chorea associated with Huntington’s disease AND • If the member has hepatic impairment, FDA labeling for use has been evaluated AND
*Austedo (deutetrabenazine) XR tablet, titration pack		• <u>For chorea associated with Huntington’s disease:</u> ○ Member has been evaluated for untreated or inadequately treated depression and member has been counseled regarding the risks of depression and suicidality associated with agents in this therapeutic class.
*Ingrezza (valbenazine) capsule, initiation pack		

* Tetrabenazine tablet		AND • <u>For tardive dyskinesia:</u> ○ If applicable, the need for ongoing treatment with 1st and 2nd generation antipsychotics, metoclopramide, or prochlorperazine has been evaluated AND ○ A baseline Abnormal Involuntary Movement Scale (AIMS) has been performed. Xenazine (tetrabenazine) Maximum dose 50 mg/day (PA available for extensive metabolizers of CYP2D6) Ingrezza (valbenazine) Quantity limits: • 40 mg: 1.767 capsules/day • 60 mg: 1 capsule/day • 80 mg: 1 capsule/day Austedo (deutetrabenazine) Maximum dose: 48 mg/day Non-preferred Movement Disorder Agents may be approved for members ≥18 years of age after trial and failure of two preferred products. Failure is defined as lack of efficacy, contraindication, allergy, intolerable side effects or significant drug-drug interaction.
------------------------	--	---

IV. Central Nervous System

Therapeutic Drug Class: ANTICONVULSANTS -Oral – Effective 4/1/2025

Therapeutic Drug Class: ANTIEPILEPTIC DRUGS (AEDs) - Class Effective 7/1/2020		
No PA Required	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i>	Members currently stabilized (in outpatient or acute care settings) on any non-preferred medication in this class may receive prior authorization approval to continue on that medication.
Barbiturates		Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.
Phenobarbital elixir, solution, tablet Primidone tablet	MYSOLINE (primidone) tablet	<u>Non-Preferred Products Newly Started for Treating Seizure Disorder or Convulsions:</u> Non-preferred medications newly started for members with a diagnosis of seizure disorder/convulsions may be approved if the following criteria are met: • The requested medication is being prescribed by a practitioner who has sufficient education and experience to safely manage treatment AND • The request meets minimum age and maximum dose limits listed in Table 1 AND
Hydantoins		
	DILANTIN (phenytoin ER), 100 mg capsules	

DILANTIN (phenytoin) 30 mg capsules, Infatab, suspension PHENYTEK (phenytoin ER) capsule Phenytoin suspension, chewable, ER capsule		- For medications indicated for use as adjunctive therapy, the medication is being used in conjunction with another medication indicated for treatment of seizure disorder/convulsions AND - The request meets additional criteria listed for any of the following: APTOM (eslicarbazepine) - Member has history of trial and failure‡ of any carbamazepine-containing product BRIVIACT (brivaracetam) - Member has history of trial and failure‡ of any levetiracetam-containing product DIACOMIT (stiripentol) - Member is concomitantly taking clobazam AND - Member has diagnosis of seizures associated with Dravet syndrome ELEPSIA XR (levetiracetam ER) tablet - Member has history of trial and failure‡ of levetiracetam ER (KEPPRA XR) EPIDIOLEX (cannabidiol) - Member has diagnosis of seizures associated with Lennox-Gastaut syndrome (LGS) or Dravet Syndrome OR - Member has a diagnosis of seizures associated with tuberous sclerosis complex (TSC). FINTEPLA (fenfluramine) - Member has a diagnosis of seizures associated with Dravet syndrome or Lennox-Gastaut syndrome OXTELLAR XR (oxcarbazepine ER) - Member is being treated for partial-onset seizures AND - Member has history of trial and failure‡ of any carbamazepine or oxcarbazepine-containing product SPRITAM (levetiracetam) tablet for suspension - Member has history of trial and failure‡ of levetiracetam solution SYMPAZAN (clobazam) film - Member has history of trial and failure‡ of clobazam tablet or solution OR - Provider attests that member cannot take clobazam tablet or solution <u>Non-Preferred Products Newly Started for Non-Seizure Disorder Diagnoses:</u> Non-preferred medications newly started for non-seizure disorder diagnoses may be approved if meeting the following criteria: - Member has history of trial and failure‡ of two preferred agents AND
Succinamides		
Ethosuximide capsule, solution	CELONTIN (methsuximide) Kapseal Methsuximide capsule ZARONTIN (ethosuximide) capsule, solution	
Benzodiazepines		
Clobazam oral syringe, tablet, suspension Clonazepam tablet, ODT	KLONOPIN (clonazepam) tablet ONFI (clobazam) suspension, tablet SYMPAZAN (clobazam) SL film	
Valproic Acid and Derivatives		
DEPAKOTE (divalproex DR) sprinkle capsule Divalproex sprinkle capsule, DR tablet, ER tablet Valproic acid capsule, solution	DEPAKOTE (divalproex DR) tablet DEPAKOTE ER (divalproex ER) tablet	
Carbamazepine Derivatives		
Carbamazepine IR tablet, ER tablet, chewable, ER capsule, suspension CARBATROL ER (carbamazepine) capsule Oxcarbazepine tablet	APTOM (eslicarbazepine) tablet Eslicarbazepine tablet EQUETRO (carbamazepine) capsule Oxcarbazepine suspension	

TEGRETOL (carbamazepine) suspension, tablet TEGRETOL XR (carbamazepine ER) tablet TRILEPTAL^{BNR} (oxcarbazepine) suspension	Oxcarbazepine ER (generic Oxtellar XR) tablet OXTELLAR XR (oxcarbazepine) tablet TRILEPTAL (oxcarbazepine) tablet	The prescription meets minimum age and maximum dose limits listed in Table 1. ‡Failure is defined as lack of efficacy, allergy, intolerable side effects, significant drug-drug interaction, documented contraindication to therapy, or inability to take preferred formulation. Members identified as HLA-B*15:02 positive, carbamazepine and oxcarbazepine should be avoided per Clinical Pharmacogenetics Implementation Consortium Guideline. This may be considered a trial for prior authorization approvals of a non-preferred agent.
Lamotrigines		
Lamotrigine IR tablet, ER tablet, chewable/dispersible tablet, ODT	LAMICTAL (lamotrigine) chewable/dispersible dose pack, tablet LAMICTAL (lamotrigine) ODT, ODT dose pack LAMICTAL XR (lamotrigine ER) tablet, dose pack Lamotrigine ER/IR/ODT dose packs	
Topiramates		
Topiramate tablet, sprinkle capsule	EPRONTIA (topiramate) solution QUDEXY XR (topiramate) capsule TOPAMAX (topiramate) tablet, sprinkle capsule Topiramate ER capsule, solution TROKENDI XR (topiramate ER) capsule	
Brivaracetam/Levetiracetam		
Levetiracetam IR tablet, ER tablet, solution	BRIVIACT (brivaracetam) solution, tablet ELEPSIA XR (levetiracetam ER) tablet	

Table 1: Non-preferred Product Minimum Age and Maximum Dose		
	Minimum Age**	Maximum Dose**
Barbiturates		
primidone (MYSOLINE)		2,000 mg per day
Benzodiazepines		
clobazam (ONFI) suspension, tablet	2 years	40 mg per day
clobazam film (SYMPAZAN)	2 years	40 mg per day
clonazepam (KLONOPIN)		20 mg per day
Brivaracetam/Levetiracetam		
brivaracetam (BRIVIACT)	1 month	200 mg per day
levetiracetam (KEPPRA)	1 month	3,000 mg per day
levetiracetam (SPRITAM)	4 years	3,000 mg per day
levetiracetam ER (ELEPSIA XR)	12 years	3,000 mg per day
levetiracetam ER (KEPPRA XR)	12 years	3,000 mg per day
Carbamazepine Derivatives		
carbamazepine (EPITOL)		1,600 mg per day
carbamazepine ER (EQUETRO)		1,600 mg per day
eslicarbazepine (APTiom)	4 years	1,600 mg per day
oxcarbazepine ER (OXTELLAR XR)	6 years	2,400 mg per day
Hydantoins		
phenytoin ER (DILANTIN) 100mg capsules, suspension, Infatab		1,000 mg loading dose 600 mg/day maintenance dose
Lamotrigines		
lamotrigine IR (LAMICTAL)	2 years	500 mg per day
lamotrigine (LAMICTAL ODT)	2 years	500 mg per day
lamotrigine ER (LAMICTAL XR)	13 years	600 mg per day
Succinamides		
ethosuximide (ZARONTIN)	3 years	1,500 mg/day
methsuximide (CELONTIN)		Not listed

	KEPPRA (levetiracetam) tablet, solution KEPRA XR (levetiracetam ER) tablet Levetiracetam 250mg tablets for suspension SPRITAM (levetiracetam) tablet	<table> <tr> <th>Valproic Acid and Derivatives</th><td></td><td></td></tr> <tr> <td>divalproex ER (DEPAKOTE ER)</td><td>10 years</td><td>60 mg/kg/day</td></tr> <tr> <th>Topiramates</th><td></td><td></td></tr> <tr> <td>topiramate (TOPAMAX)</td><td>2 years</td><td>400 mg per day</td></tr> <tr> <td>topiramate ER (QUDEXY XR)</td><td>2 years</td><td>400 mg per day</td></tr> <tr> <td>topiramate ER (TROKENDI XR)</td><td>6 years</td><td>400 mg per day</td></tr> <tr> <th>Other</th><td></td><td></td></tr> <tr> <td>cannabidiol (EPIDIOLEX)</td><td>1 year</td><td>25 mg/kg/day</td></tr> <tr> <td>cenobamate (XCOPRI)</td><td>18 years</td><td>400 mg per day</td></tr> <tr> <td>felbamate tablet, suspension</td><td>2 years</td><td>3,600 mg per day</td></tr> <tr> <td>fenfluramine (FINTEPLA)</td><td>2 years</td><td>26 mg per day</td></tr> <tr> <td>lacosamide (VIMPAT)</td><td>1 month</td><td>400 mg per day</td></tr> <tr> <td>perampanel (FYCOMPA)</td><td>4 years</td><td>12 mg per day</td></tr> <tr> <td>rufinamide (BANZEL) tablet and suspension</td><td>1 year</td><td>3,200 mg per day</td></tr> <tr> <td>stiripentol (DIACOMIT)</td><td>6 months (weighing ≥ 7 kg)</td><td>3,000 mg per day</td></tr> <tr> <td>tiagabine</td><td>12 years</td><td>56 mg per day</td></tr> <tr> <td>tiagabine (GABITRIL)</td><td>12 years</td><td>56 mg per day</td></tr> <tr> <td>vigabatrin</td><td>1 month</td><td>3,000 mg per day</td></tr> <tr> <td>vigabatrin (SABRIL)</td><td>1 month</td><td>3,000 mg per day</td></tr> <tr> <td>vigabatrin (VIGADRONE) powder packet</td><td>1 month</td><td>3,000 mg per day</td></tr> <tr> <td>zonisamide (ZONEGRAN)</td><td>16 years</td><td>600 mg per day</td></tr> <tr> <td colspan="3">**Limits based on data from FDA package insert. Approval for age/dosing that falls outside of the indicated range may be evaluated on a case-by-case basis.</td></tr> </table>	Valproic Acid and Derivatives			divalproex ER (DEPAKOTE ER)	10 years	60 mg/kg/day	Topiramates			topiramate (TOPAMAX)	2 years	400 mg per day	topiramate ER (QUDEXY XR)	2 years	400 mg per day	topiramate ER (TROKENDI XR)	6 years	400 mg per day	Other			cannabidiol (EPIDIOLEX)	1 year	25 mg/kg/day	cenobamate (XCOPRI)	18 years	400 mg per day	felbamate tablet, suspension	2 years	3,600 mg per day	fenfluramine (FINTEPLA)	2 years	26 mg per day	lacosamide (VIMPAT)	1 month	400 mg per day	perampanel (FYCOMPA)	4 years	12 mg per day	rufinamide (BANZEL) tablet and suspension	1 year	3,200 mg per day	stiripentol (DIACOMIT)	6 months (weighing ≥ 7 kg)	3,000 mg per day	tiagabine	12 years	56 mg per day	tiagabine (GABITRIL)	12 years	56 mg per day	vigabatrin	1 month	3,000 mg per day	vigabatrin (SABRIL)	1 month	3,000 mg per day	vigabatrin (VIGADRONE) powder packet	1 month	3,000 mg per day	zonisamide (ZONEGRAN)	16 years	600 mg per day	**Limits based on data from FDA package insert. Approval for age/dosing that falls outside of the indicated range may be evaluated on a case-by-case basis.		
Valproic Acid and Derivatives																																																																				
divalproex ER (DEPAKOTE ER)	10 years	60 mg/kg/day																																																																		
Topiramates																																																																				
topiramate (TOPAMAX)	2 years	400 mg per day																																																																		
topiramate ER (QUDEXY XR)	2 years	400 mg per day																																																																		
topiramate ER (TROKENDI XR)	6 years	400 mg per day																																																																		
Other																																																																				
cannabidiol (EPIDIOLEX)	1 year	25 mg/kg/day																																																																		
cenobamate (XCOPRI)	18 years	400 mg per day																																																																		
felbamate tablet, suspension	2 years	3,600 mg per day																																																																		
fenfluramine (FINTEPLA)	2 years	26 mg per day																																																																		
lacosamide (VIMPAT)	1 month	400 mg per day																																																																		
perampanel (FYCOMPA)	4 years	12 mg per day																																																																		
rufinamide (BANZEL) tablet and suspension	1 year	3,200 mg per day																																																																		
stiripentol (DIACOMIT)	6 months (weighing ≥ 7 kg)	3,000 mg per day																																																																		
tiagabine	12 years	56 mg per day																																																																		
tiagabine (GABITRIL)	12 years	56 mg per day																																																																		
vigabatrin	1 month	3,000 mg per day																																																																		
vigabatrin (SABRIL)	1 month	3,000 mg per day																																																																		
vigabatrin (VIGADRONE) powder packet	1 month	3,000 mg per day																																																																		
zonisamide (ZONEGRAN)	16 years	600 mg per day																																																																		
**Limits based on data from FDA package insert. Approval for age/dosing that falls outside of the indicated range may be evaluated on a case-by-case basis.																																																																				
Other																																																																				
*Felbamate suspension FELBATOL (felbamate) suspension FELBATOL (felbamate) ^{BNR} tablet Lacosamide solution, tablet Rufinamide tablet Zonisamide capsule	BANZEL (rufinamide) suspension, tablet DIACOMIT (stiripentol) capsule, powder packet EPIDIOLEX (cannabidiol) solution Felbamate tablet FINTEPLA (fenfluramine) solution FYCOMPA (perampanel) suspension, tablet GABITRIL (tiagabine) tablet Lacosamide UD solution MOTPOLY XR (lacosamide) capsule Perampanel tablet Rufinamide suspension SABRIL (vigabatrin) powder packet, tablet Tiagabine tablet Vigabatrin tablet, powder packet VIGAFYDE (vigabatrin) solution VIMPAT (lacosamide) solution, kit, tablet XCOPRI (cenobamate) tablet, pack																																																																			

	ZONISADE (zonisamide) suspension ZTALMY (ganaxolone) suspension	
Therapeutic Drug Class: NEWER GENERATION ANTI-DEPRESSANTS – <i>Effective 4/1/2025</i>		
No PA Required Bupropion IR, SR, XL tablet Citalopram solution, tablet Desvenlafaxine succinate ER (generic Pristiq) tablet Duloxetine (generic Cymbalta) capsule Escitalopram tablet Fluoxetine capsule, solution, 60 mg tablet Fluvoxamine tablet Mirtazapine tablet, ODT Paroxetine IR tablet Sertraline solution, tablet Trazodone tablet Venlafaxine IR tablet Venlafaxine ER capsules Vilazodone tablet	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i> APLENZIN (bupropion ER) tablet AUVELITY ER (dextromethorphan/bupropion) tablet Bupropion XL (generic Forfivo XL) tablet CELEXA (citalopram) tablet Citalopram hydrobromide capsule CYMBALTA (duloxetine) capsule Desvenlafaxine fumarate ER tablet DRIZALMA (duloxetine) sprinkle capsule EFFEXOR XR (venlafaxine ER) capsule Escitalopram solution FETZIMA (levomilnacipran ER) capsule, titration pack Fluoxetine IR tablet, DR capsule Fluvoxamine ER capsule FORFIVO XL (bupropion ER) tablet LEXAPRO (escitalopram) tablet Nefazodone tablet Paroxetine CR/ER tablet, suspension Paroxetine mesylate capsule PAXIL (paroxetine) tablet, suspension PAXIL CR (paroxetine ER) tablet	Non-preferred products may be approved for members who have failed adequate trial with two preferred newer generation anti-depressant products. If two preferred newer generation anti-depressant products are not available for indication being treated, approval of prior authorization for non-preferred products will require adequate trial of all preferred products FDA approved for that indication (failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction). Zurzuvae (zuranolone) may be approved if meeting the following criteria: • Member is ≥ 18 years of age AND • Member has a diagnosis of postpartum depression based on Diagnostic and Statistical Manual of Mental Disorders (DSM-5) criteria for a major depressive episode AND • Member is not currently pregnant AND • Prescriber attests that the member has been counseled and has been engaged in shared decision making with regard to: ○ The importance of effective contraception during zuranolone treatment, as zuranolone may cause fetal harm AND ○ Zuranolone is present in low levels in human breast milk and there are limited data on its effects on a breastfed infant AND ○ Consideration for the favorable long-term safety data associated with use of SSRIs as first-line, recommended therapies for perinatal depressive disorders by the American College of Obstetricians and Gynecologists (ACOG) or SNRIs as reasonable ACOG-recommended alternatives AND • Prescriber attests that the member has been counseled to refrain from engaging in potentially hazardous activities requiring mental alertness, including driving, for ≥ 12 hours after each zuranolone dose AND • The member has been counseled to take the medication with 400 to 1,000 calories of food containing 25% to 50% fat AND • Prescriber verifies that concomitant medications have been assessed for potential drug interactions (CNS depressants, CYP3A4 inhibitors, CYP3A4 inducers) and any needed dosage adjustments for zuranolone have been made in accordance with package labeling AND • Baseline renal and hepatic function have been assessed and prescriber verifies that dosing is appropriate in accordance with package labeling.

	PEXEVA (paroxetine mesylate) tablet PRISTIQ (desvenlafaxine succinate ER) tablet PROZAC (fluoxetine) Pulvule RALDESY (trazodone) solution REMERON (mirtazapine) Soltab (ODT), tablet Sertraline capsule TRINTELLIX (vortioxetine) tablet Venlafaxine ER tablet Venlafaxine besylate ER tablet VIIBRYD (vilazodone) tablet, dose pack WELLBUTRIN SR, XL (bupropion) tablet ZOLOFT (sertraline) tablet, oral concentrate ZURZUVAE (zuranolone) capsule	<u>Quantity Limit:</u> - Zurzuvaе 20 mg and 25 mg: 28 capsules/14 days - Zurzuvaе 30 mg: 14 capsules/14 days <u>Maximum dose:</u> 50 mg once daily <u>Duration of Approval:</u> Approval will allow 30 days to fill for one 14-day course of treatment per postpartum period Citalopram doses higher than 40mg/day for ≤60 years of age and 20mg/day for >60 years of age will require prior authorization. Please see the FDA guidance at: https://www.fda.gov/drugs/drugsafety/ucm297391.htm for important safety information. Members currently stabilized on a non-preferred newer generation antidepressant may receive approval to continue on that agent for one year if medically necessary. Verification may be provided from the prescriber or the pharmacy.
Therapeutic Drug Class: MONOAMINE OXIDASE INHIBITORS (MAOIs) – <i>Effective 4/1/2025</i>		
	PA Required EMSAM (selegiline) patch MARPLAN (isocarboxazid) tablet NARDIL (phenelzine) tablet Phenelzine tablet Tranylcypromine tablet	Non-preferred products may be approved for members who have failed adequate trial (8 weeks) with two preferred anti-depressant products. If two preferred anti-depressant products are not available for indication being treated, approval of prior authorization for non-preferred products will require adequate trial of all preferred anti-depressant products FDA approved for that indication. (Failure is defined as: lack of efficacy after 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction) Members currently stabilized on a Non-preferred MAOI antidepressant may receive approval to continue that agent for one year if medically necessary. Verification may be provided from the prescriber or the pharmacy.
Therapeutic Drug Class: TRICYCLIC ANTI-DEPRESSANTS (TCAs) – <i>Effective 4/1/2025</i>		
No PA Required Amitriptyline tablet Clomipramine capsule Desipramine tablet Doxepin 10mg, 25mg, 50mg, 75mg, 100mg, 150mg capsule, oral concentrate	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i> Amoxapine tablet ANAFRANIL (clomipramine) capsule Imipramine pamoate capsule	Non-preferred products may be approved for members who have failed adequate trial (8 weeks) with three preferred tricyclic products. If three preferred products are not available for indication being treated, approval of prior authorization for non-preferred products will require adequate trial of all tricyclic preferred products FDA approved for that indication. (Failure is defined as: lack of efficacy after 8-week trial, allergy, intolerable side effects, or significant drug-drug interaction) Members currently stabilized on a non-preferred tricyclic antidepressant may receive approval to continue on that agent for one year if medically necessary. Verification may be provided from the prescriber or the pharmacy.

Imipramine HCl tablet	NORPRAMIN (desipramine) tablet	
Nortriptyline capsule	Nortriptyline solution	
	PAMELOR (nortriptyline) capsule	
	Protriptyline tablet	
	Trimipramine capsule	
Therapeutic Drug Class: ANTI-PARKINSON'S AGENTS – Effective 4/1/2025		
Dopa decarboxylase inhibitors, dopamine precursors and combinations		
No PA Required	PA Required	
Carbidopa/Levodopa IR, ER tablet	Carbidopa tablet	Non-preferred agents may be approved with adequate trial and failure of carbidopa-levodopa IR and ER formulations (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Carbidopa/Levodopa/Entacapone tablet	Carbidopa/Levodopa ODT	Carbidopa or levodopa single agent products may be approved for members with diagnosis of Parkinson's Disease as add-on therapy to carbidopa-levodopa.
	CREXONT ER (carbidopa/levodopa) capsule	Non-preferred medications that <u>are not</u> prescribed for Parkinson's Disease (or an indication related to Parkinson's Disease) may receive approval for other FDA-labeled indications without meeting trial and failure step therapy criteria.
	DHIVY (carbidopa/levodopa) tablet	
	DUOPA (carbidopa/levodopa) suspension	
	INBRIJA (levodopa) capsule for inhalation	Members with history of trial and failure of a non-preferred Parkinson's Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred.
	LODOSYN (carbidopa) tablet	
	RYTARY ER (carbidopa/levodopa) capsule	
	SINEMET (carbidopa/levodopa) IR tablet	Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.
	STALEVO (carbidopa/levodopa/ entacapone) tablet	
MAO-B inhibitors		
No PA Required	PA Required	
Rasagiline tablet	AZILECT (rasagiline) tablet	Non-preferred agents may be approved with adequate trial and failure of selegiline capsule or tablet (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Selegiline capsule, tablet	XADAGO (safinamide) tablet	Non-preferred medications that are not prescribed for Parkinson's Disease (or an indication related to Parkinson's Disease) may receive approval for other FDA-labeled indications without meeting trial and failure step therapy criteria.
	ZELAPAR (selegiline) ODT	Members with history of trial and failure of a non-preferred Parkinson's Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form

		and active ingredient) may be considered as having met a trial and failure of the equivalent preferred. Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.
Dopamine Agonists		
No PA Required Pramipexole IR tablet Ropinirole IR tablet	PA Required APOKYN (apomorphine) SC cartridge Apomorphine SC cartridge Bromocriptine capsule, tablet MIRAPEX (pramipexole) ER tablet NEUPRO (rotigotine) patch PARLODEL (bromocriptine) capsule, tablet Pramipexole ER tablet Ropinirole ER tablet	Non-preferred agents may be approved with adequate trial and failure of ropinirole IR AND pramipexole IR (failure is defined as lack of efficacy with 4-week trial, documented contraindication to therapy, allergy, intolerable side effects or significant drug-drug interactions). APOKYN (apomorphine subcutaneous cartridge) may be approved if meeting the following: • APOKYN (apomorphine) is being used as an adjunct to other medications for acute, intermittent treatment of hypomobility, “off” episodes ("end-of-dose wearing off" and unpredictable "on/off" episodes) in patients with advanced Parkinson’s disease AND • Due to the risk of profound hypotension and loss of consciousness, member is not concomitantly using a 5HT3 antagonist such as ondansetron, granisetron, dolasetron, palonosetron or alosetron. Maximum dose: 6mg (0.6mL) three times per day KYNMOBI (apomorphine sublingual film) may be approved if meeting the following: • KYNMOBI (apomorphine) is being used for the acute, intermittent treatment of “off” episodes in patients with Parkinson’s disease AND • Due to the risk of profound hypotension and loss of consciousness, member must not be concomitantly using a 5HT3 antagonist such as ondansetron, granisetron, dolasetron, palonosetron or alosetron. Maximum dose: 30mg five times per day Non-preferred medications that <u>are not</u> prescribed for Parkinson’s Disease (or an indication related to Parkinson’s Disease) may receive approval for other FDA-labeled indications without meeting trial and failure step therapy criteria. Members with history of trial and failure of a non-preferred Parkinson’s Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred. Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.
Other Parkinson’s agents		
No PA Required	PA Required	

Amantadine capsule, solution/syrup Benztropine tablet Trihexyphenidyl tablet, elixir	Amantadine tablet COMTAN (entacapone) tablet Entacapone tablet GOCOVRI ER (amantadine ER) capsule NOURIANZ (istradefylline) tablet ONGENTYS (opicapone) capsule OSMOLEX ER (amantadine) tablet TASMAR (tolcapone) tablet Tolcapone tablet	Non-preferred agents may be approved with adequate trial and failure of two preferred agents (failure is defined as lack of efficacy with 4-week trial, documented contraindication to therapy, allergy, intolerable side effects or significant drug-drug interactions). Non-preferred medications that <u>are not</u> prescribed for Parkinson's Disease (or an indication related to Parkinson's Disease) may receive approval for other FDA-labeled indications without meeting trial and failure step therapy criteria. Members with history of trial and failure of a non-preferred Parkinson's Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred. Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.
Therapeutic Drug Class: BENZODIAZEPINES (NON-SEDATIVE HYPNOTIC) – <i>Effective 4/1/2025</i>		
No PA Required (*may be subject to age limitations) Alprazolam IR, ER tablet* Chlordiazepoxide capsule* Clonazepam tablet, ODT Clorazepate tablet* Diazepam tablet*, solution Lorazepam tablet*, oral concentrate Oxazepam capsule*	PA Required Alprazolam ODT, oral concentrate ATIVAN (lorazepam) tablet Diazepam Intensol KLONOPIN (clonazepam) tablet LOREEV (lorazepam ER) capsule XANAX (alprazolam) tablet XANAX XR (alprazolam ER) tablet	Non-preferred products may be approved following trial and failure of three preferred agents. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions. <u>Children:</u> Prior authorization will be required for all agents when prescribed for children <18 years of age (with the exception of oral solution products) and may be approved with prescriber verification of necessity of use for member age. Diazepam Intensol may be approved following trial and failure of the preferred 5 mg/5 mL oral solution. Failure is defined as intolerable side effects, drug-drug interaction, or lack of efficacy. All benzodiazepine anxiolytics will require prior authorization for members ≥ 65 years of age when exceeding 90 days of therapy. Continuation of Therapy: - Members < 65 years of age who are currently stabilized on a non-preferred benzodiazepine medication may receive approval to continue that medication. - Members < 18 years of age who are currently stabilized on a non-preferred oral solution product may receive authorization to continue that medication. Prior authorization will be required for prescribed doses that exceed the maximum (Table 1).

		Table 1 Maximum Doses		
		Product	Maximum Daily Dose	Maximum Monthly Dose
		Alprazolam tablet	<u>Adults ≥ 18 years:</u> 10 mg/day	Total of 300 mg from all dosage forms per 30 days
		Alprazolam ER tablet		
		Alprazolam ODT		
		XANAX (alprazolam) tablet		
		XANAX XR (alprazolam ER) tablet		
		Alprazolam Intensol oral concentrate 1 mg/mL		
		Clorazepate tablet	<u>≥12 years:</u> 90 mg/day <u>Children 9-12 years:</u> up to 60 mg/day	Total of 2,700 mg (adults) and 1,800 mg (children) from all tablet strengths per 30 days
		TRANXENE (clorazepate) T-Tab		
		Chlordiazepoxide capsule	<u>Adults ≥ 18 years:</u> 300 mg/day <u>Children 6-17 years:</u> up to 40 mg/day (pre-operative apprehension and anxiety)	Total of 9,000 mg (adults) and 120 mg (children, pre-op therapy) from all tablet strengths per 30 days
		Diazepam Intensol oral concentrate 5 mg/mL	<u>Adults ≥ 18 years:</u> 40 mg/day <u>Members age 6 months to 17 years:</u> up to 10 mg/day	Total of 1200 mg (adults) and 300 mg (pediatrics) from all dosage forms per 30 days
		Diazepam solution 5 mg/5 mL		
		Diazepam tablet		
		ATIVAN (lorazepam) Intensol concentrate 2 mg/mL	<u>Adults ≥ 18 years:</u> 10 mg/day <u>Children:</u> N/A	Total of 300 mg from all dosage forms per 30 days
		ATIVAN (lorazepam) tablet		
		Lorazepam oral concentrated soln 2 mg/mL		
		Lorazepam tablet		
		Oxazepam capsule	<u>Adults ≥ 18 years:</u> 120 mg/day <u>Children 6-18 years:</u> absolute dosage not established	Total of 3600 mg from all dosage forms per 30 days
		Therapeutic Drug Class: ANXIOLYTIC, NON- BENZODIAZEPINES – <i>Effective 4/1/2025</i>		
No PA Required				

Buspirone tablet		Non-preferred products may be approved following trial and failure of buspirone. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interactions.
Therapeutic Drug Class: ATYPICAL ANTI-PSYCHOTICS - Oral and Topical – <i>Effective 4/1/2025</i>		
No PA Required (unless indicated by * in criteria; all products subject to dose and age limitations)	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i>	*Vraylar (cariprazine) or Rexulti (brexpiprazole) may be approved for members after trial and failure of one preferred agent. Failure is defined as contraindication, lack of efficacy with 6-week trial, allergy, intolerable side effects, significant drug-drug interactions, or known interacting genetic polymorphism that prevents safe preferred product dosing. Non-preferred products may be approved for members meeting all of the following: • Medication is being prescribed for an FDA-Approved indication AND • Prescription meets dose and age limitations (Table 1) AND • Request meets one of the following: ○ Member has history of trial and failure of two preferred products with FDA approval for use for the prescribed indication (failure defined as lack of efficacy with 6-week trial, allergy, intolerable side effects (including rapid weight gain), contraindication, significant drug-drug interactions, or known interacting genetic polymorphism that prevents safe preferred product dosing) OR ○ Prescriber attests that within the last year (365 days) the member has trialed and failed (been unsuccessfully treated with) a preferred antipsychotic medication that was used to treat the member’s diagnosis (failure defined as lack of efficacy with 6-week trial, allergy, intolerable side effects (including rapid weight gain), significant drug-drug interactions, or known interacting genetic polymorphism that prevents safe preferred product dosing). Treatment must be under an FDA approved indication for a mental health condition or disorder. Age Limits: All products including preferred products will require a PA for members younger than the FDA approved age for the agent (Table 1). Members younger than the FDA approved age for the agent who are currently stabilized on an atypical antipsychotic will be eligible for approval. Atypical Antipsychotic prescriptions for members under 5 years of age may require a provider-provider telephone consult with a child and adolescent psychiatrist (provided at no cost to provider or member). **Quetiapine IR when given at subtherapeutic doses may be restricted for therapy. Low-dose quetiapine (<150mg/day) is only FDA approved as part of a drug titration schedule to aid patients in getting to the target quetiapine dose. PA will be required for quetiapine < 150mg per day except for utilization (when appropriate) in members 65 years or older. PA will be approved for members 10-17 years of age with approved diagnosis (Table 1) stabilized on <150mg quetiapine IR per day.
Aripiprazole tablet		
Asenapine SL tablet	ABILIFY (aripiprazole) tablet, MyCite	
Clozapine tablet	Aripiprazole oral solution, ODT	
Lurasidone tablet	CAPLYTA (lumateperone) capsule	
Olanzapine tablet, ODT	COBENFY (xanomeline/trospium) capsule, starter pack	
Paliperidone ER tablet	Clozapine ODT	
Quetiapine IR tablet**	CLOZARIL (clozapine) tablet, ODT	
Quetiapine ER tablet	FANAPT (iloperidone tablet, titration pack)	
REXULTI (brexpiprazole) dose pack, tablet*	GEODON (ziprasidone) capsule	
Risperidone ODT, oral solution, tablet	INVEGA ER (paliperidone) tablet	
VRAYLAR (cariprazine) capsule*	LATUDA (lurasidone) tablet	
Ziprasidone capsule	LYBALVI (olanzapine/samidorphan) tablet	
	NUPLAZID (pimavanserin) capsule, tablet	
	Olanzapine/Fluoxetine capsule	
	OPIPZA (aripiprazole) film	
	RISPERDAL (risperidone) tablet, oral solution	
	SAPHRIS (asenapine) SL tablet	

	SECUADO (asenapine) patch SEROQUEL IR (quetiapine IR) tablet*** SEROQUEL XR (quetiapine ER) tablet SYMBYAX (olanzapine/fluoxetine) capsule VERSACLOZ (clozapine) suspension ZYPREXA (olanzapine) tablet ZYPREXA ZYDIS (olanzapine) ODT	Aripiprazole solution: Aripiprazole <u>tablet</u> quantity limit is 2 tablets/day for pediatric members to allow for incremental dose titration and use of the preferred tablet formulation should be considered for dose titrations when possible and clinically appropriate. If incremental dose cannot be achieved with titration of the aripiprazole tablet for members < 18 years of age OR for members unable to swallow solid tablet dosage form, aripiprazole solution may be approved. For all other cases, aripiprazole solution is subject to meeting non-preferred product approval criteria listed above. Nuplazid (pimavanserin tartrate) may be approved for the treatment of hallucinations and delusions associated with Parkinson’s Disease psychosis AND following trial and failure of therapy with quetiapine or clozapine, or clinical rationale is provided supporting why these medications cannot be trialed. Failure will be defined as contraindication, intolerable side effects, drug-drug interaction, or lack of efficacy. Abilify MyCite may be approved if meeting all of the following: • Member has history of adequate trial and failure of 5 preferred agents (one trial must include aripiprazole tablet). Failure is defined as lack of efficacy with 6-week trial on maximally tolerated dose, allergy, intolerable side effects, significant drug-drug interactions AND • Information is provided regarding adherence measures being recommended by provider and followed by member (such as medication organizer or digital medication reminders) AND • Member has history of adequate trial and failure of 3 long-acting injectable formulations of atypical antipsychotics, one of which must contain aripiprazole (failure is defined as lack of efficacy with 8-week trial, contraindication, allergy, intolerable side effects, significant drug-drug interactions) AND • Abilify MyCite is being used with a MyCite patch and member is using a compatible mobile application. AND • Medication adherence information is being shared with their provider via a web portal or dashboard. <u>Quantity Limits:</u> Quantity limits will be applied to all products (Table 1). In order to receive approval for off-label dosing, the member must have an FDA approved indication and must have tried and failed on the FDA approved dosing regimen. Members currently stabilized on a non-preferred atypical antipsychotic may receive approval to continue therapy with that agent for one year.
--	---	---

Therapeutic Drug Class: ATYPICAL ANTI-PSYCHOTICS – Injectables – <i>Effective 10/1/2025</i>																																						
No PA Required ABILIFY ASIMTUFII (aripiprazole) syringe, vial ABILIFY MAINTENA (aripiprazole) syringe, vial ARISTADA ER (aripiprazole lauroxil) syringe ARISTADA INITIO (aripiprazole lauroxil) syringe Chlorpromazine ampule, vial Fluphenazine vial Fluphenazine decanoate vial HALDOL (haloperidol decanoate) ampule Haloperidol decanoate ampule, vial Haloperidol lactate syringe, vial INVEGA HAFYERA (paliperidone palmitate) syringe INVEGA SUSTENNA (paliperidone palmitate) syringe INVEGA TRINZA (paliperidone palmitate) syringe Olanzapine vial PERSERIS ER (risperidone) syringe, syringe kit	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i> GEODON (ziprasidone) vial Risperidone microspheres ER vial RYKINDO (risperidone microspheres) vial, vial kit ZYPREXA (olanzapine) vial	Preferred products do not require prior authorization. All products are subject to meeting FDA-labeled dosing quantity limits listed in Table 1. Non-preferred products may be approved for members meeting the following: • Medication is being prescribed for an FDA-Approved indication AND • Prescription meets dose limitations (Table 1) AND • Member has history of trial and failure of one preferred product with FDA approval for use for the prescribed indication (failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, contraindication, significant drug-drug interactions, or known interacting genetic polymorphism that prevents safe preferred product dosing). <table border="1"> <thead> <tr> <th colspan="3">Table 1: FDA-Labeled Dosing Quantity Limits</th></tr> <tr> <th>Long-Acting injectable</th><th>Route</th><th>Quantity Limit</th></tr> </thead> <tbody> <tr> <td>ABILIFY ASIMTUFII (aripiprazole)</td><td>IM</td><td>1 pack/2 months (56 days)</td></tr> <tr> <td>ABILIFY MAINTENA (aripiprazole)</td><td>IM</td><td>1 pack/28 days</td></tr> <tr> <td>ARISTADA ER (aripiprazole)</td><td>IM</td><td>1,064 mg: 1 pack/2 months (56 days) All other strengths: 1 pack/28 days</td></tr> <tr> <td>ARISTADA INITIO (aripiprazole)</td><td>IM</td><td>1 pack/7 weeks (49 days)</td></tr> <tr> <td>INVEGA HAFYERA (paliperidone)</td><td>IM</td><td>1 pack/6 months (168 days)</td></tr> <tr> <td>INVEGA SUSTENNA (paliperidone)</td><td>IM</td><td>156 mg: 2 packs/5 weeks (35 days) All other strengths: 1 pack/28 days</td></tr> <tr> <td>INVEGA TRINZA (paliperidone)</td><td>IM</td><td>1 pack/3 months (84 days)</td></tr> <tr> <td>PERSERIS ER (risperidone)</td><td>Subcutaneous</td><td>1 pack/28 days</td></tr> <tr> <td>RISPERDAL CONSTA (risperidone)</td><td>IM</td><td>2 packs/28 days</td></tr> <tr> <td>UZEDY ER (risperidone)</td><td>Subcutaneous</td><td>150 mg, 200 mg and 250 mg: 1 pack/2 months All other strengths: 1 pack/28 days</td></tr> </tbody> </table>	Table 1: FDA-Labeled Dosing Quantity Limits			Long-Acting injectable	Route	Quantity Limit	ABILIFY ASIMTUFII (aripiprazole)	IM	1 pack/2 months (56 days)	ABILIFY MAINTENA (aripiprazole)	IM	1 pack/28 days	ARISTADA ER (aripiprazole)	IM	1,064 mg: 1 pack/2 months (56 days) All other strengths: 1 pack/28 days	ARISTADA INITIO (aripiprazole)	IM	1 pack/7 weeks (49 days)	INVEGA HAFYERA (paliperidone)	IM	1 pack/6 months (168 days)	INVEGA SUSTENNA (paliperidone)	IM	156 mg: 2 packs/5 weeks (35 days) All other strengths: 1 pack/28 days	INVEGA TRINZA (paliperidone)	IM	1 pack/3 months (84 days)	PERSERIS ER (risperidone)	Subcutaneous	1 pack/28 days	RISPERDAL CONSTA (risperidone)	IM	2 packs/28 days	UZEDY ER (risperidone)	Subcutaneous	150 mg, 200 mg and 250 mg: 1 pack/2 months All other strengths: 1 pack/28 days
Table 1: FDA-Labeled Dosing Quantity Limits																																						
Long-Acting injectable	Route	Quantity Limit																																				
ABILIFY ASIMTUFII (aripiprazole)	IM	1 pack/2 months (56 days)																																				
ABILIFY MAINTENA (aripiprazole)	IM	1 pack/28 days																																				
ARISTADA ER (aripiprazole)	IM	1,064 mg: 1 pack/2 months (56 days) All other strengths: 1 pack/28 days																																				
ARISTADA INITIO (aripiprazole)	IM	1 pack/7 weeks (49 days)																																				
INVEGA HAFYERA (paliperidone)	IM	1 pack/6 months (168 days)																																				
INVEGA SUSTENNA (paliperidone)	IM	156 mg: 2 packs/5 weeks (35 days) All other strengths: 1 pack/28 days																																				
INVEGA TRINZA (paliperidone)	IM	1 pack/3 months (84 days)																																				
PERSERIS ER (risperidone)	Subcutaneous	1 pack/28 days																																				
RISPERDAL CONSTA (risperidone)	IM	2 packs/28 days																																				
UZEDY ER (risperidone)	Subcutaneous	150 mg, 200 mg and 250 mg: 1 pack/2 months All other strengths: 1 pack/28 days																																				

RISPERDAL CONSTA ^{BNR} (risperidone microspheres) syringe, vial UZEDY (risperidone) syringe Ziprasidone ZYPREXA RELPREVV (olanzapine pamoate) Vial kit		ZYPREXA RELPREVV (olanzapine)	IM	405 mg: 1 pack/28 days All other strengths: 1 pack/14 days
		*Requests for dosing regimens exceeding maximum may be approved for one year with pre-attestation that the member is stabilized on the requested dose and schedule.		
		<i>Note: Effective January 14, 2022, no place of service prior authorization is required for extended-release injectable medications (LAIs) used for the treatment of mental health or substance use disorders (SUD), when administered by a healthcare professional and billed under the pharmacy benefit. In addition, LAIs may be administered in any setting (pharmacy, clinic, medical office or member home) and billed to the pharmacy or medical benefit as most appropriate and in accordance with all Health First Colorado billing policies.</i>		

Table 1 Atypical Antipsychotics – FDA Approved Indication, Age Range, Quantity and Maximum Dose					
Brand	Generic	Approved Indications	Age Range	Maximum Daily Dose by Age/Indication	Quantity and Maximum Dose Limitations
ABILIFY	aripiprazole	Schizophrenia Bipolar I Disorder Bipolar I Disorder Irritability w/autistic disorder Tourette’s disorder Adjunctive treatment of MDD	≥ 13 years ≥ 18 years 10-17 years 6-17 years 6-18 years ≥ 18 years	30 mg 30 mg 30 mg 15 mg 20 mg (weight-based) 15 mg	Maximum one tablet per day (maximum of two tablets per day allowable for members < 18 years of age to accommodate for incremental dose changes)
CAPLYTA	lumateperone	Schizophrenia Bipolar I Disorder Bipolar II Disorder	≥ 18 years	42 mg	Maximum dosage of 42mg per day
CLOZARIL	clozapine	Treatment-resistant schizophrenia Recurrent suicidal behavior in schizophrenia or schizoaffective disorder	≥ 18 years	900 mg	Maximum dosage of 900mg per day

COBENFY	xanomeline and trospium	Schizophrenia	≥ 18 years	250 mg xanomeline and 60 mg trospium	Maximum two capsules per day
FANAPT	iloperidone	Schizophrenia Bipolar I Disorder	≥ 18 years	24 mg	Maximum two tablets per day
GEODON	ziprasidone	Schizophrenia Bipolar I Disorder	≥ 18 years ≥ 18 years	200 mg 160 mg	Maximum two capsules per day
INVEGA ER	paliperidone	Schizophrenia & schizoaffective disorder	≥ 12 years and weight ≥ 51 kg ≥ 12 years and weight < 51 kg	12 mg 6 mg	Maximum two 6mg tablets per day; all other strengths 1 tablet per day
LATUDA	lurasidone	Schizophrenia Schizophrenia Bipolar I disorder Bipolar I disorder	≥ 18 years 13-17 years ≥ 18 years 10-17 years	160 mg 80 mg 120 mg 80 mg	Maximum one tablet per day (If dosing 160mg for schizophrenia, then max of two tablets per day)
LYBALVI	olanzapine and samidorphan	Schizophrenia in adults Bipolar I disorder in adults	≥ 18 years ≥ 18 years	20 mg olanzapine and 10 mg samidorphan	Maximum one tablet per day
NUPLAZID	pimavanserin	Parkinson's disease psychosis	≥ 18 years	34 mg	Maximum dosage of 34mg per day
RISPERDAL	risperidone	Schizophrenia Schizophrenia Bipolar mania Irritability w/autistic disorder	≥ 18 years 13-17 years ≥ 10 years 5-17 years	16 mg 6 mg 6 mg 3 mg	Maximum dosage of 16mg/day (4 tablet/day limitation applied in claims system to allow for dose escalation and tapering)
REXULTI	brexpiprazole	Schizophrenia Adjunctive treatment of MDD Agitation associated with Alzheimer's disease (AD)	≥ 13 years ≥ 18 years	4 mg 3 mg	Maximum of 3mg/day for MDD adjunctive therapy, and agitation due to AD, Maximum of 4mg/day for schizophrenia
SAPHRIS	asenapine	Schizophrenia Bipolar mania or mixed episodes	≥ 18 years ≥ 10 years	20 mg 20 mg	Maximum two tablets per day
SECUADO	asenapine patch	Schizophrenia	≥ 18 years	7.6 mg/ 24 hours	Maximum 1 patch per day
SEROQUEL	quetiapine	Schizophrenia Schizophrenia Bipolar I mania or mixed Bipolar I mania or mixed Bipolar I depression Bipolar I Disorder Maintenance	≥ 18 years 13-17 years ≥ 18 years 10-17 years ≥ 18 years ≥ 18 years	750 mg 800 mg 800 mg 600 mg 300 mg 800 mg	Maximum three tablets per day
SEROQUEL XR	quetiapine ER	Schizophrenia Bipolar I mania Bipolar I mania Bipolar I depression Adjunctive treatment of MDD	≥ 13 years ≥ 18 years 10-17 years ≥ 18 years ≥ 18 years	800 mg 800 mg 600 mg 300 mg 300 mg	Maximum one tablet per day (for 300mg & 400mg tablets max 2 tablets per day)

SYMBYAX	olanzapine/ fluoxetine	Acute depression in Bipolar I Disorder Treatment resistant depression (MDD)	≥ 10 years	12 mg olanzapine/ 50 mg fluoxetine	Maximum three capsules per day (18mg olanzapine/75mg fluoxetine)
VERSACLOZ	clozapine	Treatment-resistant schizophrenia Recurrent suicidal behavior in schizophrenia or schizoaffective disorder	≥ 18 years ≥ 18 years	900 mg	Maximum dosage of 900 mg per day
VRAYLAR	cariprazine	Schizophrenia Acute manic or mixed episodes with Bipolar I disorder Depressive episodes with Bipolar I disorder Adjunctive treatment of MDD	≥ 18 years ≥ 18 years ≥ 18 years ≥ 18 years	6 mg 6 mg 3 mg 3 mg	Maximum dosage of 6mg/day
ZYPREXA ZYPREXA ZYDIS	olanzapine	Schizophrenia Acute manic or mixed episodes with Bipolar I disorder	≥ 13 years	20 mg	Maximum one tablet per day

Therapeutic Drug Class: CALCITONIN GENE – RELATED PEPTIDE INHIBITORS (CGRPs) – Effective 4/1/2025

PA Required for all agents		*Preferred agents may be approved if meeting the following criteria:
Preferred	Non-Preferred	
* AIMOVIG (erenumab-aooe) auto-injector * AJOVY (fremanezumab-vfrm) auto-injector, syringe * EMGALITY (galcanezumab- gnlm) pen, 120 mg syringe * NURTEC (rimegepant) ODT * UBRELVY (ubrogepant) tablet	EMGALITY (galcanezumab-gnlm) 100 mg syringe QULIPTA (atogepant) tablet ZAVZPRET (zavegepant) nasal	<u>Preferred Medications for Migraine Prevention (must meet all of the following):</u> - The requested medication is being used as preventive therapy for episodic or chronic migraine AND - Member has diagnosis of migraine with or without aura AND - Member has tried and failed 2 oral preventive pharmacological agents listed as Level A per the most current American Headache Society/American Academy of Neurology guidelines (such as divalproex, topiramate, metoprolol, propranolol). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction OR - If the prescribed medication is Nurtec, the member has tried and failed two preferred injectable product formulations. Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, significant drug-drug interaction, severe needle phobia, or member (or parent/caregiver) is unable to administer preferred CGRP inhibitor injectable formulation due to limited functional ability (such as vision impairment, limited manual dexterity and/or limited hand strength). <u>Preferred Medications for Acute Migraine Treatment (must meet all of the following):</u> - The requested medication is being used as acute treatment for migraine headache AND - Member has history of trial and failure of two triptans (failure is defined as lack of efficacy with 4-week trial, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction, severe needle phobia, or member (or parent/caregiver) is unable to administer preferred triptan injectable formulation due to limited functional ability (such as vision impairment, limited manual dexterity and/or limited hand strength).

Non-Preferred Medications for Migraine Prevention (must meet all of the following):

- The requested medication is being used as preventive therapy for episodic or chronic migraine AND
- Member has diagnosis of migraine with or without aura AND
- Member has tried and failed two oral preventive pharmacological agents listed as Level A per the most current American Headache Society/American Academy of Neurology guidelines (such as divalproex, topiramate, metoprolol, propranolol). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND
- The requested medication is not being used in combination with another CGRP medication AND
- The member has history of adequate trial and failure of three preferred products indicated for preventive therapy (failure is defined as lack of efficacy with 4-week trial, contraindication to therapy, allergy, intolerable side effects, significant drug-drug interaction, severe needle phobia, or member (or parent/caregiver) is unable to administer preferred triptan injectable formulation due to limited functional ability (such as vision impairment, limited manual dexterity and/or limited hand strength).

Non-Preferred Medications for Acute Migraine Treatment (must meet all of the following):

- Member is 18 years of age or older AND
- Medication is being prescribed to treat migraine headache with moderate to severe pain AND
- The requested medication is not being used in combination with another CGRP medication AND
- Member has history of trial and failure with all of the following (failure is defined as lack of efficacy with 4-week trial, allergy, contraindication, intolerable side effects, or significant drug-drug interaction):
 - Two triptans AND
 - One NSAID agent AND
 - One preferred agent indicated for acute migraine treatment

Non-Preferred Medications for Treatment of Episodic Cluster Headache (must meet all of the following):

- Member is 19-65 years of age AND
- Member meets diagnostic criteria for episodic cluster headache (has had no more than 8 attacks per day, a minimum of one attack every other day, and at least 4 attacks during the week prior to this medication being prescribed) AND
- Member is not taking other preventive medications to reduce the frequency of cluster headache attacks AND
- Member has history of trial and failure of all of the following (failure is defined as lack of efficacy with 4-week trial, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction):
 - Oxygen therapy AND

		○ Sumatriptan subcutaneous or intranasal OR zolmitriptan intranasal • Initial authorization will be limited to 8 weeks. Continuation (12-month authorization) will require documentation of clinically relevant improvement with no less than 30% reduction in headache frequency in a 4-week period. <u>Age Limitations:</u> All products: ≥ 18 years <table><tr><th colspan="2">Table 1. Calcitonin Gene-Related Peptide Inhibitor Quantity Limits</th></tr><tr><th>Drug Name</th><th>Maximum Dosing</th></tr><tr><td>Aimovig (erenumab)</td><td>one 140 mg autoinjector per 30 days</td></tr><tr><td>Ajovy (fremanezumab)</td><td>one 225 mg autoinjector or syringe per 30 days or three 225 mg autoinjectors or syringes every 90 days</td></tr><tr><td>Emgality 100mg (galcanezumab)</td><td>three 100 mg prefilled syringes per 30 days</td></tr><tr><td>Emgality 120 mg (galcanezumab)</td><td>two 120 mg pens or prefilled syringes once as first loading dose then one 120 mg pen or prefilled syringe per 30 days</td></tr><tr><td>Nurtec (rimegepant)</td><td>Prevention: 16 tablets/30 days; Acute Treatment: 8 tablets/30 days</td></tr><tr><td>Qulipta (atogepant)</td><td>30 tablets/30 days</td></tr><tr><td>Ubrelvy 50 mg (ubrogepant)</td><td>16 tablets/30 days</td></tr><tr><td>Ubrelvy 100 mg (ubrogepant)</td><td>16 tablets/30 days</td></tr><tr><td>ZAVZPRET (zavegepant)</td><td>6 unit-dose nasal spray devices per 30 days</td></tr></table> Members with current prior authorization approval on file for a preferred agent may receive approval for continuation of therapy with the preferred agent.	Table 1. Calcitonin Gene-Related Peptide Inhibitor Quantity Limits		Drug Name	Maximum Dosing	Aimovig (erenumab)	one 140 mg autoinjector per 30 days	Ajovy (fremanezumab)	one 225 mg autoinjector or syringe per 30 days or three 225 mg autoinjectors or syringes every 90 days	Emgality 100mg (galcanezumab)	three 100 mg prefilled syringes per 30 days	Emgality 120 mg (galcanezumab)	two 120 mg pens or prefilled syringes once as first loading dose then one 120 mg pen or prefilled syringe per 30 days	Nurtec (rimegepant)	Prevention: 16 tablets/30 days; Acute Treatment: 8 tablets/30 days	Qulipta (atogepant)	30 tablets/30 days	Ubrelvy 50 mg (ubrogepant)	16 tablets/30 days	Ubrelvy 100 mg (ubrogepant)	16 tablets/30 days	ZAVZPRET (zavegepant)	6 unit-dose nasal spray devices per 30 days
Table 1. Calcitonin Gene-Related Peptide Inhibitor Quantity Limits																								
Drug Name	Maximum Dosing																							
Aimovig (erenumab)	one 140 mg autoinjector per 30 days																							
Ajovy (fremanezumab)	one 225 mg autoinjector or syringe per 30 days or three 225 mg autoinjectors or syringes every 90 days																							
Emgality 100mg (galcanezumab)	three 100 mg prefilled syringes per 30 days																							
Emgality 120 mg (galcanezumab)	two 120 mg pens or prefilled syringes once as first loading dose then one 120 mg pen or prefilled syringe per 30 days																							
Nurtec (rimegepant)	Prevention: 16 tablets/30 days; Acute Treatment: 8 tablets/30 days																							
Qulipta (atogepant)	30 tablets/30 days																							
Ubrelvy 50 mg (ubrogepant)	16 tablets/30 days																							
Ubrelvy 100 mg (ubrogepant)	16 tablets/30 days																							
ZAVZPRET (zavegepant)	6 unit-dose nasal spray devices per 30 days																							
Therapeutic Drug Class: LITHIUM AGENTS – <i>Effective 4/1/2025</i>																								
No PA Required Lithium carbonate capsule, tablet Lithium citrate solution Lithium ER tablet	PA Required <i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i> LITHOBID ER (lithium ER) tablet	Non-preferred products may be approved with trial and failure of one preferred agent (failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, significant drug-drug interactions, intolerance to dosage form). Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.																						
Therapeutic Drug Class: NEUROCOGNITIVE DISORDER AGENTS – <i>Effective 4/1/2025</i>																								
Preferred *Must meet eligibility criteria	Non-Preferred PA Required	*Eligibility criteria for Preferred Agents – Preferred products may be approved for a diagnosis of neurocognitive disorder (eligible for AutoPA automated approval).																						

*Donepezil 5mg, 10mg tablet	ADLARITY (donepezil) patch	Non-preferred products may be approved if the member has failed treatment with one of the preferred products in the last 12 months. (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions) Members currently stabilized on a non-preferred product may receive approval to continue on that agent for one year if medically necessary and if there is a diagnosis of neurocognitive disorder.
*Donepezil ODT	ARICEPT (donepezil) tablet	
*Galantamine IR tablet	Donepezil 23mg tablet	
*Memantine IR tablet, dose pack	EXELON (rivastigmine) patch	
*Memantine ER capsule	Galantamine solution, ER capsule	
*Rivastigmine capsule, patch	Memantine IR solution	
	MESTINON (pyridostigmine) IR/ER tablet, syrup	
	Nemantine/donepezil ER capsule,	
	NAMZARIC (memantine/donepezil ER) capsule, dose pack	
	Pyridostigmine syrup, IR/ER tablet	

Therapeutic Drug Class: **SEDATIVE HYPNOTICS** – *Effective 4/1/2025*

Non-Benzodiazepines

Preferred No PA Required* (Unless age, dose, or duplication criteria apply)	Non-Preferred PA Required	
Eszopiclone tablet	AMBIEN (zolpidem) tablet	Non-preferred non-benzodiazepine sedative hypnotics may be approved for members who have failed treatment with two preferred non-benzodiazepine agents (failure is defined as lack of efficacy with a 2-week trial, allergy, intolerable side effects, or significant drug-drug interaction). <u>Children:</u> Prior authorization will be required for all agents for members < 18 years of age. <u>Duplications:</u> Only one agent in the sedative hypnotic drug class will be approved at a time (concomitant use of agents in the same sedative hypnotic class or differing classes will not be approved). All sedative hypnotics will require prior authorization for members ≥ 65 years of age when exceeding 90 days of therapy. Belsomra (suvorexant) may be approved for adult members that meet the following:
Ramelteon tablet	AMBIEN CR (zolpidem ER) tablet	
Zaleplon capsule	BELSOMRA (suvorexant) tablet	
Zolpidem IR, ER tablet	DAYVIGO (lemoborexant) tablet	
	Doxepin tablet	
	EDLUAR (zolpidem) SL tablet	
	HETLIOZ (tasimelteon) capsule	

	HETLIOZ LQ (tasimelteon) suspension LUNESTA (eszopiclone) tablet QUVIVIQ (daridorexant) tablet ROZEREM (ramelteon) tablet SILENOR (doxepin) tablet Tasimelteon capsule Zolpidem capsule, SL tablet	• Member has trialed and failed therapy with two preferred agents (failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) AND • Member is not receiving strong CYP3A4 inhibitors (such as erythromycin, clarithromycin, telithromycin, itraconazole, ketoconazole, posaconazole, fluconazole, voriconazole, delavirdine, and milk thistle) or strong CYP3A4 inducers (such as carbamazepine, oxcarbazepine, phenobarbital, phenytoin, rifampin, rifabutin, rifapentine, dexamethasone, efavirenz, etravirine, nevirapine, darunavir/ritonavir, ritonavir, and St John's Wort) AND • Member does not have a diagnosis of narcolepsy Dayvigo (lemborexant) may be approved for adult member that meet the following: • Member has trialed and failed therapy with two preferred agents AND Belsomra (suvorexant). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND • Member is not receiving strong CYP3A4 inhibitors (such as erythromycin, clarithromycin, telithromycin, itraconazole, ketoconazole, posaconazole, fluconazole, voriconazole, delavirdine, and milk thistle) or strong CYP3A4 inducers (such as carbamazepine, oxcarbazepine, phenobarbital, phenytoin, rifampin, rifabutin, rifapentine, dexamethasone, efavirenz, etravirine, nevirapine, darunavir/ritonavir, ritonavir, and St John's Wort) AND • Member does not have a diagnosis of narcolepsy Hetlioz (tasimelteon) capsules may be approved for members meeting the following criteria: • Member is ≥ 18 years of age and has a documented diagnosis of Non-24-hour sleep wake disorder (Non-24) OR • Member is ≥ 16 years of age and has a documented diagnosis of nighttime sleep disturbances in Smith-Magenis syndrome (SMS) AND • The requested medication is being prescribed by a sleep specialist or a practitioner who has sufficient education and experience to safely prescribe tasimelteon Hetlioz LQ (tasimelteon) oral suspension may be approved for members meeting the following criteria: • Member is 3 to 15 years of age and has a documented diagnosis of nighttime sleep disturbances in Smith-Magenis Syndrome (SMS) • AND the requested medication is being prescribed by a sleep specialist or a practitioner who has sufficient education and experience to safely prescribe tasimelteon. Silenor (doxepin) may be approved for adult members that meet ONE of the following criteria: • Member has tried and failed two preferred oral sedative hypnotics (Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) OR • Provider attests to the medical necessity of prescribing individual doxepin doses of less than 10 mg, OR
--	---	---

		Member's age is ≥ 65 years Prior authorization will be required for prescribed doses exceeding maximum (Table 1) below.
Benzodiazepines		
Preferred No PA Required* (Unless age, dose, or duplication criteria apply) Temazepam 15mg, 30mg capsule Triazolam tablet	Non-Preferred PA Required DORAL (quazepam) tablet Estazolam tablet Flurazepam capsule HALCION (triazolam) tablet Quazepam tablet RESTORIL (temazepam) capsule Temazepam 7.5mg, 22.5mg capsule	Non-preferred benzodiazepine sedative hypnotics may be approved for members who have trialed and failed therapy with two preferred benzodiazepine agents (failure is defined as lack of efficacy with a 2-week trial, allergy, intolerable side effects, or significant drug-drug interaction). Temazepam 22.5 mg may be approved if the member has trialed and failed temazepam 15mg or 30mg AND one other preferred product (failure is defined as lack of efficacy with a 2-week trial, allergy, intolerable side effects, or significant drug-drug interaction). Temazepam 7.5 mg may be approved if provider attests to the medical necessity of prescribing individual temazepam doses of less than 15 mg. <u>Children:</u> Prior authorization will be required for all sedative hypnotic agents when prescribed for members < 18 years of age. <u>Duplications:</u> Only one agent in the sedative hypnotic drug class will be approved at a time (concomitant use of agents in the same sedative hypnotic class or differing classes will not be approved). All sedative hypnotics will require prior authorization for member's ≥ 65 years of age when exceeding 90 days of therapy. Members currently stabilized on a non-preferred benzodiazepine medication may receive authorization to continue that medication. Prior authorization will be required for prescribed doses exceeding maximum (Table 1).

Table 1: Sedative Hypnotic Maximum Dosing		
Brand	Generic	Maximum Dose
Non-Benzodiazepine		
Ambien CR	Zolpidem CR	12.5 mg/day
Ambien IR	Zolpidem IR	10 mg/day
Belsomra	Suvorexant	20 mg/day
Dayvigo	Lemborexant	10 mg/day
Edluar	Zolpidem sublingual	10 mg/day
-	Zolpidem sublingual	Men: 3.5mg/day Women: 1.75 mg/day
Hetlioz	Tasimelteon capsule	20 mg/day
Hetlioz LQ	Tasimelteon liquid	≤ 28 kg: 0.7 mg/kg/day > 28 kg : 20 mg/day

Lunesta	Eszopiclone	3 mg/day
Quviviq	Daridorexant	50 mg/day
-	Zaleplon	20 mg/day
Rozerem	Ramelteon	8 mg/day
Benzodiazepine		
Halcion	Triazolam	0.5 mg/day
Restoril	Temazepam	30 mg/day
Silenor	Doxepin	6mg/day
-	Estazolam	2 mg/day
-	Flurazepam	30 mg/day
Doral	Quazepam	15 mg/day

Therapeutic Drug Class: SKELETAL MUSCLE RELAXANTS – Effective 4/1/2025

No PA Required (*if under 65 years of age)	PA Required	
Baclofen tablet Cyclobenzaprine tablet Methocarbamol tablet Tizanidine tablet	AMRIX ER (cyclobenzaprine ER) capsule Baclofen solution, suspension Carisoprodol tablet Carisoprodol/Aspirin tablet Chlorzoxazone tablet Cyclobenzaprine ER capsule DANTRIUM (dantrolene) capsule *Dantrolene capsule FEXMID (cyclobenzaprine) tablet FLEQSUVY (baclofen) solution LORZONE (chlorzoxazone) tablet LYVISPAH (baclofen) granules Metaxalone tablet NORGESIC/NORGESIC FORTE (orphenadrine/aspirin/ caffeine) tablet	All agents in this class will require a PA for members 65 years of age and older. The maximum allowable approval will be for a 7-day supply. Authorization for any CARISOPRODOL product will be given for a maximum 3-week one-time authorization for members with acute, painful musculoskeletal conditions who have failed treatment with three preferred products within the last 6 months. *Dantrolene may be approved for members who have trialed and failed‡ one preferred agent and meet the following criteria: • Documentation of age-appropriate liver function tests AND • One of following diagnoses: Multiple Sclerosis, Cerebral Palsy, stroke, upper motor neuron disorder, or spinal cord injury • Dantrolene will be approved for the period of one year • If a member is stabilized on dantrolene, they may continue to receive approval All other non-preferred skeletal muscle relaxants may be approved for members who have trialed and failed‡ three preferred agents. ‡Failure is defined as: lack of efficacy with 14-day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions.

	Orphenadrine ER tablet Orphenadrine/Aspirin/Caffeine tablet SOMA (carisoprodol) tablet Tizanidine capsule ZANAFLEX (tizanidine) capsule, tablet	
Therapeutic Drug Class: STIMULANTS AND RELATED AGENTS – Effective 4/1/2025		
Preferred *No PA Required (if age, max daily dose, and diagnosis met) <i>Brand/generic changes effective 08/08/2024</i> Amphetamine salts, mixed ER (generic Adderall XR) capsule Amphetamine salts, mixed (generic Adderall IR) tablet Armodafinil tablet Atomoxetine capsule Clonidine ER tablet DAYTRANA^{BNR} (methylphenidate) patch Dexmethylphenidate IR tablet Dexmethylphenidate ER capsule Guanfacine ER tablet Methylphenidate (generic Methylin/Ritalin) solution, tablet Methylphenidate ER tablet (generic Concerta)	Non-Preferred PA Required ADDERALL IR (amphetamine salts, mixed IR) tablet ADDERALL XR (amphetamine salts, mixed ER) capsule ADZENYS XR-ODT (amphetamine) Amphetamine tablet (generic Evekeo) APTENSIO XR (methylphenidate ER) capsule AZSTARYS (serdexmethylphenidate/dexmethylphenidate) capsule CONCERTA (methylphenidate ER) tablet COTEMPLA XR-ODT (methylphenidate ER) DESOXYN (methamphetamine) tablet DEXEDRINE (dextroamphetamine) Spansule Dextroamphetamine ER capsule, solution, tablet DYANAVEL XR (amphetamine) suspension, tablet EVEKEO (amphetamine) ODT, tablet FOCALIN (dexmethylphenidate) tablet, XR capsule	*Preferred medications may be approved through AutoPA for indications listed in Table 1 (preferred medications may also receive approval for off-label use for fatigue associated with multiple sclerosis). Non-preferred medications may be approved for members meeting the following criteria (for Sunosi (solriamfetol) and Wakix (pitolisant), refer to specific criteria listed below): • Prescription meets indication/age limitation criteria (Table 1) AND • <u>If member is ≥ 6 years of age:</u> ○ Has documented trial and failure‡ with three preferred products in the last 24 months AND ○ If the member is unable to swallow solid oral dosage forms, two of the trials must be methylphenidate solution, dexmethylphenidate ER, Vyvanse, Adderall XR, or any other preferred product that can be taken without the need to swallow a whole capsule. OR • <u>If member is 3–5 years of age:</u> ○ Has documented trial and failure‡ with one preferred product in the last 24 months AND ○ If the member is unable to swallow solid oral dosage forms, the trial must be methylphenidate solution, dexmethylphenidate ER, Vyvanse, Adderall XR, or any other preferred product that can be taken without the need to swallow a whole capsule. SUNOSI (solriamfetol) prior authorization may be approved if member meets the following criteria: • Member is 18 years of age or older AND • Member has diagnosis of either narcolepsy or obstructive sleep apnea (OSA) and is experiencing excessive daytime sleepiness AND • Member does not have end stage renal disease AND • If Sunosi is being prescribed for OSA, member has 1 month trial of CPAP AND • Member has trial and failure‡ of modafinil AND armodafinil AND one other agent in stimulant PDL class.

Modafinil tablet VYVANSE^{BNR} (lisdexamfetamine) capsule	INTUNIV (guanfacine ER) tablet JORNAY PM (methylphenidate) capsule Lisdexamfetamine capsule, chewable tablet Methamphetamine tablet METHYLIN (methylphenidate) solution Methylphenidate CD/ER/LA capsule, chewable tablet, ER tablet (generic Relexxi/Ritalin), patch MYDAYIS ER (dextroamphetamine/amphetamine) capsule NUVIGIL (armodafinil) tablet ONYDA XR (Clonidine) suspension PROCENTRA (dextroamphetamine) solution PROVIGIL (modafinil) tablet QELBREE (viloxazine ER) capsule QUILLICHEW ER (methylphenidate) chewable tablet, XR suspension RELEXXII (methylphenidate ER) tablet RITALIN (methylphenidate) IR/ER tablet, ER capsule STRATTERA (atomoxetine) capsule SUNOSI (solriamfetol) tablet VYVANSE (lisdexamfetamine) chewable tablet WAKIX (pitolisant) tablet XELSTRYM (dextroamphetamine) patch	WAKIX (pitolisant) prior authorization may be approved if member meets the following criteria: • Member is 6 years of age or older AND • Member has diagnosis of narcolepsy and is experiencing excessive daytime sleepiness AND • Member does not have end stage renal disease (eGFR <15 mL/minute) AND • Member does not have severe hepatic impairment AND • Member has trial and failure[‡] of modafinil AND armodafinil AND one other agent in the stimulant PDL class AND • Member has been counseled that Wakix may reduce the efficacy of hormonal contraceptives and counseled regarding use of an alternative non-hormonal method of contraception during Wakix therapy and for at least 21 days after discontinuing treatment. Maximum Dose (all products): See Table 2 Exceeding Maximum Dose: Prior authorization may be approved for doses that are higher than the listed maximum dose (Table 2) for members meeting the following criteria: • Member is taking medication for indicated use listed in Table 1 AND • Member has 30-day trial and failure[‡] of three different preferred or non-preferred agents at maximum doses listed in Table 2 AND • Documentation of member's symptom response to maximum doses of three other agents is provided AND • Member is not taking a sedative hypnotic medication (such as temazepam, triazolam, or zolpidem from the Sedative Hypnotic PDL class). [‡]Failure is defined as: lack of efficacy with 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction.
---	--	---

	ZENZEDI (dextroamphetamine) tablet	
Table 1: Diagnosis and Age Limitations		
- Approval for medically accepted indications <u>not</u> listed in Table 1 may be given with prior authorization review and may require submission of peer-reviewed literature or medical compendia showing safety and efficacy of the medication used for the prescribed indication. - Preferred medications may also receive approval for off-label use for fatigue associated with multiple sclerosis if meeting all other criteria for approval. Bolded drug names are preferred (subject to preferential coverage changes for brand/generic equivalents)		
Drug	Diagnosis and Age Limitations	
Stimulants–Immediate Release		
Amphetamine sulfate (EVEKEO)	ADHD (Age ≥ 3 years), Narcolepsy (Age ≥ 6 years)	
Dexmethylphenidate IR (FOCALIN)	ADHD (Age ≥ 6 years)	
Dextroamphetamine IR tablet (ZENZEDI)	ADHD (Age 3 to 16 years), Narcolepsy (Age ≥ 6 years)	
Dextroamphetamine solution (PROCENTRA)	ADHD (Age 3 to 16 years), Narcolepsy (Age ≥ 6 years)	
Methamphetamine (DESOXYN)	ADHD (Age ≥ 6 years)	
methylphenidate IR (generic METHYLIN, RITALIN)	ADHD (Age ≥ 6 years [†]), Narcolepsy (Age ≥ 6 years), OSA. [†] Prior Authorization for members 3-6 years of age with a diagnosis of ADHD may be approved with prescriber attestation to the following: - Member’s symptoms have not significantly improved despite adequate behavior interventions AND - Member experiences moderate-to-severe continued disturbance in functioning AND - Prescriber has determined that the potential benefits of starting methylphenidate before the age of 6 years outweigh the potential harm of delaying treatment.	
Mixed amphetamine salts IR (generic ADDERALL)	ADHD (Age ≥ 3 years), Narcolepsy (Age ≥ 6 years)	
Stimulants –Extended-Release		
Amphetamine ER (ADZENYS XR-ODT and ADZENYS ER suspension)	ADHD (Age ≥ 6 years)	
Amphetamine ER (DYANAVEL XR)	ADHD (Age ≥ 6 years)	
Mixedamphetamine salts ER (ADDERALL XR)	ADHD (Age ≥ 6 years)	
Dexmethylphenidate ER (generic Focalin XR)	ADHD (Age ≥ 6 years)	
Dextroamphetamine ER (DEXEDRINE)	ADHD (Age 6 to 16 years), Narcolepsy (Age ≥ 6 years)	
Dextroamphetamine ER/amphetamine ER (MYDAYIS ER)	ADHD (Age ≥ 13 years)	
Dextroamphetamine ER patch (XELSTRYM)	ADHD (Age ≥ 6 years)	
Lisdexamfetamine dimesylate (VYVANSE capsule, Vyvanse chewable)	ADHD (Age ≥ 6 years), Moderate to severe binge eating disorder in adults (Age ≥ 18 years)	
Methylphenidate ER OROS (CONCERTA)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years), OSA	
Methylphenidate patch (DAYTRANA)	ADHD (Age ≥ 6 years)	
Methylphenidate SR (METADATE ER)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years)	
Methylphenidate ER (METADATE CD)	ADHD (Age ≥ 6 years)	

Methylphenidate ER (QUILLICHEW ER)	ADHD (Age 6 years to ≤ 65 years), Narcolepsy (Age ≥ 6 years)
Methylphenidate ER (QUILLIVANT XR)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years)
Methylphenidate ER (RELEXXI ER)	ADHD (Age 6 to 65 years)
Methylphenidate ER (RITALIN LA)	ADHD (Age ≥ 6 years) [†] Prior Authorization for members 4-6 years of age with a diagnosis of ADHD may be approved with prescriber attestation to the following: • Member's symptoms have not significantly improved despite adequate behavior interventions AND • Member experiences moderate-to-severe continued disturbance in functioning AND Prescriber has determined that the potential benefits of starting methylphenidate before the age of 6 years outweigh the potential harm of delaying treatment.
Methylphenidate ER (ADHANSIA XR)	ADHD (Age ≥ 6 years)
Methylphenidate ER (JORNAY PM)	ADHD (Age ≥ 6 years)
Methylphenidate XR (APTENSIO XR)	ADHD (Age ≥ 6 years)
Methylphenidate XR ODT (COTEMPLA XR-ODT)	ADHD (Age 6 to 17 years)
Serdexmethylphenidate/dexmethylphenidate (AZSTARYS)	ADHD (Age ≥ 6 years)
Non-Stimulants	
Atomoxetine (generic STRATTERA)	ADHD (Age ≥ 6 years)
Clonidine ER	ADHD as monotherapy OR adjunctive therapy to stimulants (Age ≥ 6 years)
Guanfacine ER (generic INTUNIV)	ADHD as monotherapy OR adjunctive therapy to stimulants (Age ≥ 6 years)
Viloxazine ER (QELBREE)	ADHD (Age ≥ 6 years)
Wakefulness-promoting Agents	
Armodafinil (generic NUVIGIL)	Excessive sleepiness associated with narcolepsy, OSA, SWD, and adjunct therapy to treat fatigue and sleepiness in patients with major depressive disorder (MDD) (Age ≥ 18 years)
Modafinil (PROVIGIL)	Excessive sleepiness associated with narcolepsy, OSA, SWD, and adjunct therapy to treat fatigue and sleepiness in patients with major depressive disorder (MDD), antipsychotic medication-related fatigue (Age ≥ 18 years)
Pitolisant (WAKIX)	Excessive sleepiness associated with narcolepsy (Age ≥ 6 years)
Solriamfetol (SUNOSI)	Excessive sleepiness associated with narcolepsy, OSA (Age ≥ 18 years)
KEY: ADHD —attention-deficit/hyperactivity disorder, OSA —obstructive sleep apnea, SWD —shift work disorder	

Table 2: Maximum Dose

Drug	Maximum Daily Dose
ADDERALL	60 mg
ADDERALL XR	60 mg
ADHANSIA XR	85 mg
ADZENYS XR ODT	18.8 mg (age 6-12)

ADZENYS ER SUSPENSION	12.5 mg (age ≥ 13)
AMPHETAMINE SALTS	40 mg
APTENSIO XR	60 mg
CONCERTA	54 mg (age 6-12) or 72 mg (≥ age 13)
AZSTARYS	52.3 mg serdexmethylphenidate and 10.4 mg dexmethylphenidate
CLONIDINE ER	0.4 mg
COTEMPLA XR-ODT	51.8 mg
DEXTROAMPHETAMINE ER	60 mg
DAYTRANA	30 mg/9 hour patch (3.3 mg/hr)
DESOXYN	25 mg
DEXEDRINE	60 mg
DYANAVEL XR	20 mg
EVEKEO	60 mg
FOCALIN	20 mg
FOCALIN XR	40 mg
GUANFACINE ER	4 mg (age 6-12) or 7 mg (age ≥ 13)
INTUNIV ER	4 mg (age 6-12) or 7 mg (age ≥ 13)
JORNAY PM	100 mg
METADATE CD	60 mg
METADATE ER	60 mg
METHYLIN	60 mg
METHYLIN ER	60 mg
METHYLIN SUSPENSION	60 mg
METHYLPHENIDATE	60 mg
METHYLPHENIDATE ER	60 mg
MYDAYIS ER	25 mg (age 13-17) or 50 mg (age ≥ 18)
NUVIGIL	250 mg
PROCENTRA	60 mg
PROVIGIL	400 mg
QELBREE	400 mg (age 6-17) or 600 mg (age ≥ 18)
QUILLICHEW ER	60 mg
QUILLIVANT XR	60 mg
RELEXXII	54 mg (ages 6-12) or 72 mg (≥ age 13)
RITALIN IR	60 mg
RITALIN SR	60 mg
RITALIN LA	60 mg
STRATTERA	100mg
SUNOSI	150 mg
VYVANSE CAPSULES AND CHEWABLE TABLETS	70 mg
WAKIX	35.6 mg
XELSTRYM ER PATCH	18 mg/9 hours

ZENZEDI		60 mg											
Therapeutic Drug Class: TRIPTANS, DITANS AND OTHER MIGRAINE TREATMENTS - Oral – <i>Effective 4/1/2025</i>													
No PA Required (Quantity limits may apply) Eletriptan tablet (generic Relpax) Naratriptan tablet (generic Amerge) Rizatriptan tablet, ODT (generic Maxalt) Sumatriptan tablet (generic Imitrex) Zolmitriptan tablet (generic Zomig)	PA Required Almotriptan tablet FROVA (frovatriptan) tablet Frovatriptan tablet IMITREX (sumatriptan) tablet MAXALT/MAXALT MLT (rizatriptan) tablet, ODT RELPAX (eletriptan) tablet REYVOW (lasmiditan) tablet Sumatriptan/Naproxen tablet SYMBRAVO (rizatriptan/meloxicam) tablet Zolmitriptan ODT ZOMIG (zolmitriptan) tablet	Reyvow (lasmiditan) may be approved if meeting the following: - Member has trialed and failed three preferred products OR member is unable to use triptan therapy due to cardiovascular risk factors AND - Member has trialed and failed two preferred agents in the CGRP Inhibitors drug class indicated for the acute treatment of migraine. All other non-preferred oral products may be approved for members who have trialed and failed three preferred oral products. Failure is defined as lack of efficacy with 4-week trial, allergy, documented contraindication to therapy, intolerable side effects, or significant drug-drug interaction. Quantity Limits: <table><tr><td>Amerge (naratriptan), Frova (frovatriptan), Imitrex (sumatriptan), Zomig (zolmitriptan)</td><td>9 tabs/30 days</td></tr><tr><td>Treximet (sumatriptan/naproxen)</td><td>9 tabs/30 days</td></tr><tr><td>Axert (almotriptan) and Relpax (eletriptan)</td><td>6 tabs/30 days</td></tr><tr><td>Maxalt (rizatriptan)</td><td>12 tabs/30 days</td></tr><tr><td>Reyvow (lasmiditan)</td><td>8 tabs/30 days</td></tr></table>		Amerge (naratriptan), Frova (frovatriptan), Imitrex (sumatriptan), Zomig (zolmitriptan)	9 tabs/30 days	Treximet (sumatriptan/naproxen)	9 tabs/30 days	Axert (almotriptan) and Relpax (eletriptan)	6 tabs/30 days	Maxalt (rizatriptan)	12 tabs/30 days	Reyvow (lasmiditan)	8 tabs/30 days
Amerge (naratriptan), Frova (frovatriptan), Imitrex (sumatriptan), Zomig (zolmitriptan)	9 tabs/30 days												
Treximet (sumatriptan/naproxen)	9 tabs/30 days												
Axert (almotriptan) and Relpax (eletriptan)	6 tabs/30 days												
Maxalt (rizatriptan)	12 tabs/30 days												
Reyvow (lasmiditan)	8 tabs/30 days												
Therapeutic Drug Class: TRIPTANS, DITANS, AND OTHER MIGRAINE TREATMENTS - Non-Oral – <i>Effective 4/1/2025</i>													
No PA Required (Quantity limits may apply) Dihydroergotamine nasal spray IMITREX (sumatriptan) nasal spray Sumatriptan cartridge, pen injector MIGRANAL ^{BNR} (dihydroergotamine) nasal spray Sumatriptan nasal spray*, vial	PA Required Dihydroergotamine injection IMITREX (sumatriptan) cartridge, pen injector TOSYMRA (sumatriptan) nasal spray TRUDHESA (dihydroergotamine) nasal spray ZEMBRACE SYMTOUCH (sumatriptan) auto-injector Zolmitriptan nasal spray ZOMIG (zolmitriptan) nasal spray	Zembrace Symtouch injection, Tosymra nasal spray, or Onzetra Xsail nasal powder may be approved for members who have trialed and failed one preferred non-oral triptan products AND two oral triptan agents with different active ingredients. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects, significant drug-drug interaction, or documented inability to take alternative dosage form. All other non-preferred products may be approved for members who have trialed and failed one preferred non-oral triptan product AND one preferred oral triptan product. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions, documented inability to tolerate dosage form. Quantity Limits: <table><tr><td>Dihydroergotamine mesylate vial 1mg/mL</td><td>24 vials/ 28 days</td></tr><tr><td>Imitrex (sumatriptan) injection</td><td>4 injectors / 30 days</td></tr><tr><td>Imitrex (sumatriptan) nasal spray</td><td>6 inhalers / 30 days</td></tr></table>		Dihydroergotamine mesylate vial 1mg/mL	24 vials/ 28 days	Imitrex (sumatriptan) injection	4 injectors / 30 days	Imitrex (sumatriptan) nasal spray	6 inhalers / 30 days				
Dihydroergotamine mesylate vial 1mg/mL	24 vials/ 28 days												
Imitrex (sumatriptan) injection	4 injectors / 30 days												
Imitrex (sumatriptan) nasal spray	6 inhalers / 30 days												

		Migranal (dihydroergotamine mesylate) nasal spray	8 nasal spray devices/ 30 days
		Onzetra Xsail (sumatriptan) nasal powder	16 nosepieces / 30 days
		Tosymra (sumatriptan) nasal spray	12 nasal spray devices / 30 days
		Zembrace Symtouch (sumatriptan) injection	36mg / 30 days
		Zomig (zolmitriptan) nasal spray	6 inhalers / 30 days
		Members currently utilizing a non-oral dihydroergotamine product formulation (based on recent claims history) may receive one year approval to continue therapy with that medication.	

V. Dermatological

Therapeutic Drug Class: ACNE AGENTS– Topical – Effective 10/1/2025

Preferred No PA Required (if age and diagnosis criteria are met*)	Non-Preferred PA Required	Authorization will not be approved for acne agents prescribed solely for cosmetic purposes.
*Adapalene gel	ACANYA (clindamycin/benzoyl peroxide) gel, pump	Preferred topical clindamycin and erythromycin products may be approved by AutoPA verification of ICD-10 diagnosis code for acne vulgaris, psoriasis, cystic acne, comedonal acne, disorders of keratinization, neoplasms, folliculitis, hidradenitis suppurativa, or perioral dermatitis (erythromycin only). Approval of preferred topical clindamycin and erythromycin products for other medically accepted indications may be considered following clinical prior authorization review by a call center pharmacist.
*Adapalene/benzoyl peroxide gel (generic Epiduo), gel pump (generic Epiduo Forte)	Adapalene cream, gel pump, solution ALTRENO (tretinoin) lotion	
*Clindamycin phosphate gel, lotion, solution, medicated swab/pledget	ARAZLO (tazarotene) lotion ATRALIN (tretinoin) gel	All other preferred topical acne agents may be approved if meeting the following criteria: - For members > 25 years of age, may be approved following prescriber verification that the medication is not being utilized for cosmetic purposes AND prescriber verification that the indicated use is for acne vulgaris, psoriasis, cystic acne, disorders of keratinization, neoplasms, or comedonal acne. These medications are only eligible for prior authorization approval for the aforementioned diagnoses. - For members ≤ 25 years of age, may be approved for a diagnosis of acne vulgaris, psoriasis, cystic acne, disorders of keratinization, neoplasms, or comedonal acne. Diagnosis will be verified through automated verification (AutoPA) of the appropriate corresponding ICD-10 diagnosis code related to the indicated use of the medication.
*Clindamycin/benzoyl peroxide gel jar (generic Benzaclin)	BENZAMYCIN (erythromycin/benzoyl peroxide) gel	
*Clindamycin/benzoyl peroxide gel tube (generic Duac)	BP (sulfacetamide sodium/sulfur/urea) cleansing wash	Non-preferred topical products may be approved for members meeting all of the following criteria: - Member has trialed/failed three preferred topical products with different mechanisms (such as tretinoin, antibiotic). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND - Prescriber verification that the medication is being prescribed for one of the following diagnoses: acne vulgaris, psoriasis, cystic acne, disorders of keratinization, neoplasms, or comedonal acne.
*Dapsone gel	CABTREO (adapalene/benzoyl peroxide/clindamycin) gel	
*Erythromycin solution	CLEOCIN-T (clindamycin) lotion	
*Erythromycin/Benzoyl peroxide gel (generic Benzamycin)	CLINDACIN ETZ/PAC (clindamycin phosphate) kit	
*Sulfacetamide sodium suspension	CLINDAGEL gel	
*Tretinoin cream	Clindamycin phosphate foam	

*Tretinoin gel (Mylan only)	Clindamycin/Benzoyl peroxide gel pump Clindamycin/tretinoin gel Dapsone gel pump ERY/ERYGEL (erythromycin/ethanol) gel, medicated swabs/pads Erythromycin gel EVOCLIN (clindamycin) foam FABIOR (tazarotene) foam KLARON (sulfacetamide) suspension NEUAC (clindamycin/benzoyl peroxide/emollient) kit ONEXTON (clindamycin/benzoyl peroxide) gel, gel pump RETIN-A MICRO (tretinoin) (all products) ROSULA (sulfacetamide sodium/sulfur) cloths, wash SSS 10-5 (sulfacetamide sodium/sulfur) foam Sulfacetamide sodium cleanser, cleansing gel, lotion, shampoo, wash Sulfacetamide sodium/sulfur cleanser, cream, pad, suspension, wash SUMADAN/XLT (sulfacetamide sodium/sulfur) kit, wash SUMAXIN/ CP/TS (sulfacetamide sodium/sulfur) kit, pads, suspension, wash Tazarotene cream, foam, gel Tretinoin gel (all other manufacturers)	
-----------------------------	---	--

	Tretinoin microspheres (all products) WINLEVI (clascoterone) cream ZIANA (clindamycin/tretinoin) gel	
Therapeutic Drug Class: ACNE AGENTS– ORAL ISOTRETINOIN – <i>Effective 7/1/2025</i>		
PA Required for all agents		Preferred products may be approved for adults and children ≥ 12 years of age for treating severe acne vulgaris or for treating moderate acne vulgaris in members unresponsive to conventional therapy. Non-preferred products may be approved for members meeting the following: - Member has trialed/failed one preferred agent (failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) AND - Member is an adult or child ≥ 12 years of age with severe, recalcitrant nodulocystic acne and has been unresponsive to conventional therapy.
Preferred AMNESTEEM capsule CLARAVIS capsule Isotretinoin 10 mg, 20 mg, 30 mg, 40 mg capsule (<i>Mayne-Pharma, Upsher-Smith, Zydus only</i>) ZENATANE capsule	Non-Preferred ABSORICA capsule ABSORICA LD capsule Isotretinoin 10 mg, 20 mg, 30 mg, 40 mg capsule (<i>All manufacturers except Mayne-Pharma, Upsher-Smith, Zydus</i>) Isotretinoin 25 mg, 35 mg capsule MYORISAN capsule	
Therapeutic Drug Class: ANTI-PSORIATICS - Oral – <i>Effective 7/1/2025</i>		
No PA Required Acitretin capsule	PA Required Methoxsalen capsule	Prior authorization for non-preferred oral agents may be approved with failure of two preferred anti-psoriatic agents, one of which must be a preferred oral agent. Failure is defined as lack of efficacy of a 4-week trial, allergy, intolerable side effects or significant drug-drug interaction.
Therapeutic Drug Class: ANTI-PSORIATICS -Topical – <i>Effective 7/1/2025</i>		
No PA Required Calcipotriene cream, foam, ointment, solution Calcipotriene/betamethasone dipropionate ointment	PA Required Calcipotriene/betamethasone dipropionate suspension Calcitriol ointment DUOBRII (halobetasol/tazarotene) lotion ENSTILAR (calcipotriene/betamethasone) foam	Preferred and non-preferred products that contain a corticosteroid ingredient (such as betamethasone) will be limited to 4 weeks of therapy. Continued use will require one week of steroid-free time in between treatment periods. Non-preferred topical agents may be approved with failure of two preferred topical agents. If non-preferred topical agent being requested is a combination product, trial of two preferred agents must include a preferred combination agent. Failure is defined as lack of efficacy of a 4-week trial, allergy, intolerable side effects or significant drug-drug interaction.

TACLONEX SCALP ^{BNR} (calcipotriene/betamethasone) suspension TACLONEX (calcipotriene/betamethasone) ointment	SORILUX (calcipotriene) foam VTAMA (tapinarof) cream ZORYVE 0.3% (roflumilast) cream, 0.3% foam	Members with >30% of their body surface area affected may not use Enstilar (calcipotriene/betamethasone DP) foam or Taclonex (calcipotriene/betamethasone DP) ointment products as safety and efficacy have not been established. ZORYVE (roflumilast) 0.3% cream may receive approval if meeting the following based on prescribed indication: <u>Plaque psoriasis</u> (0.3% cream formulation only): • Member is ≥ 6 years of age AND • Member has a diagnosis of plaque psoriasis AND • Member has body surface area (BSA) involvement of $\leq 20\%$ AND • Member does not have moderate or severe hepatic impairment (Child-Pugh B or C) AND • Medication is being prescribed by or in consultation with a dermatologist AND • <u>If the affected area is limited to the scalp:</u> ○ Prescriber attests that member has been counseled regarding alternative treatment options, including over-the-counter (OTC) emollients, vitamin D analogs, and coal tar shampoo when appropriate AND ○ Member has documented trial and failure (with a minimum 2-week treatment period) of a topical corticosteroid. Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction. • <u>If the affected area includes the face or body:</u> ○ Member has documented trial and failure (with a minimum 2-week treatment period) of at least one product from ALL of the following categories. (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction): ▪ Topical corticosteroid ▪ Topical calcineurin inhibitor (such as pimecrolimus, tacrolimus) Members may not apply Zoryve (roflumilast) cream to >20% of affected body surface area, as safety and efficacy have not been established. <u>Quantity limit:</u> 60 grams/30 days <u>Initial approval:</u> 8 weeks <u>Reauthorization:</u> Reauthorization for one year may be approved based on provider attestation that member's symptoms improved during the initial 8 weeks of treatment and continuation of therapy is justified.
--	--	---

		<u>Plaque psoriasis (0.3% foam formulation only):</u> • Member is ≥ 12 years of age AND • Member has a diagnosis of plaque psoriasis AND • Member has body surface area (BSA) involvement of $\leq 20\%$ AND • Member does not have moderate or severe hepatic impairment (Child-Pugh B or C) AND • Medication is being prescribed by or in consultation with a dermatologist AND • If the affected area is limited to the scalp: ○ Prescriber attests that member has been counseled regarding alternative treatment options, including over-the-counter (OTC) emollients, vitamin D analogs, and coal tar shampoo when appropriate AND ○ Member has documented trial and failure (with a minimum 2-week treatment period) of a topical corticosteroid. Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction. • If the affected area includes the face or body: ○ Member has documented trial and failure (with a minimum 2-week treatment period) of at least one product from ALL of the following categories (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction): ▪ Topical corticosteroid ▪ Topical calcineurin inhibitor (such as pimecrolimus, tacrolimus) <u>Quantity limit:</u> 60 grams/30 days <u>Initial approval:</u> 8 weeks <u>Reauthorization:</u> Reauthorization for one year may be approved based on provider attestation that member's symptoms improved during the initial 8 weeks of treatment and continuation of therapy is justified.
Therapeutic Drug Class: IMMUNOMODULATORS, TOPICAL – <i>Effective 7/15/2025</i>		
Atopic Dermatitis		
No PA Required (Unless indicated*) ELIDEL (pimecrolimus) cream *EUCRISA (crisaborole) ointment *OPZELURA (ruxolitinib) cream	PA Required ANZUPGO (delgocitinib) cream VTAMA (tapinarof) 1% cream ZORYVE (roflumilast) 0.15% cream, 0.3% foam	EUCRISA (crisaborole) may be approved if the following criteria are met: • Member is ≥ 3 months of age AND • Member has a diagnosis of mild to moderate atopic dermatitis AND • Member tried and failed‡ one preferred agent OR one medium-to-very high potency topical corticosteroid AND • Eucrisa (crisaborole) is being prescribed by or in consultation with a dermatologist or allergist/immunologist.

Pimecrolimus cream Tacrolimus ointment		OPZELURA (ruxolitinib) cream may be approved if the following criteria are met based on prescribed indication: <u>Atopic Dermatitis</u> • Member is ≥ 12 years of age AND • Member has a diagnosis of mild to moderate atopic dermatitis AND • Medication is being prescribed by or in consultation with a dermatologist or allergist/immunologist AND • Member has trialed and failed‡ one preferred agent OR one medium potency to very high potency topical corticosteroid (such as mometasone furoate, betamethasone dipropionate, or fluocinonide) or prescriber verifies that member is not a candidate for topical corticosteroids. <u>Nonsegmental Vitiligo</u> • Member is ≥ 12 years of age AND • Member is immunocompetent AND • Member has a diagnosis of stable nonsegmental vitiligo, defined as no increase in the size of existing lesions and the absence of new lesions in the previous 3 to 6 months, AND • Medication is being prescribed by or in consultation with a dermatologist AND • Member has trialed and failed‡ one preferred agent AND one medium potency to very high potency topical corticosteroid (such as mometasone furoate, betamethasone dipropionate, or fluocinonide) or prescriber verifies that member is not a candidate for topical corticosteroids <u>Quantity limit:</u> 60 grams/week Non-preferred topical immunomodulator products may be approved for atopic dermatitis following adequate trial and failure‡ of one prescription topical corticosteroid AND two preferred agents. ZORYVE (roflumilast) 0.15% cream and 0.3% foam may receive approval if meeting the following based on prescribed indication: <u>Atopic dermatitis (0.15% cream formulation only):</u> • 6 years of age and older AND • Member has a diagnosis of mild atopic dermatitis in adult and pediatric patients AND • Request meets trial and failure criteria for non-preferred agents listed above <u>Seborrheic dermatitis (0.3% foam formulation only):</u>
--	--	--

		• Member is ≥ 9 years of age AND • Member has a diagnosis of seborrheic dermatitis AND • Member does not have moderate or severe hepatic impairment (Child-Pugh B or C) AND • Medication is being prescribed by or in consultation with a dermatologist AND • Member has been counseled that Zoryve foam is flammable. Fire, flame, or smoking during and immediately following application must be avoided. • <u>If the affected area is limited to the scalp:</u> ○ Prescriber attests that member has been counseled regarding alternative treatment options, including over-the-counter (OTC) antifungal shampoo (such as selenium sulfide, zinc pyrithione) and OTC coal tar shampoo, when appropriate) AND ○ Member has documented trial and failure (with a minimum 2-week treatment period) of at least one prescription product for seborrheic dermatitis, such as ketoconazole 2% antifungal shampoo or a topical corticosteroid. Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction. • <u>If the affected area includes the face or body:</u> ○ Member has documented trial and failure (with a minimum 2-week treatment period) with at least one product from ALL of the following categories (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction): ▪ Topical antifungal (such as ketoconazole, ciclopirox) ▪ Topical corticosteroid ▪ Topical calcineurin inhibitor (such as pimecrolimus, tacrolimus) <u>Quantity limit:</u> 60 grams/30 days <u>Initial approval:</u> 8 weeks Members may not apply Zoryve (roflumilast) cream to >20% of affected body surface area, as safety and efficacy have not been established. <u>Reauthorization:</u> Reauthorization for one year may be approved based on provider attestation that member's symptoms improved during the initial 8 weeks of treatment and continuation of therapy is justified.
--	--	---

		‡Failure is defined as a lack of efficacy with a 2-week trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction
Antineoplastic Agents		
Preferred No PA Required (Unless indicated*) *Diclofenac 3% gel (generic Solaraze) Fluorouracil 5% cream (generic Efudex) Fluorouracil 2%, 5% solution	Non-Preferred PA Required Bexarotene gel CARAC (fluorouracil) cream EFUDEX (fluorouracil) cream Fluorouracil 0.5% (generic Carac) cream PANRETIN (alitretinoin) gel TARGRETIN (bexarotene) gel VALCHLOR (mechlorethamine) gel	*Diclofenac 3% gel (generic Solaraze) may be approved if the member has a diagnosis of actinic keratosis (AK). TARGRETIN (bexarotene) gel or VALCHLOR (mechlorethamine) gel may be approved for members who meet the following criteria: • Member is ≥ 18 years of age AND • Member has been diagnosed with Stage IA or IB cutaneous T-cell lymphoma (CTCL) AND • Member has refractory or persistent CTCL disease after other therapies OR has not tolerated other therapies AND • Member and partners have been counseled on appropriate use of contraception Non-preferred agents may be approved for members who have failed an adequate trial of all preferred products FDA-approved for that indication. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.
Other Agents		
No PA Required Imiquimod (generic Aldara) cream Podofilox gel, solution	PA Required CONDYLOX (podofilox) gel HYFTOR (sirolimus) gel Imiquimod (generic Zyclara) cream, cream pump VEREGEN (sinecatechins) ointment ZYCLARA (imiquimod) cream, cream pump	Hyftor (sirolimus) gel • Member has a diagnosis of facial angiofibroma associated with tuberous sclerosis AND • Member is ≥ 6 years of age AND • Provider has evaluated, and member has received, all age-appropriate vaccinations as recommended by current immunization guidelines prior to initiating treatment with HYFTOR <u>Initial approval:</u> 6 months <u>Reauthorization:</u> An additional 6 months may be approved based on provider attestation that symptoms improved during the initial 6 months of treatment and the provider has assessed use of all vaccinations recommended by current immunization guidelines. <u>Maximum dose:</u> one 10-gram tube/28 days Veregen (sinecatechins) may be approved if the following criteria are met: • Member has a diagnosis of external genital and/or perianal warts (Condylomata acuminata) AND • Member is ≥ 18 years of age AND Member is immunocompetent AND

		- Member has tried and failed two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. Zyclara (imiquimod) 2.5% cream may be approved if the following criteria are met: - Member has a diagnosis of clinically typical visible or palpable actinic keratoses (AK) of the full face or balding scalp AND - Member is ≥ 18 years of age AND - Member is immunocompetent AND - Member has tried and failed one preferred product in the Antineoplastic Agents class (such as diclofenac gel or fluorouracil) AND the preferred imiquimod (generic Aldara) product. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. Zyclara (imiquimod) 3.75% cream may be approved for: - Treatment of clinically typical visible or palpable, actinic keratoses (AK) of the full face or balding scalp if the following criteria are met: - Member is ≥ 18 years of age AND - Member is immunocompetent AND - Member has tried and failed one preferred product from the Antineoplastic Agents class (such as diclofenac gel or fluorouracil) AND the preferred imiquimod (generic Aldara) product. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. OR - Treatment of external genital and/or perianal warts (Condylomata acuminata) if the following criteria are met: - Member is ≥ 12 years of age AND - Member has tried and failed two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. All other non-preferred products may be approved for members who have trialed and failed all preferred products that are FDA-approved for use for the prescribed indication. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. <u>Quantity Limits:</u> Aldara (imiquimod) cream has a quantity limit of 12 packets/28 days.
Therapeutic Drug Class: ROSACEA AGENTS – <i>Effective 7/1/2025</i>		
No PA Required	PA Required	
Azelaic acid gel	Brimonidine gel pump	Prior authorization for non-preferred products in this class may be approved if meeting the following criteria for the prescribed diagnosis:

FINACEA (azelaic acid) gel FINACEA (azelaic acid) foam Metronidazole cream, lotion Metronidazole 0.75% gel	*Doxycycline monohydrate DR capsule (generic Oracea) Ivermectin cream Metronidazole 1% gel, gel pump MIRVASO (Brimonidine gel pump) NORITATE (metronidazole) cream RHOFADE (oxymetazoline) cream ROSADAN (metronidazole/skin cleanser) cream kit, gel kit	<u>Rosacea:</u> - Member has a diagnosis of persistent (non-transient) facial erythema with inflammatory papules and pustules due to rosacea AND - Prescriber attests that medication is not being used solely for cosmetic purposes AND - Member has tried and failed two preferred agents of different mechanisms of action (Failure is defined as lack of efficacy with 4-week trial, allergy, contraindication, or intolerable side effects) <u>Demodex Blepharitis:</u> - Requests for non-preferred topical ivermectin cream may be approved for treatment of moderate to severe Demodex blepharitis *Doxycycline monohydrate DR (generic Oracea) may be approved if the following criteria are met: - Member has taken generic doxycycline for a minimum of three months and failed therapy in the last 6 months. Failure is defined as: lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions AND - Member has history of an adequate trial/failure (8 weeks) of 2 other preferred agents (oral or topical). Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions AND - Member is ≥ 18 years of age and has been diagnosed with rosacea with inflammatory lesions (papules and pustules)
Therapeutic Drug Class: TOPICAL STEROIDS – <i>Effective 7/1/2025</i>		
Low potency		
No PA Required DERMA-SMOOTH-FS (fluocinolone) 0.01% body oil/scalp oil^{BNR} Desonide 0.05% cream, ointment Fluocinolone 0.01% cream, 0.01% solution Hydrocortisone (Rx) cream, lotion, ointment	PA Required Alclometasone 0.05% cream, ointment CAPEX (fluocinolone) 0.01% shampoo Desonide 0.05% lotion Fluocinolone 0.01% body oil, 0.01% scalp oil PROCTOCORT (hydrocortisone) (Rx) 1% cream SYNALAR (fluocinolone) 0.01% solution	Non-preferred Low Potency topical corticosteroids may be approved following adequate trial and failure of two preferred agents in the Low Potency class (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).

	SYNALAR TS (fluocinolone/skin cleanser) Kit TEXACORT (hydrocortisone) 2.5% solution	
Medium potency		
No PA Required	PA Required	Non-preferred Medium Potency topical corticosteroids may be approved following adequate trial and failure of two preferred agents in the Medium Potency class (failure is defined as: lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
Betamethasone dipropionate 0.05% cream, lotion, ointment Betamethasone valerate 0.1% cream, ointment Fluocinolone 0.025% cream, 0.05% cream, 0.005% ointment Fluticasone cream, ointment Hydrocortisone valerate 0.2% cream Mometasone 0.1% cream, 0.1% ointment, 0.1% solution Triamcinolone acetonide 0.025% cream, 0.1% cream, 0.025% ointment, 0.05% ointment, 0.1% ointment, 0.025% lotion, 0.1% lotion Triamcinolone 0.1% dental paste	BESER (fluticasone) lotion, emollient kit Betamethasone valerate 0.1% lotion, 0.12% foam Clocortolone 0.1% cream, cream pump CLODERM (clocortolone) 0.1% cream, cream pump CUTIVATE (fluticasone) 0.05% cream, lotion Diflorasone 0.05% cream Fluocinolone 0.025% ointment Fluocinonide-E 0.05% cream Flurandrenolide 0.05% cream, lotion, ointment Fluticasone 0.05% lotion Hydrocortisone butyrate 0.1% cream, lotion, solution, ointment, lipid/lipocream Hydrocortisone valerate 0.2% ointment KENALOG (triamcinolone) spray LOCOID (hydrocortisone butyrate) 0.1% lotion LOCOID LIPOCREAM (hydrocortisone butyrate-emollient) 0.1% cream LUXIQ (betamethasone valerate) 0.12% foam ORALONE (Triamcinolone) 0.1% dental paste PANDEL (hydrocortisone probutate) 0.1% cream	

	Prednicarbate 0.1% cream, ointment PSORCON (diflorasone) 0.05% cream SYNALAR (fluocinolone) 0.025% cream/kit, ointment/kit Triamcinolone 0.147 mg/gm spray	
High potency		
No PA Required (*unless exceeds duration of therapy) * Betamethasone dipropionate 0.05% ointment *Betamethasone dipropionate/propylene glycol (augmented) 0.05% cream *Fluocinonide 0.05% cream, 0.05% gel, 0.05% ointment, 0.05% solution *Triamcinolone acetonide 0.5% cream, 0.5% ointment	PA Required Amcinonide 0.1% cream, lotion APEXICON-E (diflorasone/emollient) 0.05% cream Desoximetasone 0.05%, 0.25% cream, 0.05% gel, 0.05%, 0.25% ointment Diflorasone 0.05% ointment Halcinonide 0.1% cream HALOG (halcinonide) 0.1% cream, ointment, solution TOPICORT (desoximetasone) 0.05%, 0.25% cream, 0.05% gel, 0.05%, 0.25% ointment	Non-preferred High Potency topical corticosteroids may be approved following adequate trial and failure of two preferred agents in the High Potency class (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions). *All High Potency topical corticosteroids will require prior authorization beyond 4 weeks of therapy. The provider will be encouraged to transition to a medium or low potency topical steroid after this time has elapsed. Claims for compounded products containing high-potency topical steroids will be limited to a maximum of 60 grams or 60 mL of a high-potency ingredient per 4-week treatment period. Claims exceeding this quantity limit will require prior authorization with prescriber's justification for use of the product at the prescribed dose.
Very high potency		
No PA Required (Unless exceeds duration of therapy*) *Betamethasone dipropionate/propylene glycol (augmented), 0.05% lotion 0.05% ointment *Clobetasol 0.05% cream, 0.05% gel, 0.05% ointment, 0.05% solution *Fluocinonide 0.1% cream	PA Required Betamethasone dipropionate/propylene glycol (augmented) 0.05% gel BRYHALI (halobetasol) 0.01% lotion Clobetasol emollient/emulsion 0.05% cream, foam Clobetasol 0.05% lotion, foam, spray, shampoo CLODAN (clobetasol) 0.05% cleanser kit Desoximetasone 0.25% spray	Non-preferred Very High Potency topical corticosteroids may be approved following adequate trial and failure of clobetasol propionate in the same formulation as the product being requested (if the formulation of the requested non-preferred product is not available in preferred clobetasol product options, then trial and failure of any preferred clobetasol product formulation will be required). Failure is defined as lack of efficacy with 2-week trial, allergy, intolerable side effects or significant drug-drug interactions. *All Very High Potency topical corticosteroids will require prior authorization beyond 2 weeks of therapy. If clobetasol propionate shampoo is being used to treat plaque psoriasis, then prior authorization will be required beyond 4 weeks of therapy. The provider will be encouraged to transition to a medium or low potency topical steroid after this time has elapsed.

	DIPROLENE (betamethasone dipropionate/propylene glycol, augmented) 0.05% ointment Halobetasol 0.05% cream, foam, ointment IMPEKLO (clobetasol) 0.05% lotion LEXETTE (halobetasol) 0.05% foam OLUX (clobetasol) 0.05% foam TOPICORT (desoximetasone) 0.25% spray TOVET EMOLLIENT (clobetasol) 0.05% foam ULTRAVATE (halobetasol) 0.05% lotion VANOS (fluocinonide) 0.1% cream	
--	--	--

VI. Endocrine

Therapeutic Drug Class: **ANDROGENIC AGENTS, Topical, Injectable, Oral** – *Effective 10/1/2025*

PA Required for all agents in this class		
Preferred	Non-Preferred	
Testosterone cypionate IM injection	ANDROGEL (testosterone) gel packet	<u>Hypogonadotropic or Primary Hypogonadism (may be secondary to Klinefelter Syndrome):</u> Preferred products may be approved for members meeting the following: - Member is a male patient ≥ 16 years of age with a documented diagnosis of hypogonadotropic or primary hypogonadism OR ≥ 12 years of age with a diagnosis of hypogonadotropic or hypogonadism secondary to Klinefelter Syndrome (all other diagnoses will require manual review) AND - Member has two documented low serum testosterone levels below the lower limit of normal range for testing laboratory prior to initiation of therapy AND - Member does not have a diagnosis of breast or prostate cancer AND
Testosterone gel packet	ANDROGEL (testosterone) gel 1.62% pump	
Testosterone 1.62% gel pump	DEPO-TESTOSTERONE (testosterone cypionate) IM injection	
	JATENZO (testosterone undecanoate) capsule	
<i>Injectable testosterone cypionate is a pharmacy benefit when</i>	KYZATREX (testosterone undecanoate) capsule	

<i>self-administered.</i> <i>Administration in an office setting is a medical benefit.</i>	METHITEST (methyltestosterone) tablet Methyltestosterone capsule NATESTO (testosterone) nasal spray TESTIM (testosterone) gel Testosterone 1% gel tube, 30 mg/1.5 ml pump Testosterone enanthate IM injection TLANDO (testosterone undecanoate) capsule UNDECATREX (testosterone undecanoate) capsule XYOSTED (testosterone enanthate) SC injection	• If the member is > 40 years of age, has prostate-specific antigen (PSA) < 4 ng/mL or has no palpable prostate nodule AND • Member has baseline hematocrit < 50% Reauthorization Criteria (requests for renewal of a currently expiring prior authorization for a preferred product may be approved for members meeting the following criteria): • Member is a male patient ≥ 16 years of age with a documented diagnosis of hypogonadotropic or primary hypogonadism OR ≥ 12 years of age with a diagnosis of hypogonadotropic or hypogonadism secondary to Klinefelter Syndrome AND • Serum testosterone is being regularly monitored (at least annually) to achieve total testosterone level in the middle tertile of the normal reference range AND • Member does not have a diagnosis of breast or prostate cancer AND • Member has a hematocrit < 54% <u>Gender Transition/Affirming Hormone Therapy:</u> Preferred androgenic drugs may be approved for members meeting the following: 1. Female sex assigned at birth and has reached Tanner stage 2 of puberty AND 2. Is undergoing female to male transition AND 3. Has a negative pregnancy test prior to initiation AND 4. Hematocrit (or hemoglobin) is being monitored. Non-Preferred Products: Non-preferred topical androgenic agents may be approved for members meeting the above criteria with trial and failed$\ddagger$ therapy with two preferred topical androgen formulations. Non-preferred injectable androgenic agents may be approved for members meeting the above criteria with trial and failed$\ddagger$ therapy with a preferred injectable androgenic drug. Prior authorization for oral androgen agents (tablet, capsule, buccal) may be approved if member has trialed and failed$\ddagger$ therapy with a preferred topical agent AND testosterone cypionate injection. $\ddagger$Failure is defined as lack of efficacy with 8 week trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction. For all agents and diagnoses, members < 16 years of age will require a manual prior authorization review by a pharmacist (with exception of members ≥ 12 years of age with a diagnosis of hypogonadotropic or hypogonadism secondary to Klinefelter Syndrome).
Therapeutic Drug Class: BONE RESORPTION SUPPRESSION AND RELATED AGENTS – <i>Effective 10/1/2025</i>		
Bisphosphonates		
No PA Required Alendronate solution, tablet	PA Required ACTONEL (risedronate) tablet	

Ibandronate tablet	ATELVIA (risedronate) tablet	Non-preferred bisphosphonates may be approved for members who have failed treatment with one preferred product at treatment dose. Failure is defined as lack of efficacy with a 12-month trial, allergy, intolerable side effects, or significant drug-drug interaction. For members who have a low risk of fracture, discontinuation of bisphosphonate therapy and drug holiday should be considered following 5 years of treatment. Low risk is defined as having a bone mineral density, based on the most recent T-score, of greater than (better than) -2.5 AND no history of low trauma or fragility fracture.
Risedronate tablet	BINOSTO (alendronate) effervescent tablet	
	FOSAMAX (alendronate) tablet	
	FOSAMAX plus D (alendronate/vit D) tablet	
Non-Bisphosphonates		
PREFERRED	Non-Preferred	*FORTEO (teriparatide) or generic teriparatide may be approved if the following criteria are met: - Member has one of the following diagnoses: - Male primary or hypogonadal osteoporosis (BMD T-score of -2.5 or less) - Osteoporosis due to corticosteroid use - Postmenopausal osteoporosis AND - Member is at very high risk for fracture† OR member has history of trial and failure of one preferred bisphosphonate. Failure is defined as lack of efficacy with a 12-month trial, allergy, intolerable side effects, or significant drug-drug interaction. - Prior authorization will be given for one year and total exposure of parathyroid hormone analogs (teriparatide and abaloparatide) shall not exceed two years TYMLOS (abaloparatide) may be approved if the member meets the following criteria: - Member has a diagnosis of postmenopausal osteoporosis (BMD T-score of -2.5 or less) AND - Member is post-menopausal with very high risk for fracture† OR member has history of trial and failure of FORTEO (teriparatide). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. AND - Prior authorization will be given for one year and total exposure of parathyroid hormone analogs (teriparatide and abaloparatide) shall not exceed two years. All other non-preferred non-bisphosphonates may be approved for FDA-labeled indications for members who have failed treatment with one preferred bisphosphonate or non-bisphosphonate product at treatment dose. Failure is defined as lack of efficacy with a 12-month trial, allergy, unable to use oral therapy, intolerable side effects, or significant drug-drug interaction. †Members at very high risk for fracture: Members will be considered at very high risk for fracture if they meet <u>one</u> of the following: - A history of fracture within the past 12 months OR - Fractures experienced while receiving guideline-supported osteoporosis therapy OR - A history of multiple fractures OR
FORTEO (teriparatide) SC pen ^{BNR*}	BONSITY (teriparatide) SC pen	
Raloxifene tablet	Calcitonin salmon nasal spray	
	EVISTA (raloxifene) tablet	
	Teriparatide SC pen	
	TYMLOS (abaloparatide) SC pen	

- A history of fractures experienced while receiving medications that cause skeletal harm (such as long-term glucocorticoids) **OR**
- A very low T-score (less than -3.0) **OR**
- A high risk for falls or a history of injurious falls **OR**
- A very high fracture probability by FRAX (> 30% for a major osteoporosis fracture or > 4.5% for hip fracture).

Non-Bisphosphonate Product	FDA-approved Maximum Dose
Calcitonin salmon nasal spray	1 metered dose spray (200 units) daily
Evista (raloxifene) oral tablet	60 mg daily
Forteo (teriparatide) subcutaneous injection	20 mcg daily
Tymlos (abaloparatide) subcutaneous injection	80 mcg daily

Note: Prior authorization criteria for Prolia (denosumab) and other injectable bone resorption and related agents are listed on Appendix P.

Therapeutic Drug Class: CONTRACEPTIVES - Topical – Effective 07/10/2025

Effective 01/14/22, topical contraceptive patch products are eligible for coverage with a written prescription by an enrolled pharmacist. Additional information regarding pharmacist enrollment can be found at <https://hcpf.colorado.gov/pharm-serv>.

No PA Required	PA Required	
ANNOVERA (segesterone acetate/EE) vaginal ring	Etonorgestrel/EE vaginal ring (<i>all other manufacturers</i>)	Non-preferred topical contraceptive products may be approved following a trial and failure of one preferred topical contraceptive product. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.
Etonorgestrel/EE vaginal ring (<i>Prasco Labs</i>)	Norelgestromin/EE TD patch (generic XULANE)	*PHEXXI (lactic acid/citric/potassium) vaginal gel quantity limit: 120 grams per 30 days
*PHEXXI (lactic acid/citric/potassium) vaginal gel	NUVARING (etonorgestrel/EE) vaginal ring	Effective 7/1/2022: Prescriptions are eligible to be filled for up to a twelve-month supply.
TWIRLA (levonorgestrel/EE) TD patch	ZAFEMY (norelgestromin/EE) TD patch	<i>Note: IUD and select depot product formulations are billed through the medical benefit</i>

XULANE (norelgestromin/EE) TD patch		
Therapeutic Drug Class: DIABETES MANAGEMENT CLASSES, INSULINS – <i>Effective 02/27/2025</i>		
Rapid-Acting		
No PA Required HUMALOG (insulin lispro) cartridge, vial Insulin aspart cartridge, pen, vial Insulin lispro Kwikpen, Jr. Kwikpen, vial <i>(Eli Lilly)</i> NOVOLOG (insulin aspart) cartridge, FlexPen, vial	PA Required ADMELOG (insulin lispro) Solostar pen, vial AFREZZA (regular insulin) cartridge, unit APIDRA (insulin glulisine) Solostar pen, vial FIASP (insulin aspart) FlexPen, PenFill, pump cartridge, vial HUMALOG (insulin lispro) 200 U/mL Kwikpen HUMALOG Tempo Pen 100 U/mL HUMALOG 100U/mL KwikPen, vial HUMALOG Jr. (insulin lispro) KwikPen Insulin lispro 100 U/mL vial <i>(all other manufacturers)</i> KIRSTY (insulin aspart-xjhz) Kwikpen, vial, Tempo pen LYUMJEV (insulin lispro-aabc) Kwikpen, vial, Tempo pen MERILOG (insulin aspart-szjj) pen, vial	All non-preferred products may be approved following trial and failure of treatment with two preferred products, one of which is the same rapid-acting insulin analog (lispro or aspart) as the non-preferred product being requested. (Failure is defined as allergy [hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, and angioedema] or intolerable side effects). Afrezza (human insulin) may be approved if meeting the following criteria: • Member is 18 years or older AND • Member has trialed and failed treatment with two preferred products (failure is defined as allergy [hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, or angioedema] or intolerable side effects) AND • Member must not have chronic lung disease such as COPD or asthma AND • If member has type 1 diabetes, must use in conjunction with long-acting insulin AND • Prescriber acknowledges that Afrezza is not recommended in patients who smoke or have recently stopped smoking.
Short-Acting		
No PA Required HUMULIN R U-100 (insulin regular) vial (OTC)	PA Required NOVOLIN R U-100 (insulin regular) vial (OTC)	Non-preferred products may be approved following trial and failure of treatment with one preferred product (failure is defined as allergy or intolerable side effects).

NOVOLIN R U-100 (insulin regular) FlexPen (OTC)		
Intermediate-Acting		
No PA Required	PA Required	Non-preferred products may be approved following trial and failure of treatment with one preferred product (failure is defined as allergy or intolerable side effects).
HUMULIN N U-100 (insulin NPH), KwikPen (OTC), vial (OTC)	NOVOLIN N U-100 (insulin NPH) vial (OTC)	
NOVOLIN N U-100 (insulin NPH) FlexPen (OTC)		
Long-Acting		
Preferred	Non-Preferred	*Preferred Tresiba pen and vial formulations may be approved for members who have trialed and failed‡ Lantus. Non-preferred products may be approved if the member has tried and failed‡ treatment with Lantus AND a preferred insulin degludec product. ‡Failure is defined as lack of efficacy, allergy, or intolerable side effects.
LANTUS ^{BNR} (insulin glargine) Solostar, vial	BASAGLAR (insulin glargine) Kwikpen, Tempo pen	
TRESIBA ^{BNR} (insulin degludec)* FlexTouch, vial	Insulin degludec FlexTouch, vial Insulin glargine solostar, vial Insulin glargine MAX solostar Insulin glargine-yfgn pen, vial LEVEMIR (insulin detemir) FlexTouch, vial REZVOGLAR (insulin glargine-aglr) Kwikpen SEMGLEE (insulin glargine-yfgn) pen, vial TOUJEO (insulin glargine) Solostar TOUJEO MAX (insulin glargine) Solostar	
Concentrated		
No PA Required	PA Required	Non-preferred products may be approved following trial and failure of treatment with one preferred product (failure is defined as allergy or intolerable side effects).
HUMULIN R U-500 (insulin regular) concentrated vial, Kwikpen		
Mixtures		
No PA Required	PA Required	
HUMALOG MIX 50/50 Kwikpen		

HUMALOG MIX 75/25 vial HUMULIN 70/30 (OTC) Kwikpen, vial Insulin aspart protamine/insulin aspart 70/30 FlexPen, vial (generic Novolog Mix) Insulin lispro protamine/insulin lispro 75/25 Kwikpen (generic Humalog Mix) NOVOLOG MIX 70/30 FlexPen, vial	HUMALOG MIX 75/25 Kwikpen NOVOLIN 70/30 FlexPen, vial (OTC)	Non-preferred products may be approved if the member has failed treatment with two of the preferred products (failure is defined as: allergy or intolerable side effects).
Therapeutic Drug Class: DIABETES MANAGEMENT CLASSES, NON- INSULINS – 5/9/2025		
Amylin		
	PA Required SYMLIN (pramlintide) pen	SYMLIN (pramlintide) may be approved following trial and failure of metformin AND trial and failure of a DPP-4 inhibitor or GLP-1 analogue. Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen) following 3-month trial, allergy, intolerable side effects, or a significant drug-drug interaction. Prior authorization may be approved for Symlin (pramlintide) products for members with a diagnosis of Type 1 diabetes without requiring trial and failure of other products. Maximum Dose: Prior authorization will be required for doses exceeding FDA-approved dosing listed in product package labeling.
Biguanides		
No PA Required Metformin IR tablets Metformin ER 500mg, 750mg tablets (generic Glucophage XR)	PA Required GLUMETZA ER (metformin) tablet Metformin 625 mg tablets Metformin ER (generic Fortamet, Glumetza, Bayshore Pharma) Metformin solution (generic Riomet) RIOMET (metformin) solution RIOMET ER (metformin) suspension	Non-preferred products may be approved for members who have failed treatment with two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. Liquid metformin may be approved for members that are unable to use a solid oral dosage form.
Dipeptidyl Peptidase 4 Enzyme inhibitors (DPP-4 Inhibitors)		
Preferred	Non-Preferred	

TRADJENTA (linagliptin) tablet	PA Required		Non-preferred DPP-4 inhibitors may be approved after a member has failed a 3-month trial of one preferred product. Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen), allergy, contraindication, intolerable side effects, or a significant drug-drug interaction.													
	Alogliptin tablet		<u>Continuation of therapy:</u> Members currently stabilized on Januvia (sitagliptin) may receive approval for continuation of therapy with that agent.													
	BRYNOVIN (Sitagliptin) tablet for suspension															
	JANUVIA (sitagliptin) tablet															
	NESINA (alogliptin) tablet		<u>Maximum Dose:</u> Prior authorization will be required for doses exceeding the FDA-approved maximum dosing listed in the following table:													
	ONGLYZA (saxagliptin) tablet															
	Saxagliptin tablet															
	Sitagliptin (generic Zituvio)															
	ZITUVIO (sitagliptin tablet)															
		<table><tr><th>DPP-4 Inhibitor</th><th>FDA-Approved Maximum Daily Dose</th></tr><tr><td>Alogliptin (generic Nesina)</td><td>25 mg/day</td></tr><tr><td>Januvia (sitagliptin)</td><td>100 mg/day</td></tr><tr><td>Nesina (alogliptin)</td><td>25 mg/day</td></tr><tr><td>Onglyza (saxagliptin)</td><td>5 mg/day</td></tr><tr><td>Tradjenta (linagliptin)</td><td>5 mg/day</td></tr><tr><td>Zituvio (sitagliptin)</td><td>100 mg/day</td></tr></table>	DPP-4 Inhibitor	FDA-Approved Maximum Daily Dose	Alogliptin (generic Nesina)	25 mg/day	Januvia (sitagliptin)	100 mg/day	Nesina (alogliptin)	25 mg/day	Onglyza (saxagliptin)	5 mg/day	Tradjenta (linagliptin)	5 mg/day	Zituvio (sitagliptin)	100 mg/day
DPP-4 Inhibitor	FDA-Approved Maximum Daily Dose															
Alogliptin (generic Nesina)	25 mg/day															
Januvia (sitagliptin)	100 mg/day															
Nesina (alogliptin)	25 mg/day															
Onglyza (saxagliptin)	5 mg/day															
Tradjenta (linagliptin)	5 mg/day															
Zituvio (sitagliptin)	100 mg/day															
DPP-4 Inhibitors – Combination with Metformin																
Preferred	Non-Preferred PA Required	Non-preferred combination products may be approved for members who have been stable on the two individual ingredients of the requested combination for three months AND have had adequate three-month trial and failure of a preferred combination agent. Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen), contraindication, allergy, intolerable side effects, or a significant drug-drug interaction.														
JENTADUETO (linagliptin/metformin) tablet	Alogliptin/metformin tablet															
JENTADUETO XR (linagliptin/metformin) tablet	JANUMET (sitagliptin/metformin) tablet															
	JANUMET XR (sitagliptin/metformin) tablet															
	KAZANO (alogliptin/metformin) tablet	<u>Continuation of therapy:</u> Members currently stabilized on Janumet (sitagliptin/metformin) or Janumet XR (sitagliptin ER/metformin ER) may receive approval for continuation of therapy with those agents.														

	KOMBIGLYZE XR (saxagliptin/metformin) Saxagliptin/metformin tablet Sitagliptin/metformin (generic Zituvimet)	Maximum Dose: Prior authorization will be required for doses exceeding the FDA-approved maximum dosing listed in the following table: <table><tr><th>DPP-4 Inhibitor Combination</th><th>FDA Approved Maximum Daily Dose</th></tr><tr><td>Alogliptin/metformin tablet</td><td>25 mg alogliptin/2,000 mg metformin</td></tr><tr><td>Janumet and Janumet XR (sitagliptin/metformin)</td><td>100 mg sitagliptin/ 2,000 mg of metformin</td></tr><tr><td>Jentadueto and Jentadueto XR (linagliptin/metformin)</td><td>5 mg linagliptin/ 2,000 mg metformin</td></tr><tr><td>Kazano (alogliptin/metformin)</td><td>25 mg alogliptin/ 2,000 mg metformin</td></tr><tr><td>Kombiglyze XR (saxagliptin ER/metformin ER) tablet</td><td>5 mg saxagliptin/ 2,000 mg metformin</td></tr></table>	DPP-4 Inhibitor Combination	FDA Approved Maximum Daily Dose	Alogliptin/metformin tablet	25 mg alogliptin/2,000 mg metformin	Janumet and Janumet XR (sitagliptin/metformin)	100 mg sitagliptin/ 2,000 mg of metformin	Jentadueto and Jentadueto XR (linagliptin/metformin)	5 mg linagliptin/ 2,000 mg metformin	Kazano (alogliptin/metformin)	25 mg alogliptin/ 2,000 mg metformin	Kombiglyze XR (saxagliptin ER/metformin ER) tablet	5 mg saxagliptin/ 2,000 mg metformin
DPP-4 Inhibitor Combination	FDA Approved Maximum Daily Dose													
Alogliptin/metformin tablet	25 mg alogliptin/2,000 mg metformin													
Janumet and Janumet XR (sitagliptin/metformin)	100 mg sitagliptin/ 2,000 mg of metformin													
Jentadueto and Jentadueto XR (linagliptin/metformin)	5 mg linagliptin/ 2,000 mg metformin													
Kazano (alogliptin/metformin)	25 mg alogliptin/ 2,000 mg metformin													
Kombiglyze XR (saxagliptin ER/metformin ER) tablet	5 mg saxagliptin/ 2,000 mg metformin													
Glucagon-like Peptide-1 Receptor Agonists (GLP-1 Analogues)														
Preferred *Must meet eligibility criteria *BYDUREON BCISE (exenatide ER) autoinjector *BYETTA^{BNR} (exenatide) pen *Liraglutide pen (<i>Teva</i>) *OZEMPIC (semaglutide) pen *TRULICITY (dulaglutide) pen *VICTOZA (liraglutide) pen **WEGOVY (semaglutide) pen	Non-Preferred PA Required Exenatide pen Liraglutide pen (<i>all other manufacturers</i>) MOUNJARO (tirzepatide) pen RYBELSUS (semaglutide) oral tablet ZEPBOUND (tirzepatide)	*Preferred products may be approved for members with a diagnosis of type 2 diabetes. **WEGOVY (semaglutide) may be approved if meeting the following criteria: - Member is 18 years of age or older AND - Member has established cardiovascular disease (history of myocardial infarction, stroke, or symptomatic peripheral arterial disease) and either obesity or overweight (defined as a BMI ≥ 25 kg/m²) AND - Member does not have a diagnosis of Type 1 or Type 2 diabetes AND - Wegovy (semaglutide) is being prescribed to decrease the risk of adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND - Member has been counseled regarding implementation of lifestyle interventions (diet modification and exercise) to promote weight loss. ZEPBOUND (tirzepatide) may be approved if meeting the following criteria: - Member is 18 years of age or older AND - Member has a documented diagnosis of moderate to severe obstructive sleep apnea (OSA) AND - Member has a BMI ≥ 30 kg/m² indicating obesity documented in medical chart notes AND - A polysomnogram has been performed at baseline with a documented result of Apnea-Hypopnea Index (AHI) ≥ 15 events/hour AND - Member is not pregnant or planning to become pregnant AND - Member has been counseled regarding the risk of medullary thyroid cancer (MTC) with the use of Zepbound (tirzepatide) and does not have a personal or family history of MTC or Multiple												

		Endocrine Neoplasia syndrome type 2 (MEN 2) AND • The requested medication is being prescribed by or in consultation with a neurologist, pulmonologist, otolaryngologist, or other sleep medicine specialist AND • Member has been counseled regarding and is engaged in implementation of lifestyle interventions (diet modification and exercise) to promote weight loss AND • Member has failed a 6-month trial of continuous positive airway pressure (CPAP) or has a contraindication to the use of PAP therapy. Requests for GLP-1 analogues that are FDA-indicated for the treatment of metabolic dysfunction-associated steatohepatitis (MASH) may be approved if meeting the following: • A diagnosis of MASH with moderate to advanced fibrosis has been confirmed through liver biopsy AND • Member meets the FDA-labeled minimum age requirement for the prescribed product AND • Member does not have cirrhosis or significant liver disease other than MASH AND • The requested medication is being prescribed for use for the FDA-labeled indication and as outlined in product package labeling AND • Medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider AND • Requests for non-preferred agents will be subject to meeting non-preferred criteria listed below. All other non-preferred products may be approved for members with an FDA-labeled diagnosis (excluding labeled use solely for weight loss) following a trial and failure‡ of three preferred agents that are FDA-labeled for use for the prescribed indication <u>Continuation of therapy:</u> Members that are currently stabilized on a non-preferred product may receive approval for continuation of therapy with that product. <u>Maximum Dose:</u> Prior authorization is required for all products exceeding maximum dose listed in product package labeling. <table><tr><th colspan="2">Table 1: GLP-1 Analogue Maximum Dose</th></tr><tr><td>Bydureon Bcise (exenatide ER)</td><td>2 mg weekly</td></tr><tr><td>Byetta (exenatide)</td><td>20 mcg daily</td></tr><tr><td>Mounjaro (tirzepatide)</td><td>15 mg weekly</td></tr><tr><td>Ozempic (semaglutide)</td><td>2 mg weekly</td></tr><tr><td>Rybelsus (semaglutide)</td><td>14 mg daily</td></tr><tr><td>Trulicity (dulaglutide)</td><td>4.5 mg weekly</td></tr><tr><td>Victoza (liraglutide)</td><td>1.8 mg daily</td></tr><tr><td>Wegovy (semaglutide)</td><td>2.4 mg weekly</td></tr></table> ‡Failure is defined as lack of efficacy with a 3-month trial (such as not meeting hemoglobin A1C goal despite adherence to regimen), allergy, intolerable side effects, limited dexterity resulting in the inability	Table 1: GLP-1 Analogue Maximum Dose		Bydureon Bcise (exenatide ER)	2 mg weekly	Byetta (exenatide)	20 mcg daily	Mounjaro (tirzepatide)	15 mg weekly	Ozempic (semaglutide)	2 mg weekly	Rybelsus (semaglutide)	14 mg daily	Trulicity (dulaglutide)	4.5 mg weekly	Victoza (liraglutide)	1.8 mg daily	Wegovy (semaglutide)	2.4 mg weekly
Table 1: GLP-1 Analogue Maximum Dose																				
Bydureon Bcise (exenatide ER)	2 mg weekly																			
Byetta (exenatide)	20 mcg daily																			
Mounjaro (tirzepatide)	15 mg weekly																			
Ozempic (semaglutide)	2 mg weekly																			
Rybelsus (semaglutide)	14 mg daily																			
Trulicity (dulaglutide)	4.5 mg weekly																			
Victoza (liraglutide)	1.8 mg daily																			
Wegovy (semaglutide)	2.4 mg weekly																			

		to administer doses of a preferred product, or a significant drug-drug interaction. <i>Note: Prior Authorization for GLP-1 analogues prescribed solely for weight loss will not be approved.</i>
Other Hypoglycemic Combinations		
	PA Required Alogliptin/pioglitazone tablet Glipizide/metformin tablet Glyburide/metformin tablet GLYXAMBI (empagliflozin/linagliptin) tablet OSENI (alogliptin/pioglitazone) tablet Pioglitazone/glimepiride tablet QTERN (dapagliflozin/saxagliptin) tablet SOLIQUA (insulin glargine/lixisenatide) pen STEGLUJAN (ertugliflozin/sitagliptin) tablet TRIJARDY XR tablet(empagliflozin/linagliptin/metformin) XULTOPHY (insulin degludec/liraglutide) pen	Non-preferred products may be approved for members who have been stable on each of the individual ingredients in the requested combination for 3 months (including cases where the ingredients are taken as two separate 3-month trials or when taken in combination for at least 3 months). SOLIQUA (insulin glargine/lixisenatide) may be approved if member has had a trial and failure with one preferred GLP-1 AND one preferred insulin glargine product (Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen), allergy, intolerable side effects, or significant drug-drug interaction.)
Meglitinides		
Repaglinide tablet	PA Required Nateglinide tablet	Non-preferred products may be approved for members who have failed treatment with one preferred product. Failure is defined as: lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen), allergy, intolerable side effects, or significant drug-drug interaction.
Meglitinides Combination with Metformin		
	PA Required Repaglinide/metformin	Non-preferred products may be approved for members who have been stable on the two individual ingredients of the requested combination for 3 months.
Sodium-Glucose Cotransporter Inhibitors (SGLT inhibitors)		
No PA Required FARXIGA ^{BNR} (dapagliflozin) tablet	PA Required Dapagliflozin tablet	Non-preferred products may receive approval following trial and failure with one preferred product. Failure is defined as lack of efficacy with 3-month trial (such as not meeting hemoglobin A1C goal despite adherence to regimen), contraindication, allergy, intolerable side effects, or a significant drug-drug interaction.

	INPEFA (sotagliflozin) tablet INVOKANA (canagliflozin) tablet JARDIANCE (empagliflozin) tablet STEGLATRO (ertugliflozin) tablet	<u>Maximum Dose:</u> Prior authorization is required for all products exceeding maximum dose listed in product package labeling.
SGLT Inhibitor Combinations with Metformin		
No PA Required XIGDUO XR ^{BNR} (dapagliflozin/metformin) tablet	PA Required Dapagliflozin/Metformin XR tablet INVOKAMET (canagliflozin/metformin) tablet INVOKAMET XR (canagliflozin/metformin) tablet SEGLUROMET (ertugliflozin/metformin) tablet SYNJARDY (empagliflozin/metformin) tablet SYNJARDY XR (empagliflozin/metformin) tablet	Non-preferred products may be approved for members who have been stable on the two individual ingredients of the requested combination for 3 months. INVOKAMET, INVOKAMET XR, SEGLUROMET, SYNJARDY, SYNJARDY XR and XIGDUO XR are contraindicated in patients with an eGFR less than 30 mL/min/1.73 m ² or on dialysis.
Thiazolidinediones (TZDs)		
No PA Required Pioglitazone tablet	PA Required ACTOS (pioglitazone) tablet	Non-preferred agents may be approved following trial and failure of one preferred product. Failure is defined as lack of efficacy (such as not meeting hemoglobin A1C goal despite adherence to regimen) with a 3-month trial, allergy, intolerable side effects, or a significant drug-drug interaction.
Thiazolidinediones Combination with Metformin		
Pioglitazone/metformin tablet	PA Required ACTOPLUS MET (pioglitazone/metformin) TABLET	Non-preferred products may be approved for members who have been stable on the two individual ingredients of the requested combination for 3 months.
Therapeutic Drug Class: ESTROGEN AGENTS -Effective 10/1/2025		
No PA Required	PA Required	Non-preferred parenteral estrogen agents may be approved with trial and failure of one preferred parenteral agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.
Parenteral		
DELESTROGEN ^{BNR} (estradiol valerate) vial	Estradiol valerate 10mg/mL vial	

DEPO-ESTRODIOL (estradiol cypionate) vial Estradiol valerate 40mg/mL vial, 20mg/mL vial		Non-preferred oral estrogen agents may be approved with trial and failure of one preferred oral agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. Non-preferred transdermal estrogen agents may be approved with trial and failure of two preferred transdermal agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.																				
Oral/Transdermal																						
Estradiol oral tablet Estradiol (generic Climara) weekly patch MINIVELLE ^{BNR} (estradiol) patch VIVELLE-DOT ^{BNR} (estradiol) patch	CLIMARA (estradiol) patch DOTTI (estradiol) patch ESTRACE (estradiol) oral tablet Estradiol bi-weekly patch LYLLANA (estradiol) patch MENOSTAR (estradiol) patch	<table><tr><th colspan="2">Table 1: Transdermal Estrogen FDA-Labeled Dosing</th></tr><tr><td>ALORA (estradiol) patch</td><td>2/week</td></tr><tr><td>CLIMARA (estradiol) patch</td><td>1/week</td></tr><tr><td>DOTTI (estradiol) patch</td><td>2/week</td></tr><tr><td>Estradiol patch (once weekly)</td><td>1/week</td></tr><tr><td>Estradiol patch (twice weekly)</td><td>2/week</td></tr><tr><td>LYLLANA (estradiol) patch</td><td>2/week</td></tr><tr><td>MENOSTAR (estradiol) patch</td><td>1/week</td></tr><tr><td>MINIVELLE (estradiol) patch</td><td>2/week</td></tr><tr><td>VIVELLE-DOT (estradiol) patch</td><td>2/week</td></tr></table> <i>Note: Estrogen agents are a covered benefit for gender affirming hormone therapy and treating clinicians and mental health providers should be knowledgeable about the diagnostic criteria for gender-affirming hormone treatment and have sufficient training and experience in assessing related mental health conditions.</i>	Table 1: Transdermal Estrogen FDA-Labeled Dosing		ALORA (estradiol) patch	2/week	CLIMARA (estradiol) patch	1/week	DOTTI (estradiol) patch	2/week	Estradiol patch (once weekly)	1/week	Estradiol patch (twice weekly)	2/week	LYLLANA (estradiol) patch	2/week	MENOSTAR (estradiol) patch	1/week	MINIVELLE (estradiol) patch	2/week	VIVELLE-DOT (estradiol) patch	2/week
Table 1: Transdermal Estrogen FDA-Labeled Dosing																						
ALORA (estradiol) patch	2/week																					
CLIMARA (estradiol) patch	1/week																					
DOTTI (estradiol) patch	2/week																					
Estradiol patch (once weekly)	1/week																					
Estradiol patch (twice weekly)	2/week																					
LYLLANA (estradiol) patch	2/week																					
MENOSTAR (estradiol) patch	1/week																					
MINIVELLE (estradiol) patch	2/week																					
VIVELLE-DOT (estradiol) patch	2/week																					
Therapeutic Drug Class: GLUCAGON, SELF-ADMINISTERED – <i>Effective 11/8/2024</i>																						
Preferred No PA Required	Non-Preferred PA Required																					
BAQSIMI (glucagon) nasal spray Glucagon Emergency Kit (<i>Eli Lilly, Fresenius, Amphastar</i>) ZEGALOGUE (dasiglucagon) autoinjector	GVOKE (glucagon) Hypopen, Syringe, vial ZEGALOGUE (dasiglucagon) syringe	Non-preferred products may be approved if the member has failed treatment with two preferred products (failure is defined as allergy to ingredients in product, intolerable side effects, contraindication, or inability to administer dosage form). Quantity limit for all products: 2 doses per year unless used/ damaged/ lost																				
Therapeutic Drug Class: GROWTH HORMONES – <i>Effective 10/1/2025</i>																						

Preferred	Non-Preferred	All preferred products may be approved if the member has one of the qualifying diagnoses listed below (diagnosis may be verified through AutoPA) AND if prescription does not exceed limitations for maximum dosing (Table 1). *Second line preferred products (NGENLA, SKYTROFA) require trial and failure of Genotropin (somatropin) OR Norditropin (somatropin). Non-preferred Growth Hormone products may be approved if the following criteria are met: - Member failed treatment with one preferred growth hormone product (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions) AND - Member has a qualifying diagnosis that includes any of the following conditions: - Prader-Willi Syndrome (PWS) - Chronic renal insufficiency/failure requiring transplantation (defined as Creatinine Clearance < 30mL/min) - Turner's Syndrome - Hypopituitarism: as a result of pituitary disease, hypothalamic disease, surgery, radiation therapy or trauma verified by one of the following: - Has failed at least one GH stimulation test (peak GH level < 10 ng/mL) - Has at least one documented low IGF-1 level (below normal range for patient's age – refer to range on submitted lab document) - Has deficiencies in ≥ 3 pituitary axes (such as TSH, LH, FSH, ACTH, ADH) - Cachexia associated with AIDS - Noonan Syndrome - Short bowel syndrome - Neonatal symptomatic growth hormone deficiency (limited to 3-month PA approval) AND - Prescription does not exceed limitations for FDA-labeled maximum dosing for prescribed indication (Table 1) based on prescriber submission/verification of patient weight from most recent clinical documentation															
GENOTROPIN (somatropin) cartridge, Miniquick pen NGENLA (Somatrogen-ghla)* pen NORDITROPIN (somatropin) Flexpro pen SKYTROFA (lonapegsomatropin-tcgd)* cartridge	HUMATROPE (somatropin) cartridge NUTROPIN AQ (somatropin) Nuspin injector OMNITROPE (somatropin) cartridge, vial SAIZEN (somatropin) cartridge, vial SEROSTIM (somatropin) vial SOGROYA (somapacitan-beco) pen ZOMACTON (somatropin) vial	Table 1: Growth Hormone Product Maximum Dosing* <table> <tr> <th>Medication</th><th>Pediatric Maximum Dosing per week (age < 18 years)</th><th>Adult Maximum Dosing per week (age ≥ 18 years)</th></tr> <tr> <td>Genotropin</td><td>0.48 mg/kg/week</td><td>0.08 mg/kg/week</td></tr> <tr> <td>Humatrope</td><td>0.47 mg/kg/week</td><td>0.0875 mg/kg/week</td></tr> <tr> <td>Ngenla</td><td>0.66 mg/kg/week</td><td>Not Indicated</td></tr> <tr> <td>Norditropin Flexpro</td><td>0.47 mg/kg/week</td><td>0.112 mg/kg/week</td></tr> </table>	Medication	Pediatric Maximum Dosing per week (age < 18 years)	Adult Maximum Dosing per week (age ≥ 18 years)	Genotropin	0.48 mg/kg/week	0.08 mg/kg/week	Humatrope	0.47 mg/kg/week	0.0875 mg/kg/week	Ngenla	0.66 mg/kg/week	Not Indicated	Norditropin Flexpro	0.47 mg/kg/week	0.112 mg/kg/week
Medication	Pediatric Maximum Dosing per week (age < 18 years)	Adult Maximum Dosing per week (age ≥ 18 years)															
Genotropin	0.48 mg/kg/week	0.08 mg/kg/week															
Humatrope	0.47 mg/kg/week	0.0875 mg/kg/week															
Ngenla	0.66 mg/kg/week	Not Indicated															
Norditropin Flexpro	0.47 mg/kg/week	0.112 mg/kg/week															

		Nutropin AQ Nuspin	0.7 mg/kg/week	0.175 mg/kg/week for ≤35 years of age 0.0875 mg/kg/week for >35 years of age	
		Omnitrope	0.48 mg/kg/week	0.08 mg/kg/week	
		Saizen	0.18 mg/kg/week	0.07 mg/kg/week	
		Serostim	Not Indicated	42 mg/week for HIV wasting or cachexia (in combination with antiretroviral therapy)	
		Skytrofa	1.68 mg/kg/week	Not Indicated	
		Sogroya	Dose Individualized for each patient, based on growth response	8 mg/week	
		Zomacton	0.47 mg/kg/week	0.0875 mg/kg/week	
		Zorbtive	Not Indicated	56 mg/week for up to 4 weeks for short bowel syndrome only	
		*Based on FDA labeled indications and dosing			

VII. Gastrointestinal

Therapeutic Drug Class: **BILE SALTS** – *Effective 7/1/2025*

No PA Required	PA Required	
Ursodiol capsule	BYLVAY (odevixibat) capsule, pellet	Actigall (ursodiol) may be approved for members who meet the following criteria: - Member is ≥ 18 years of age AND - Member has tried and failed therapy with a 12-month trial of a preferred ursodiol product (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions). Chenodal (chenodiol) may be approved for members who meet the following criteria: - Member is > 18 years of age AND - Member has tried and failed therapy with a 12-month trial of a preferred ursodiol product (failure is defined as lack of efficacy, contraindication, allergy, intolerable side effects or significant drug-drug interactions). If chenodiol is
Ursodiol tablet	CHENODAL (chenodiol) tablet	
	CHOLBAM (cholic acid) capsule	
	LIVMARLI (maralixibat) solution, tablet	
	OICALIVA (obeticholic acid) tablet	

	RELTONE (ursodiol) capsule URSO (ursodiol) tablet URSO FORTE (ursodiol) tablet	being prescribed for treatment of cerebrotendinous xanthomatosis, no trial and failure of ursodiol is required. Cholbam (cholic acid) may be approved for members who meet the following criteria: • Bile acid synthesis disorders: ○ Member age must be greater than 3 weeks old AND ○ Member has a diagnosis for bile acid synthesis disorder due to single enzyme defect (Single Enzyme-Defect Disorders: Defective sterol nucleus synthesis, 3β-hydroxy-Δ-c27-steroid oxidoreductase deficiency, AKR1D1 deficiency, CYP7A1 deficiency, Defective side-chain synthesis, CYP27A1 deficiency (cerebrotendinous xanthomatosis), 2-methylacyl-CoA racemase deficiency (AMACR), 25-hydroxylation pathway (Smith–Lemli–Opitz). • Peroxisomal disorder including Zellweger spectrum disorders: ○ Member age must be greater than 3 weeks old AND ○ Member has diagnosis of peroxisomal disorders (PDs) including Zellweger spectrum disorders AND ○ Member has manifestations of liver disease, steatorrhea or complications from decreased fat-soluble vitamin absorption. Ocaliva (obeticholic acid) may be approved for members meeting the following criteria: • Member is ≥ 18 years of age AND • Medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider AND • Member has the diagnosis of primary biliary cholangitis without cirrhosis OR a diagnosis of primary biliary cholangitis with compensated cirrhosis with no evidence of portal hypertension AND • Member has failed treatment with a preferred ursodiol product for at least 6 months due to an inadequate response, intolerable side effects, drug-drug interaction, or allergy to preferred ursodiol formulations. Reltone (ursodiol) may be approved for members meeting the following criteria: • Member is ≥ 18 years of age AND • The requested medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider AND • The requested medication is being prescribed for one of the following: ○ Treatment of radiolucent, noncalcified gallbladder stones < 20 mm in greatest diameter AND elective cholecystectomy would be undertaken except for the presence of increased surgical risk due to systemic disease, advanced age, idiosyncratic reaction to general anesthesia, or for those patients who refuse surgery OR ○ Prevention of gallstone formation in obese patients experiencing rapid weight loss AND
--	---	---

		• No compelling reasons for the member to undergo cholecystectomy exist, including unremitting acute cholecystitis, cholangitis, biliary obstruction, gallstone pancreatitis, or biliary-gastrointestinal fistula, AND • Member has trialed and failed treatment with a preferred ursodiol product for at least 6 months due to an inadequate response, intolerable side effects, drug-drug interaction, or allergy to inactive ingredients contained in the preferred ursodiol formulations. <u>Initial approval:</u> 1 year <u>Reauthorization:</u> May be reauthorized for 1 additional year with provider attestation that partial or complete stone dissolution was observed after completion of the initial year of Reltone therapy. Maximum cumulative approval per member is 24 months. Urso (ursodiol) and Urso Forte (ursodiol) may be approved for members meeting the following criteria: • Member is ≥ 18 years of age AND • Medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider AND • Member has the diagnosis of Primary Biliary Cholangitis as evidenced by two of the following at the time of diagnosis: ○ Evidence of cholestasis with an alkaline phosphatase elevation of at least 1.5 times the upper limit of normal ○ Presence of antimitochondrial antibody with titer of 1:40 or higher ○ Histologic evidence of nonsuppurative destruction cholangitis and destruction of interlobular bile ducts AND • Member has failed treatment with a preferred ursodiol product for at least 6 months due to an inadequate response, intolerable side effects, drug-drug interaction, or allergy to inactive ingredients contained in the preferred ursodiol formulations. Requests for drug products that are FDA-indicated for the treatment of nonalcoholic steatohepatitis (NASH) may be approved if meeting the following: • A diagnosis of NASH has been confirmed through liver biopsy AND • Member meets the FDA-labeled minimum age requirement for the prescribed product AND • Member does not have significant liver disease other than NASH, AND • The requested medication is being prescribed for use for the FDA-labeled indication and as outlined in product package labeling AND • Medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider. Non-preferred products prescribed for FDA-labeled indications not identified above may receive approval for use as outlined in product package labeling.
--	--	---

Therapeutic Drug Class: ANTI-EMETICS, Oral – <i>Effective 7/1/2025</i>		
No PA Required DICLEGIS DR ^{BNR} tablet (doxylamine/pyridoxine) Meclizine (Rx) 12.5 mg, 25 mg tablet Metoclopramide solution, tablet Ondansetron ODT; 4mg, 8mg tablet Ondansetron oral suspension/ solution Prochlorperazine tablet Promethazine syrup, tablet	PA Required AKYNZEO (netupitant/palonosetron) capsule ANTIVERT (meclizine) 50 mg tablet ANZEMET (dolasetron) tablet Aprepitant capsule, tripack BONJESTA ER (doxylamine/pyridoxine) tablet Doxylamine/pyridoxine tablet (generic Diclegis) Dronabinol capsule EMEND (aprepitant) capsule, powder for suspension, dose/tri-pack Granisetron tablet MARINOL (dronabinol) capsule Ondansetron 16mg tablet REGLAN (metoclopramide) tablet Trimethobenzamide capsule ZOFRAN (ondansetron) tablet	Emend (aprepitant) TriPack or Emend (aprepitant) powder kit may be approved following trial and failure of two preferred products AND Emend (aprepitant) capsule. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction. Doxylamine/pyridoxine tablet (generic) or Bonjesta (doxylamine/pyridoxine) may be approved for 9 months if meeting the following criteria: • Member has nausea and vomiting associated with pregnancy AND • Member has trialed and failed DICLEGIS DR tablet AND one of the following (failure is defined as lack of efficacy with a 7-day trial, allergy, intolerable side effects, or significant drug-drug interaction): ○ Antihistamine (such as diphenhydramine, dimenhydrinate, meclizine) OR ○ Dopamine antagonist (such as metoclopramide, prochlorperazine, promethazine) OR ○ Serotonin antagonist (ondansetron, granisetron) All other non-preferred products may be approved for members who have trialed and failed treatment with two preferred products. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction. Dronabinol prior authorization may be approved for members meeting above non-preferred criteria OR via AutoPA for members with documented HIV diagnosis. Promethazine product formulations require prior authorization for members < 2 years of age due to risk of fatal respiratory depression.
Therapeutic Drug Class: ANTI-EMETICS, Non-Oral – <i>Effective 7/1/2025</i>		
No PA Required Prochlorperazine 25 mg suppository	PA Required PROMETHEGAN 50 mg (Promethazine) suppository SANCUSO (granisetron) patch	Non-preferred products may be approved for members who have trialed and failed treatment with two preferred products. Failure is defined as lack of efficacy with 14-day trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.

Promethazine 12.5 mg, 25 mg suppository	TRANSDERM-SCOP (scopolamine) patch	
Scopolamine patch		
Therapeutic Drug Class: GI MOTILITY, CHRONIC – <i>Effective 7/1/2025</i>		
PA Required for all agents in this class		All agents will only be approved for FDA labeled indications and up to FDA approved maximum doses listed below.
Preferred	Non-Preferred	Preferred agents may be approved if the member meets the following criteria: - Has diagnosis of Irritable Bowel Syndrome – Constipation (IBS-C), Chronic Idiopathic Constipation (CIC), Functional Constipation (FC), or Opioid Induced Constipation (OIC) in patients with opioids prescribed for noncancer pain AND - Member does not have a diagnosis of GI obstruction AND - For indication of OIC, member opioid use must exceed 4 weeks of treatment AND - For indications of CIC, OIC, IBS-C; member must have documentation of adequate trial of two or more over-the-counter motility agents (polyethylene glycol, docusate or bisacodyl, for example). OR If the member cannot take oral medications, then the member must fail a 7-day trial with a nonphosphate enema (docusate or bisacodyl enema). Failure is defined as a lack of efficacy for a 7-day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction AND - For indication of IBS-D, must have documentation of adequate trial and failure with loperamide and trial and failure with dicyclomine or hyoscyamine. Failure is defined as a lack of efficacy for a 7-day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction. Non-preferred agents may be approved if the member meets the following criteria: - Member meets all listed criteria for preferred agents AND - Member has trialed and failed two preferred agents OR if the indication is OIC caused by methadone, then a non-preferred agent may be approved after an adequate trial of MOVANTIK (naloxegol). Failure is defined as a lack of efficacy for a 7-day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction AND - If prescribed Viberzi (eluxadoline) or Lotronex (alosetron), member meets the additional criteria for those agents listed below. VIBERZI (eluxadoline) may be approved for members who meet the following additional criteria: - Diagnosis of Irritable Bowel Syndrome – Diarrhea (IBS-D) AND - Member has a gallbladder AND - Member does not have severe hepatic impairment (Child-Pugh C), history of
LINZESS (linaclotide) capsule	Alosetron tablet	
Lubiprostone capsule	AMITIZA (lubiprostone) capsule	
MOVANTIK (naloxegol) tablet	IBSRELA tablet	
	LOTROXEX (alosetron) tablet	
	MOTTEGRITY (prucalopride) tablet	
	Prucalopride tablet	
	RELISTOR (methylnaltrexone) syringe, tablet, vial	
	SYMPROIC (naldemedine) tablet	
	TRULANCE (plecanatide) tablet	
	VIBERZI (eluxadoline) tablet	

		severe constipation, known mechanical gastrointestinal obstruction, biliary duct obstruction, history of pancreatitis or structural disease of the pancreas AND - Member does not drink more than 3 alcoholic drinks per day LOTIRONEX (alosetron) and generic alosetron may be approved for members who meet the following additional criteria: - Member is a female with Irritable Bowel Syndrome – Diarrhea (IBS-D) with symptoms lasting 6 months or longer AND - Member does not have severe hepatic impairment (Child-Pugh C), history of severe constipation or ischemic colitis, hypercoagulable state, Crohn’s disease or ulcerative colitis, or known mechanical gastrointestinal obstruction.
--	--	--

Medication	FDA approved indication	FDA Max Dose
Amitiza (lubiprostone)	IBS-C (females only), CIC, OIC (not caused by methadone)	48mcg/day
Linzess (linaclotide)	IBS-C, CIC (≥ 18 years)	290mcg/day
Movantik (naloxegol)	OIC, FC (6 to 17 years)	25mg/day (OIC), 72mcg/day (FC)
Viberzi (eluxadoline)	IBS-D	200mg/day
Relistor subcutaneous injection (methylnaltrexone)	OIC	12mg/day
Relistor oral (methylnaltrexone)	OIC	450mg/day
Lotronex (alosetron)	IBS-D (women only)	2mg/day (women only)
Symproic (Naldemedine)	OIC	0.2mg/day
Trulance (plecanatide)	CIC, IBS-C	3mg/day
Motegrity (prucalopride)	CIC	2mg/day

CIC – chronic idiopathic constipation, FC – functional constipation, OIC – opioid induced constipation, IBS – irritable bowel syndrome, D – diarrhea predominant, C – constipation predominant

Therapeutic Drug Class: **H. PYLORI TREATMENTS** – Effective 7/1/2025

No PA Required	PA Required	
PYLERA ^{BNR} capsule (bismuth subcitrate/metronidazole tetracycline)	Amoxicillin/lansoprazole/clarithromycin pack Bismuth subcitrate/metronidazole tetracycline capsule OMECLAMOX-PAK (amoxicillin/omeprazole/clarithromycin) TALICIA (omeprazole/amoxicillin/ rifabutin) tablet	Non-preferred <i>H. pylori</i> treatments should be used as individual product ingredients unless one of the individual products is not commercially available, then a PA for the combination product may be given.

	VOQUEZNA DUAL (vonoprazan/amoxicillin) dose pack	
	VOQUEZNA TRIPLE (vonoprazan/amoxicillin/clarithromycin dose pack	
Therapeutic Drug Class: HEMORRHOIDAL, ANORECTAL, AND RELATED TOPICAL ANESTHETIC AGENTS – <i>Effective 7/1/2025</i>		
Hydrocortisone single agent		Non-preferred products may be approved following trial and failure of therapy with 3 preferred products (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).
No PA Required	PA Required	
ANUSOL-HC (hydrocortisone) 2.5% cream with applicator	CORTENEMA (hydrocortisone) enema	
CORTIFOAM (hydrocortisone) 10% aerosol	PROCORT cream	
Hydrocortisone 1% cream with applicator		
Hydrocortisone 2.5% cream with applicator		
Hydrocortisone enema		
Lidocaine single agent		
No PA Required	PA Required	
Lidocaine 3% cream, 5% ointment		
Other and Combinations		
No PA Required	PA Required	RECTIV (nitroglycerin) ointment may be approved if meeting the following:
Lidocaine-Hydrocortisone 3-0.5% cream with applicator	ANALPRAM HC (Hydrocortisone-Pramoxine) 1%-1% cream, 2.5%-1% cream	- Member has a diagnosis of anal fissure AND - Prescriber attests that member has trialed and maximized use of appropriate supportive therapies including sitz bath, fiber, topical analgesics (such as lidocaine), and stool softeners/laxatives.
Lidocaine-Prilocaine Cream (<i>all other manufacturers</i>)	EPIFOAM (Hydrocortisone-Pramoxine) 1%-1% foam	
PROCTOFOAM-HC (hydrocortisone-pramoxine) 1%-1% foam	Hydrocortisone-Pramoxine 1%-1%, 2.5%-1% cream	
	Lidocaine-Hydrocortisone in Coleus 2%-2% cream kit	
	Lidocaine-Hydrocortisone 2.8%-0.55% gel	

	Lidocaine-Hydrocortisone 3%-0.5% cream w/o applicator, cream kit Lidocaine-Hydrocortisone 3%-1% cream kit Lidocaine-Hydrocortisone 3%-2.5% gel kit Lidocaine-Prilocaine Cream (<i>Fougera only</i>) PLIAGLIS (lidocaine-tetracaine) 7%-7% cream PROCORT (Hydrocortisone-Pramoxine) 1.85%-1.15% cream RECTIV (nitroglycerin) 0.4% ointment	
Therapeutic Drug Class: PANCREATIC ENZYMES – <i>Effective 7/1/2025</i>		
No PA Required CREON (pancrelipase) capsule VIOKACE (pancrelipase) tablet ZENPEP (pancrelipase) capsule	PA Required PERTZYE (pancrelipase) capsule	Non-preferred products may be approved for members who have failed an adequate trial (4 weeks) with at least two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction.
Therapeutic Drug Class: PROTON PUMP INHIBITORS – <i>Effective 7/1/2025</i>		
No PA Required Esomeprazole DR packet for oral suspension, capsule (RX) Lansoprazole DR capsules (RX) Lansoprazole ODT (RX) <i>(for members under 2 years)</i> Omeprazole DR capsule (RX) Pantoprazole tablet PROTONIX (pantoprazole DR) packet for oral suspension ^{BNR}	PA Required ACIPHEX (rabeprazole) tablet, sprinkle capsule DEXILANT (dexlansoprazole) capsule Dexlansoprazole capsule Esomeprazole DR 49.3 capsule (RX), (OTC) capsule KONVOMEF (Omeprazole/Na bicarbonate) suspension Lansoprazole DR capsule OTC Lansoprazole ODT (OTC) NEXIUM (esomeprazole) capsule (RX), oral suspension packet, 24HR (OTC)	For members treating GERD symptoms that are controlled on PPI therapy, it is recommended that the dose of the PPI be re-evaluated or step-down with an H2 blocker (such as famotidine) be trialed in order to reduce long-term PPI use. Prior authorization for non-preferred proton pump inhibitors may be approved if all of the following criteria are met: • Member has a qualifying diagnosis (below) AND • Member has trialed and failed therapy with three preferred agents within the last 24 months. (Failure is defined as: lack of efficacy following 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction) AND • Member has been diagnosed using one of the following diagnostic methods: ○ Diagnosis made by GI specialist ○ Endoscopy ○ X-ray ○ Biopsy ○ Blood test ○ Breath Test Qualifying Diagnoses:

	Omeprazole/Na bicarbonate capsule, packet for oral suspension Omeprazole DR tablet (OTC), ODT (OTC) Pantoprazole packet for oral suspension PREVACID (lansoprazole) capsule, Solutab, suspension PRILOSEC (omeprazole) suspension PROTONIX (pantoprazole DR) tablet Rabeprazole tablet VOQUEZNA (vonoprazan) tablet ZEGERID (omeprazole/Na bicarbonate) capsule, packet for oral suspension	Barrett's esophagus, duodenal ulcer, erosive esophagitis, gastric ulcer, GERD, GI Bleed, H. pylori infection, hypersecretory conditions (Zollinger-Ellison), NSAID-induced ulcer, pediatric esophagitis, requiring mechanical ventilation, requiring a feeding tube Quantity Limits: All agents will be limited to once daily dosing except when used for the following diagnoses: Barrett's esophagus, GI Bleed, H. pylori infection, hypersecretory conditions (Zollinger-Ellison), or members who have spinal cord injury with associated acid reflux. Adult members with GERD on once daily, high-dose PPI therapy who continue to experience symptoms may receive initial prior authorization approval for a 4-week trial of twice daily, high-dose PPI therapy. Continuation of the twice daily dosing regimen for GERD beyond 4 weeks will require additional prior authorization approval verifying adequate member response to the dosing regimen and approval may be placed for one year. If a member with symptomatic GERD does not respond to twice daily, high-dose PPI therapy, this should be considered a treatment failure. Pediatric members (< 18 years of age) on once daily dosing of a PPI who continue to experience symptoms may receive one-year prior authorization approval for twice daily PPI therapy. Age Limits: Nexium 24H and Zegerid will not be approved for members less than 18 years of age. Prevacid Solutab may be approved for members < 2 years of age OR for members ≥ 2 years of age with a feeding tube. <u>Continuation of Care:</u> Members currently taking Dexilant (dexlansoprazole) capsules may continue to receive approval for that medication.
Therapeutic Drug Class: NON-BIOLOGIC ULCERATIVE COLITIS AGENTS- Oral – Effective 7/1/2025		
No PA Required	PA Required	
APRISO (mesalamine ER) capsule	AZULFIDINE (sulfasalazine) Entab, tablet	Prior authorization for non-preferred oral formulations will require trial and failure of two preferred oral products with different active ingredients AND one preferred rectal product. If inflammation is not within reach of topical therapy, trial of preferred rectal product is not required. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. Uceris (budesonide) tablet: Prior authorization may be approved following trial and failure of one preferred oral product AND one preferred rectal product. If inflammation is not within reach of topical therapy, trial of preferred rectal product is not required. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. Approval will be placed for 8 weeks. Further prior authorization may be approved if 7 days of steroid-free time has elapsed, and member continues to meet the above criteria.
Mesalamine DR tablet (generic Lialda) (<i>Takeda only</i>)	Balsalazide capsule	
Mesalamine ER capsule (generic Apriso) (<i>Teva only</i>)	Budesonide DR tablet	
PENTASA ^{BNR} (mesalamine) capsule	COLAZAL (balsalazide) capsule	
Sulfasalazine IR and DR tablet	DELZICOL (mesalamine DR) capsule	
	DIPENTUM (olsalazine) capsule	
	LIALDA (mesalamine DR) tablet	

	Mesalamine DR tablet (generic Asacol HD, Lialda) Mesalamine DR/ER capsule (generic Delzicol and Pentasa) UCERIS (budesonide) tablet	
--	--	--

Therapeutic Drug Class: NON-BIOLOGIC ULCERATIVE COLITIS AGENTS- Rectal – Effective 7/1/2025

No PA Required	PA Required	
Mesalamine suppository Mesalamine 4gm/60 ml enema (generic SF ROWASA) SF ROWASA enema, kit (mesalamine)	Budesonide foam CANASA (mesalamine) suppository Mesalamine enema, kit ROWASA enema, kit (mesalamine) UCERIS (budesonide) foam	Prior authorization for non-preferred rectal formulations will require trial and failure of one preferred rectal formulation and one preferred oral formulation (Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction). Uceris (budesonide) foam: If the above criteria are met, Uceris (budesonide) foam prior authorization may be approved for 6 weeks. Further prior authorization may be approved if 7 days of steroid-free time has elapsed, and member continues to meet the above criteria.

VIII. Hematological

Therapeutic Drug Class: ANTICOAGULANTS- Oral – Effective 7/1/2025

No PA Required	PA Required	
Dabigatran capsule ELIQUIS (apixaban) tablet, tablet pack Warfarin tablet XARELTO (rivaroxaban)^{BNR} 10 mg, 15 mg, 20 mg tablet, dose pack	PRADAXA (dabigatran) capsule, pellet Rivaroxaban tablet Rivaroxaban oral suspension SAVAYSA (edoxaban) tablet XARELTO (rivaroxaban) 2.5 mg tablet XARELTO (rivaroxaban) oral suspension	SAVAYSA (edoxaban) may be approved if all the following criteria have been met: • The member has failed therapy with two preferred agents. (Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction) AND • Member is not on dialysis AND • Member does not have CrCl > 95 mL/min AND • The member has a diagnosis of deep vein thrombosis (DVT), pulmonary embolism (PE) OR • The member has a diagnosis of non-valvular atrial fibrillation AND • The member does not have a mechanical prosthetic heart valve XARELTO 2.5mg (rivaroxaban) may be approved for members meeting all of the following criteria: • Xarelto 2.5mg is being prescribed to reduce major CV events in members diagnosis of chronic coronary artery disease (CAD) or peripheral artery disease AND • Xarelto 2.5mg is being taken twice daily and in combination with aspirin 75-100mg daily AND

		- Member must not be receiving dual antiplatelet therapy, other non-aspirin antiplatelet therapy, or other oral anticoagulant AND - Member must not have had an ischemic, non-lacunar stroke within the past month AND - Member must not have had a hemorrhagic or lacunar stroke at any time XARELTO (rivaroxaban) oral suspension may be approved without prior authorization for members <18 years of age who require a rivaroxaban dose of less than 10 mg OR with prior authorization verifying the member is unable to use the solid oral dosage form. All other non-preferred oral agents require trial and failure of two preferred oral agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction. Continuation of Care: Members with current prior authorization approval on file for a non-preferred <u>oral</u> anticoagulant medication may continue to receive approval for that medication
Therapeutic Drug Class: ANTICOAGULANTS- Parenteral – Effective 7/1/2025		
No PA Required Enoxaparin syringe Enoxaparin vial	PA Required ARIXTRA (fondaparinux) syringe Fondaparinux syringe FRAGMIN (dalteparin) vial, syringe LOVENOX (enoxaparin) syringe, vial	Non-preferred parenteral anticoagulants may be approved if member has trial and failure of one preferred parenteral agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction ARIXTRA (fondaparinux) may be approved if the following criteria have been met: - Member is 18 years of age or older AND - Member has a CrCl > 30 ml/min AND - Member weighs > 50 kg AND - Member has a documented history of heparin induced-thrombocytopenia OR - Member has a contraindication to enoxaparin Members currently stabilized on fondaparinux (Arixtra) or dalteparin (Fragmin) may receive prior authorization approval to continue receiving that medication.
Therapeutic Drug Class: ANTI-PLATELETS – Effective 4/8/2025		
No PA Required Aspirin/dipyridamole ER capsule BRILINTA (ticagrelor) tablet ^{BNR} Cilostazol tablet Clopidogrel tablet	PA Required EFFIENT (prasugrel) tablet PLAVIX (clopidogrel) tablet Ticagrelor tablet	Zontivity (vorapaxar) may be approved for patients with a diagnosis of myocardial infarction or peripheral artery disease without a history of stroke, transient ischemic attack, intracranial bleeding, or active pathological bleeding. Patients must also be taking aspirin and/or clopidogrel concomitantly. Non-preferred products without criteria will be reviewed on a case-by-case basis.

Dipyridamole tablet		
Pentoxifylline ER tablet		
Prasugrel tablet		
Therapeutic Drug Class: COLONY STIMULATING FACTORS – Effective 7/1/2025		
PA Required for all agents in this class*		*Prior authorization for preferred agents may be approved if meeting the following criteria: - Medication is being used for one of the following indications: - Patient with cancer receiving myelosuppressive chemotherapy –to reduce incidence of infection (febrile neutropenia) (Either the post nadir ANC is less than 10,000 cells/mm3 or the risk of neutropenia for the member is calculated to be greater than 20%) - Acute Myeloid Leukemia (AML) patients receiving chemotherapy - Bone Marrow Transplant (BMT) - Peripheral Blood Progenitor Cell Collection and Therapy - Hematopoietic Syndrome of Acute Radiation Syndrome - Severe Chronic Neutropenia (Evidence of neutropenia infection exists or ANC is below 750 cells/mm3) Prior authorization for non-preferred agents may be approved if meeting the following criteria: - Medication is being used for one of the following indications: - Patient with cancer receiving myelosuppressive chemotherapy –to reduce incidence of infection (febrile neutropenia) (Either the post nadir ANC is less than 10,000 cells/mm3 or the risk of neutropenia for the member is calculated to be greater than 20%) - Acute Myeloid Leukemia (AML) patients receiving chemotherapy - Bone Marrow Transplant (BMT) - Peripheral Blood Progenitor Cell Collection and Therapy - Hematopoietic Syndrome of Acute Radiation Syndrome - Severe Chronic Neutropenia (Evidence of neutropenia infection exists or ANC is below 750 cells/mm3) AND - Member has history of trial and failure of Neupogen AND one other preferred agent. Failure is defined as a lack of efficacy with a 3-month trial, allergy, intolerable side effects, significant drug-drug interactions, or contraindication to therapy. Trial and failure of Neupogen will not be required if meeting one of the following: - Member has limited access to caregiver or support system for assistance with medication administration OR - Member has inadequate access to healthcare facility or home care interventions.
Preferred	Non-Preferred	
FULPHILA (pegfilgrastim-jmdb) syringe	FYLNETRA (pegfilgrastim-jmdb) syringe	
NEUPOGEN (filgrastim) vial, syringe	GRANIX (tbo-filgrastim) syringe, vial	
	LEUKINE (sargramostim) vial	
	NEULASTA (pegfilgrastim) kit, syringe	
	NIVESTYM (filgrastim-aafi) syringe, vial	
	NYVEPRIA (pegfilgrastim-apgf) syringe	
	RELEUKO (filgrastim-ayow) syringe, vial	
	RYZNEUTA (efbemalenograstim alfa-vuxw) syringe	
	STIMUFEND (pegfilgrastim-fpgk) syringe	
	UDENYCA (pegfilgrastim-cbqv) autoinjector, On-Body, syringe	
	ZARXIO (filgrastim-sndz) syringe	
	ZIEXTENZO (pegfilgrastim-bmez) syringe	

Therapeutic Drug Class: ERYTHROPOIESIS STIMULATING AGENTS – <i>Effective 7/1/2025</i>		
PA Required for all agents in this class*		
Preferred	Non-Preferred	
EPOGEN (epoetin alfa) vial RETACRIT (epoetin alfa-epbx) (<i>Pfizer only</i>) vial	ARANESP (darbepoetin alfa) syringe, vial MIRCERA (methoxy peg-epoetin beta) syringe PROCRT (epoetin alfa) vial RETACRIT (epoetin alfa-epbx) (<i>Vifor only</i>) vial	*Prior Authorization is required for all products and may be approved if meeting the following: - Medication is being administered in the member's home or in a long-term care facility AND - Member meets <u>one</u> of the following: - A diagnosis of cancer, currently receiving chemotherapy, with chemotherapy-induced anemia, and hemoglobin[†] of 10g/dL or lower OR - A diagnosis of chronic renal failure, and hemoglobin[†] below 10g/dL OR - A diagnosis of hepatitis C, currently taking ribavirin and failed response to a reduction of ribavirin dose, and hemoglobin[†] less than 10g/dL (or less than 11g/dL if symptomatic) OR - A diagnosis of HIV, currently taking zidovudine, hemoglobin[†] less than 10g/dL, and serum erythropoietin level of 500 mU/mL or less OR - Member is undergoing elective, noncardiac, nonvascular surgery and medication is given to reduce receipt of allogenic red blood cell transfusions, hemoglobin[†] is greater than 10g/dL, but less than or equal to 13g/dL and high risk for perioperative blood loss. Member is not willing or unable to donate autologous blood pre-operatively AND - For any non-preferred product, member has trialed and failed treatment with one preferred product. Failure is defined as lack of efficacy with a 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction. [†]Hemoglobin results must be from the last 30 days.

IX. Immunological

Therapeutic Drug Class: IMMUNE GLOBULINS – <i>Effective 1/1/2025</i>		
PA Required for all agents in this class*		
Preferred	Non-Preferred	
CUVITRU 20% SQ liquid GAMMAGARD 10% IV/SQ liquid GAMUNEX-C 10% IV/SQ liquid HIZENTRA 20% SQ syringe, vial	ALYGLO 10% IV liquid BIVIGAM 10% IV liquid CUTAQUIG 16.5% SQ liquid FLEBOGAMMA DIF 5%, 10% IV liquid GAMMAGARD S/D vial	Preferred agents may be approved for members meeting at least one of the approved conditions listed below for prescribed doses not exceeding maximum (Table 1). Non-preferred agents may be approved for members meeting the following: - Member meets at least one of the approved conditions listed below AND - Member has history of trial and failure of two preferred agents (failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions) AND - Prescribed dose does not exceed listed maximum (Table 1) Approved Conditions for Immune Globulin Use: - Primary Humoral Immunodeficiency disorders including: - Common Variable Immunodeficiency (CVID) - Severe Combined Immunodeficiency (SCID)

PRIVIGEN 10% IV liquid <i>If immune globulin is being administered in a long-term care facility or in a member's home by a home healthcare provider, it should be billed as a pharmacy claim. All other claims must be submitted through the medical benefit.</i>	GAMMAKED 10% IV/SQ liquid GAMMAPLEX 5%, 10% IV liquid HYQVIA 10% SQ liquid OCTAGAM 5%, 10% IV liquid PANZYGA 10% IV liquid XEMBIFY 20% IV liquid	○ X-Linked Agammaglobulinemia ○ X-Linked with Hyperimmunoglobulin M (IgM) Immunodeficiency ○ Wiskott-Aldrich Syndrome ○ Members < 13 years of age with pediatric Human Immunodeficiency Virus (HIV) and CD-4 count > 200/mm3 • Neurological disorders including: ○ Guillain-Barré Syndrome ○ Relapsing-Remitting Multiple Sclerosis ○ Chronic Inflammatory Demyelinating Polyneuropathy ○ Myasthenia Gravis ○ Polymyositis and Dermatomyositis ○ Multifocal Motor Neuropathy • Kawasaki Syndrome • Chronic Lymphocytic Leukemia (CLL) • Autoimmune Neutropenia (AN) with absolute neutrophil count < 800 mm and history of recurrent bacterial infections • Autoimmune Hemolytic Anemia (AHA) • Liver or Intestinal Transplant • Immune Thrombocytopenia Purpura (ITP) including: ○ Requiring preoperative therapy for undergoing elective splenectomy with platelet count < 20,000/mcL ○ Members with active bleeding & platelet count <30,000/mcL ○ Pregnant members with platelet counts <10,000/mcL in the third trimester ○ Pregnant members with platelet count 10,000 to 30,000/mcL who are bleeding • Multisystem Inflammatory Syndrome in Children (MIS-C) <table><tr><th colspan="2">Table 1: FDA-Approved Maximum Immune Globulin Dosing</th></tr><tr><td>Asceniv – IV admin</td><td>800 mg/kg every 3 to 4 weeks</td></tr><tr><td>Bivigam – IV admin</td><td>800 mg/kg every 3 to 4 weeks</td></tr><tr><td>Cuvitru –subcutaneous admin</td><td>12 grams protein/site for up to four sites weekly (48grams/week)</td></tr><tr><td>Flebogamma DIF – IV admin</td><td>600 mg/kg every 3 weeks</td></tr><tr><td>Gammaplex 5% – IV admin</td><td>1 gram/kg for 2 consecutive days</td></tr><tr><td>Gammagard liquid subcutaneous or IV admin</td><td>2.4 grams/kg/month</td></tr><tr><td>Gammaked –subcutaneous or IV admin</td><td>600 mg/kg every 3 weeks</td></tr><tr><td>Gamunex-C –subcutaneous or IV admin</td><td>600 mg/kg every 3 weeks</td></tr><tr><td>Hizentra –subcutaneous admin</td><td>0.4 g/kg per week</td></tr><tr><td>Octagam – IV admin</td><td>2 grams/kg every 4 weeks</td></tr></table>	Table 1: FDA-Approved Maximum Immune Globulin Dosing		Asceniv – IV admin	800 mg/kg every 3 to 4 weeks	Bivigam – IV admin	800 mg/kg every 3 to 4 weeks	Cuvitru –subcutaneous admin	12 grams protein/site for up to four sites weekly (48grams/week)	Flebogamma DIF – IV admin	600 mg/kg every 3 weeks	Gammaplex 5% – IV admin	1 gram/kg for 2 consecutive days	Gammagard liquid subcutaneous or IV admin	2.4 grams/kg/month	Gammaked –subcutaneous or IV admin	600 mg/kg every 3 weeks	Gamunex-C –subcutaneous or IV admin	600 mg/kg every 3 weeks	Hizentra –subcutaneous admin	0.4 g/kg per week	Octagam – IV admin	2 grams/kg every 4 weeks
Table 1: FDA-Approved Maximum Immune Globulin Dosing																								
Asceniv – IV admin	800 mg/kg every 3 to 4 weeks																							
Bivigam – IV admin	800 mg/kg every 3 to 4 weeks																							
Cuvitru –subcutaneous admin	12 grams protein/site for up to four sites weekly (48grams/week)																							
Flebogamma DIF – IV admin	600 mg/kg every 3 weeks																							
Gammaplex 5% – IV admin	1 gram/kg for 2 consecutive days																							
Gammagard liquid subcutaneous or IV admin	2.4 grams/kg/month																							
Gammaked –subcutaneous or IV admin	600 mg/kg every 3 weeks																							
Gamunex-C –subcutaneous or IV admin	600 mg/kg every 3 weeks																							
Hizentra –subcutaneous admin	0.4 g/kg per week																							
Octagam – IV admin	2 grams/kg every 4 weeks																							

		<table><tr><td>Panzyga – IV admin</td><td>2 g/kg every 3 weeks</td></tr><tr><td>Privigen – IV admin</td><td>2 g/kg over 2 to 5 consecutive days</td></tr></table> Members currently receiving a preferred or non-preferred immunoglobulin product may receive approval to continue therapy with that product at prescribed doses not exceeding maximum (Table 1).	Panzyga – IV admin	2 g/kg every 3 weeks	Privigen – IV admin	2 g/kg over 2 to 5 consecutive days
Panzyga – IV admin	2 g/kg every 3 weeks					
Privigen – IV admin	2 g/kg over 2 to 5 consecutive days					
Therapeutic Drug Class: NEWER GENERATION ANTIHISTAMINES – <i>Effective 1/1/2025</i>						
No PA Required Cetirizine (OTC) syrup/solution (OTC/RX), tablet Desloratadine tablet (RX) Levocetirizine tablet (RX/OTC) Loratadine tablet (OTC), syrup/solution (OTC)	PA Required Cetirizine (OTC) chewable tablet, softgel, UD cups solution CLARINEX (desloratadine) tablet Desloratadine ODT (RX) Fexofenadine tablet (OTC), suspension (OTC) Levocetirizine solution (RX) Loratadine chewable (OTC), ODT (OTC)	Non-preferred single agent antihistamine products may be approved for members who have failed treatment with two preferred products in the last 6 months. For members with respiratory allergies, an additional trial of an intranasal corticosteroid will be required in the last 6 months. Failure is defined as lack of efficacy with a 14-day trial, allergy, intolerable side effects, or significant drug-drug interaction.				
Therapeutic Drug Class: ANTI HISTAMINE/DECONGESTANT COMBINATIONS – <i>Effective 1/1/2025</i>						
No PA Required Loratadine-D (OTC) tablet	PA Required Cetirizine-PSE (OTC) CLARINEX-D (desloratadine-D) Fexofenadine/PSE (OTC)	Non-preferred antihistamine/decongestant combinations may be approved for members who have failed treatment with the preferred product in the last 6 months. For members with respiratory allergies, an additional trial of an intranasal corticosteroid will be required in the last 6 months. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.				
Therapeutic Drug Class: INTRANASAL RHINITIS AGENTS – <i>Effective 1/1/2025</i>						
No PA Required Azelastine 137 mcg Budesonide (OTC) DYMISTA (azelastine/ fluticasone) ^{BNR} Fluticasone (RX)	PA Required Azelastine (Astepro) 0.15% Azelastine/Fluticasone BECONASE AQ (beclomethasone dipropionate) Flunisolide 0.025%	Non-preferred products may be approved following trial and failure of treatment with three preferred products (failure is defined as lack of efficacy with a 2-week trial, allergy, intolerable side effects or significant drug-drug interactions). Non-preferred combination agents may be approved following trial of individual products with same active ingredients AND trial and failure of one additional preferred agent (failure is defined as lack of efficacy with 2-week trial, allergy, intolerable side effects or significant drug-drug interactions).				

Ipratropium Olopatadine Triamcinolone acetonide (OTC)	Fluticasone (OTC) Mometasone NASONEX (mometasone) OMNARIS (ciclesonide) PATANASE (olopatadine) QNASL (beclomethasone) RYALTRIS (olopatadine/mometasone) XHANCE (fluticasone) ZETONNA (ciclesonide)	
Therapeutic Drug Class: LEUKOTRIENE MODIFIERS – <i>Effective 1/1/2025</i>		
No PA Required Montelukast tablet, chewable	PA Required ACCOLATE (zafirlukast) tablet Montelukast granules SINGULAIR (montelukast) tablet, chewable, granules Zafirlukast tablet Zileuton ER tablet ZYFLO (zileuton) tablet	Non-preferred products may be approved if meeting the following criteria: - Member has trialed and failed treatment with one preferred product (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions) AND - Member has a diagnosis of asthma. Montelukast granules may be approved if a member has tried and failed montelukast chewable tablets AND has difficulty swallowing.
Therapeutic Drug Class: METHOTREXATE PRODUCTS – <i>Effective 1/1/2025</i>		
No PA Required Methotrexate tablet, vial	PA Required JYLAMVO (methotrexate) solution OTREXUP (methotrexate) auto-injector RASUVO (methotrexate) auto-injector REDITREX (methotrexate) syringe TREXALL (methotrexate) tablet	OTREXUP, REDITREX or RASUVO may be approved if meeting the following criteria: - Member has diagnosis of severe, active rheumatoid arthritis OR active polyarticular juvenile idiopathic arthritis (pJIA) OR inflammatory bowel disease (IBD) AND - Member has trialed and failed preferred methotrexate tablet formulation (failure is defined as lack of efficacy, allergy, intolerable side effects, inability to take oral product formulation, or member has a diagnosis of pJIA and provider has determined that the subcutaneous formulation is necessary to optimize methotrexate therapy) AND - Member (or parent/caregiver) is unable to administer preferred methotrexate vial formulation due to limited functional ability (such as vision impairment, limited manual dexterity and/or limited hand strength).

	XATMEP (methotrexate) solution	TREXALL may be approved if meeting the following criteria: - Member has trialed and failed preferred methotrexate tablet formulation. Failure is defined as allergy or intolerable side effects. XATMEP may be approved for members who meet the following criteria: - Member is < 18 years of age - Member has a diagnosis of acute lymphoblastic leukemia OR - Member has a diagnosis of active polyarticular juvenile idiopathic arthritis (pJIA) and has had an insufficient therapeutic response to, or is intolerant to, an adequate trial of first-line therapy including full dose non-steroidal anti-inflammatory agents (NSAIDs) AND - Member has a documented swallowing difficulty due to young age and/or a medical condition and is unable to use the preferred methotrexate tablet formulation <i>Methotrexate can cause serious embryo-fetal harm when administered during pregnancy and it is contraindicated for use during pregnancy for the treatment of non-malignant diseases. Advise members of reproductive potential to use effective contraception during and after treatment with methotrexate, according to FDA product labeling.</i> Members currently stabilized on a non-preferred methotrexate product may receive approval to continue that agent.
Therapeutic Drug Class: MULTIPLE SCLEROSIS AGENTS – Effective 6/5/2025		
Disease Modifying Therapies		
Preferred No PA Required (Unless indicated*) AVONEX (interferon beta 1a) pen, syringe BETASERON (interferon beta 1b) injection COPAXONE (glatiramer) 20mg injection ^{BNR} Dimethyl fumarate tablet, starter pack Fingolimod capsule Glatiramer 40mg injection	Non-Preferred PA Required AUBAGIO (teriflunomide) tablet BAFIERTAM (monomethyl fumarate DR) capsule COPAXONE (glatiramer) 40mg injection EXTAVIA (interferon beta 1b) kit, vial GILENYA (fingolimod) capsule Glatiramer 20mg GLATOPA (glatiramer) injection MAVENCLAD (cladribine) tablet	*Kesimpta (ofatumumab) may be approved if member has trialed and failed treatment with one preferred agent (failure is defined as intolerable side effects, contraindication to therapy, drug-drug interaction, or lack of efficacy). <u>Non-Preferred Products:</u> Non-preferred products may be approved if meeting the following: - Member has a diagnosis of a relapsing form of multiple sclerosis AND - Member has previous trial and failure with three preferred agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND - Prescribed dose does not exceed the maximum FDA-approved dose for the medication being ordered AND - If indicated in the product labeling, a negative pre-treatment pregnancy test has been documented, AND - If indicated in the product labeling, an ophthalmologic examination has been performed and documented prior to medication initiation, AND - The request meets additional criteria listed for any of the following:

*KESIMPTA (ofatumumab) pen **2nd Line** Teriflunomide tablet	MAYZENT (siponimod) tablet, pack PLEGRIDY (peg-interferon beta 1a) pen, syringe PONVORY (ponesimod) tablet, pack REBIF (interferon beta 1a) syringe REBIF REDIDOSE (interferon beta 1a) pen TASCENSO ODT (fingolimod) tablet TECFIDERA (dimethyl fumarate) tablet, pack VUMERITY (diroximel DR) capsule ZEPOSIA (ozanimod) capsule, kit, starter pack	Mayzent (siponimod): - Member has previous trial and failure of three preferred agents, one of which must be Gilenya (fingolimod). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. Mavenclad (cladribine): - Member has history of ≥ 1 relapse in the 12 months preceding initiation of therapy AND - Member has previous trial and failure of three other therapies for relapsing forms of multiple sclerosis (failure is defined as lack of efficacy with 3-month trial, allergy, intolerable side effects, or significant drug-drug interactions) Vumerity (diroximel fumarate) or Bafiertam (monomethyl fumarate DR): - Member has previous trial and failure of three preferred agents, one of which must be Tecfidera (dimethyl fumarate). Failure is defined as lack of efficacy, allergy, significant drug-drug interactions, intolerable side effects (if GI adverse events, must meet additional criteria below) AND - If the requested medication is being prescribed due to GI adverse events with Tecfidera therapy (and no other reason for failure of Tecfidera is given), then the following additional criteria must be met: - Member has trialed a temporary dose reduction of Tecfidera (with maintenance dose being resumed within 4 weeks) AND - Member has trialed taking Tecfidera with food AND - GI adverse events remain significant despite maximized use of gastrointestinal symptomatic therapies (such as calcium carbonate, bismuth subsalicylate, PPIs, H2 blockers, anti-bloating/anti-constipation agents, anti-diarrheal, and centrally acting anti-emetics) AND - Initial authorization will be limited to 3 months. Continuation (12-month authorization) will require documentation of clinically significant reduction in GI adverse events. Members currently stabilized on a preferred second line (Kesimpta) or non-preferred product (may receive approval to continue therapy with that agent).
Symptom Management Therapies		
No PA Required Dalfampridine ER tablet	PA Required AMPYRA ER (dalfampridine) tablet	Non-preferred products may be approved with prescriber attestation that there is clinical rationale supporting why the preferred brand/generic equivalent product formulation is unable to be used. <u>Maximum Dose:</u> Ampyra (dalfampridine) 10mg twice daily

Therapeutic Drug Class: TARGETED IMMUNE MODULATORS – <i>Effective 7/15/2025</i>		
<i>Preferred agents:</i> Adalimumab-aaty and adbm; ADBRY (tralokinumab-ldrm); Cyltezo (adalimumab-adbm); DUPIXENT (dupilumab); ENBREL (etanercept); FASENRA (benralizumab) pen; HADLIMA (adalimumab- bwwd); HUMIRA (adalimumab); OTEZLA (apremilast) tablet; KEVZARA (sarilumab); TALTZ (ixekizumab); TEZSPIRE (tezepelumab-ekko) pen; XELJANZ IR (tofacitinib) tablet; XOLAIR (omalizumab) syringe		
Rheumatoid Arthritis, all other Arthritis (except psoriatic arthritis, see below), and Ankylosing Spondylitis		
Preferred No PA Required (If diagnosis met) (*Must meet eligibility criteria) Adalimumab-aaty pen, syringe Adalimumab-adbm pen, syringe CYLTEZO (adalimumab-adbm) pen, syringe ENBREL (etanercept) HADLIMA (adalimumab-bwwd) Pushtouch, syringe HUMIRA (adalimumab) *KEVZARA (sarilumab) pen, syringe *TALTZ (ixekizumab) 80 mg syringe, autoinjector *TYENNE (tocilizumab-aazg) pen, syringe XELJANZ IR (tofacitinib) tablet	Non-Preferred PA Required ABRILADA (adalimumab-afzb) pen, syringe ACTEMRA (tocilizumab) syringe, Actpen Adalimumab-aacf pen, syringe Adalimumab-adaz pen, syringe Adalimumab-fkjp pen, syringe Adalimumab-ryvk auto-injector AMJEVITA (adalimumab-atto) auto-injector, syringe BIMZELX (bimekizumab-bkzx) pen CIMZIA (certolizumab pegol) syringe, vial COSENTYX (secukinumab) syringe, pen-injector HULIO (adalimumab-fkjp) pen, syringe HYRIMOZ (adalimumab-adaz) pen, syringe IDACIO (adalimumab-aacf) pen, syringe ILARIS (canakinumab) vial KINERET (anakinra) syringe OLUMIANT (baricitinib) tablet ORENCIA (abatacept) clickject, syringe RINVOQ (upadacitinib), solution, tablet	First line preferred agents (preferred adalimumab products, ENBREL, and XELJANZ IR) may receive approval for use for FDA-labeled indications. *TALTZ (ixekizumab) may receive approval for use for FDA-labeled indications following trial and failure‡ of a preferred adalimumab product or ENBREL. *KEVZARA (sarilumab) may receive approval for use for FDA-labeled indications following trial and failure‡ of: • A preferred adalimumab product or ENBREL AND • XELJANZ IR. *TYENNE (tocilizumab-aazg) may receive approval for use for FDA-labeled indications following trial and failure‡ of: • A preferred adalimumab product or ENBREL AND • XELJANZ IR. <u>Quantity Limit:</u> XELJANZ IR is limited to 2 tablets per day or 60 tablets for a 30-day supply <u>Non-Preferred Agents:</u> COSENTYX (secukinumab) may receive approval for: • FDA-labeled indications following trial and failure‡ of all indicated preferred agents OR • Treatment of enthesitis-related arthritis if meeting the following: ○ Member is ≥ 4 years of age and weighs ≥ 15 kg AND ○ Member has had trialed and failed‡ NSAID therapy and ENBREL and a preferred adalimumab product KINERET (anakinra) may receive approval for: • Treatment of systemic juvenile idiopathic arthritis (sJIA) or Adult-Onset Still's Disease (AOSD) OR • Treatment of rheumatoid arthritis following trial and failure‡ of ○ A preferred adalimumab product or ENBREL AND ○ XELJANZ IR ILARIS (canakinumab) may receive approval if meeting the following:

	SIMLANDI (adalimumab-ryvk) auto-injector SIMPONI (golimumab) pen, syringe SKYRIZI (risankizumab-rzaa) OnBody, SC pen, syringe XELJANZ (tofacitinib) solution XELJANZ XR (tofacitinib ER) tablet YUFLYMA (adalimumab-aaty) auto-injector, syringe YUSIMRY (adalimumab-aqvh) pen <i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i>	- Medication is being prescribed for systemic juvenile idiopathic arthritis (sJIA) or Adult-Onset Still's Disease (AOSD), AND - Member has trialed and failed‡ a tocilizumab product. Quantity Limit: 300mg (2mL) every 4 weeks XELJANZ (tofacitinib) XR approval will require verification of the clinically relevant reason for use of the XELJANZ XR formulation versus the XELJANZ IR formulation, in addition to meeting non-preferred criteria listed below. XELJANZ (tofacitinib) oral solution may be approved when the following criteria are met: - Member has a diagnosis of polyarticular course juvenile idiopathic arthritis (pJIA) who require a weight-based dose for <40 kg following trial and failure‡ of a preferred adalimumab product or ENBREL OR - Member cannot swallow a tofacitinib tablet All other non-preferred agents may receive approval for FDA-labeled indications following trial and failure‡ of all preferred agents that are FDA-indicated or have strong evidence supporting use for the prescribed indication from clinically recognized guideline compendia (only one preferred adalimumab product trial required). Non-preferred agents that are being prescribed per FDA labeling to treat non-radiographic axial spondyloarthritis (nr-axSpA) will require trial and failure‡ of preferred agents that are FDA-labeled for treating an axial spondyloarthritis condition, including ankylosing spondylitis (AS) or nr-axSpA. <u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent may receive approval for continuation of therapy with the prescribed agent. ‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction. Note that trial and failure of preferred TNF inhibitors will not be required when prescribed to treat polyarticular juvenile idiopathic arthritis (pJIA) in members with documented clinical features of lupus. <i>The Department would like to remind providers that many products are associated with patient-centered programs that are available to assist with drug administration, education, and emotional support related to our members' various disease states.</i>
Psoriatic Arthritis		
Preferred	Non-Preferred	

No PA Required (If diagnosis met) (*Must meet eligibility criteria)	PA Required	
Adalimumab-aaty pen, syringe Adalimumab-adbm pen, syringe CYLTEZO (adalimumab-adbm) pen, syringe ENBREL (etanercept) HADLIMA (adalimumab-bwwd) Pushtouch, syringe HUMIRA (adalimumab) *OTEZLA (apremilast) tablet *TALTZ (ixekizumab) 80 mg syringe XELJANZ IR (tofacitinib) tablet	ABRILADA (adalimumab-afzb) pen, syringe Adalimumab-aacf pen, syringe Adalimumab-adaz pen, syringe Adalimumab-fkjp pen, syringe Adalimumab-ryvk auto-injector AMJEVITA (adalimumab-atto) auto-injector, syringe BIMZELX (bimekizumab-bkzx) pen CIMZIA (certolizumab pegol) syringe, vial COSENTYX (secukinumab) syringe, pen-injector HULIO (adalimumab-fkjp) pen, syringe HYRIMOZ (adalimumab-adaz) pen, syringe IDACIO (adalimumab-aacf) pen, syringe IMULDOSA (ustekinumab-SRLF) syringe, vial ORENCIA (abatacept) syringe, clickject OTULFI (ustekinumab-aaaz) syringe PYZCHIVA (ustekinumab-ttwe) syringe RINVOQ (upadacitinib) tablet RINVOQ LQ (upadacitinib) solution SELARSDI (ustekinumab-AEKN) syringe SIMLANDI (adalimumab-ryvk) auto-injector SIMPONI (golimumab) pen, syringe	First line preferred agents (HADLIMA, HUMIRA, ENBREL, XELJANZ IR) may receive approval for psoriatic arthritis indication. *OTEZLA (apremilast) may receive approval for psoriatic arthritis indication following trial and failure‡ of: - A preferred adalimumab product or ENBREL AND - XELJANZ IR or TALTZ. *TALTZ (ixekizumab) may receive approval for psoriatic arthritis indication following trial and failure‡ of: - A preferred adalimumab product or ENBREL AND - XELJANZ IR or OTEZLA. Quantity Limit: XELJANZ IR is limited to 2 tablets per day or 60 tablets for a 30-day supply <u>Non-Preferred Agents:</u> COSENTYX (secukinumab) may receive approval for psoriatic arthritis indication for members ≥ 2 years of age and weighing ≥ 15 kg following trial and failure‡ of: - A preferred adalimumab product or ENBREL AND - XELJANZ IR AND - TALTZ or OTEZLA. USTEKINUMAB (Stelara brand/generic and biosimilar agents) syringe for subcutaneous use may receive approval if meeting the following: - The request meets one of the following: - The prescribed agent is one of the following favored Ustekinumab products: Imuldosa, Otulfi, Pyzchiva, Selarsdi, Steqeyma, Ustekinumab (generic Stelara), Ustekinumab-AEKN, Yesintek OR - If the prescribed agent is brand Stelara or a product that is not favored Ustekinumab product, then the member has trialed and failed‡ at least one favored Ustekinumab product AND - Member has trial and failure‡ of: - A preferred adalimumab product or ENBREL AND - XELJANZ IR AND - TALTZ or OTEZLA AND - Prior authorization approval may be given for an initial 16-week supply and authorization approval for continuation may be provided based on clinical response.

	SKYRIZI (risankizumab-rzaa) OnBody, pen, syringe STELARA (ustekinumab) syringe STEQEYMA (ustekinumab-stba) syringe TREMFYA (guselkumab) pen, injector, syringe Ustekinumab (generic Stelara, TTWE, AEKN) syringe, vial WEZLANA (ustekinumab-auub) syringe, vial XELJANZ (tofacitinib) solution XELJANZ XR (tofacitinib ER) tablet YESINTEK (ustekinumab-kfce) syringe, vial YUFLYMA (adalimumab-aaty) auto-injector, syringe YUSIMRY (adalimumab-aqvh) pen <i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i>	XELJANZ (tofacitinib) XR approval will require verification of the clinically relevant reason for use of the XELJANZ XR formulation versus the XELJANZ IR formulation, in addition to meeting non-preferred criteria listed below. All other non-preferred agents may receive approval for psoriatic arthritis following trial and failure‡ of: • A preferred adalimumab product or ENBREL AND • XELJANZ IR AND • TALTZ or OTEZLA. ‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction. <u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent may receive approval for continuation of therapy with the prescribed agent. <i>The Department would like to remind providers that many products are associated with patient-centered programs that are available to assist with drug administration, education, and emotional support related to our members' various disease states.</i>
Plaque Psoriasis		
Preferred No PA Required (If diagnosis met) (*Must meet eligibility criteria) Adalimumab-aaty pen, syringe Adalimumab-adbm pen, syringe CYLTEZO (adalimumab-adbm) pen, syringe ENBREL (etanercept) HADLIMA (adalimumab-bwwd) Pushtouch, syringe	Non-Preferred PA Required ABRILADA (adalimumab-afzb) pen, syringe Adalimumab-aacf pen, syringe Adalimumab-adaz pen, syringe Adalimumab-fkjp pen, syringe Adalimumab-ryvk auto-injector AMJEVITA (adalimumab-atto) auto-injector, syringe BIMZELX (bimekizumab-bkzx) pen	First line preferred agents (preferred adalimumab products, ENBREL) may receive approval for plaque psoriasis indication. *Second line preferred agents (TALTZ, OTEZLA) may receive approval for plaque psoriasis indication following trial and failure‡ of a preferred adalimumab product OR ENBREL. <u>Non-Preferred Agents:</u> USTEKINUMAB (Stelara brand/generic and biosimilar agents) syringe for subcutaneous use may receive approval if meeting the following: • The request meets one of the following: ○ The prescribed agent is one of the following favored Ustekinumab products: Imuldosa, Otulfi, Pyzchiva, Selarsdi, Steqeyma, Ustekinumab (generic Stelara), Ustekinumab-AEKN, Yesintek OR

HUMIRA (adalimumab) *OTEZLA (apremilast) tablet *TALTZ (ixekizumab) 80 mg syringe TYENNE (tocilizumab-aazg) pen, syringe	CIMZIA (certolizumab pegol) syringe, vial COSENTYX (secukinumab) syringe, pen-injector HULIO (adalimumab-fkjp) pen, syringe HYRIMOZ (adalimumab-adaz) pen, syringe IDACIO (adalimumab-aacf) pen, syringe IMULDOSA (ustekinumab-SRLF) syringe, vial ORENCIA (abatacept) syringe, clickject OTULFI (ustekinumab-aauz) syringe PYZCHIVA (ustekinumab-ttwe) syringe SELARSDI (ustekinumab-AEKN) syringe SILIQ (brodalumab) syringe SIMLANDI (adalimumab-ryvk) auto-injector SKYRIZI (risankizumab-rzaa) OnBody, pen, syringe SOTYKTU (ducravacitinib) oral tablet STELARA (ustekinumab) syringe STEQEYMA (ustekinumab-stba) syringe TALTZ (ixekizumab) 20mg, 40mg syringe TREMFYA (guselkumab) injector, syringe Ustekinumab (generic Stelara, TTWE, AEKN) syringe, vial WEZLANA (ustekinumab-auub) syringe, vial YESINTEK (ustekinumab-kfce) syringe, vial	○ If the prescribed agent is brand Stelara or a product that is not favored Ustekinumab product, then the member has trialed and failed‡ at least one favored Ustekinumab product AND • Member has trial and failure‡ of one indicated first line agent (preferred adalimumab products, ENBREL) AND two indicated second line agents (TALTZ, OTEZLA) AND • Prior authorization approval may be given for an initial 16-week supply and authorization approval for continuation may be provided based on clinical response. All other non-preferred agents may receive approval for plaque psoriasis indication following trial and failure‡ of one indicated first line agent (a preferred adalimumab product, ENBREL) AND two second line agents (TALTZ, OTEZLA). ‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction. <u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent may receive approval for continuation of therapy with the prescribed agent. <i>The Department would like to remind providers that many products are associated with patient-centered programs that are available to assist with drug administration, education, and emotional support related to our members' various disease states.</i>
--	--	--

	YUFLYMA (adalimumab-aaty) auto-injector, syringe YUSIMRY (adalimumab-aqvh) pen <i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i>	
Crohn's Disease and Ulcerative Colitis		
Preferred No PA Required (If diagnosis met) (*Must meet eligibility criteria) Adalimumab-aaty pen, syringe Adalimumab-adbm pen, syringe CYLTEZO (adalimumab-adbm) pen, syringe HADLIMA (adalimumab-bwwd) Pushtouch, syringe HUMIRA (adalimumab) *XELJANZ IR (tofacitinib) tablet	Non-Preferred PA Required ABRILADA (adalimumab-afzb) pen, syringe Adalimumab-aacf pen, syringe Adalimumab-adaz pen, syringe Adalimumab-fkjp pen, syringe Adalimumab-ryvk auto-injector AMJEVITA (adalimumab-atto) auto-injector, syringe CIMZIA (certolizumab pegol) syringe, vial COSENTYX (secukinumab) syringe, pen-injector ENTYVIO (vedolizumab) pen HULIO (adalimumab-fkjp) syringe HYRIMOZ (adalimumab-adaz) pen, syringe IDACIO (adalimumab-aacf) pen, syringe IMULDOSA (ustekinumab-SRLF) syringe, vial OLUMIANT (baricitinib) tablet OMVOH (mirikizumab-mrkz) pen OTULFI (ustekinumab-aauz) syringe	Preferred agents (preferred adalimumab products, XELJANZ IR) may receive approval for Crohn's disease and ulcerative colitis indications. Quantity Limit: XELJANZ IR is limited to 2 tablets per day or 60 tablets for a 30-day supply Non-Preferred Agents: ENTYVIO (vedolizumab) pen for subcutaneous injection may receive approval if the following criteria are met: - For treatment of moderately-to-severely active Crohn's disease, member has trial and failure‡ of one preferred adalimumab product OR for treatment of moderately-to-severely active ulcerative colitis, member has trial and failure‡ of one preferred adalimumab product and XELJANZ IR AND - Member is ≥ 18 years of age AND - Prescriber acknowledges that administration of IV induction therapy prior to approval of ENTYVIO (vedolizumab) pen for subcutaneous injection using the above criteria should be avoided and will not result in an automatic approval of requests for these formulations. OMVOH (mirikizumab-mrkz) pen for subcutaneous injection may receive approval if the following criteria are met: - The requested medication is being prescribed for treatment of moderately-to-severely active ulcerative colitis AND - Member is ≥ 18 years of age AND - Member has trial and failure‡ of one preferred adalimumab product AND XELJANZ IR AND ENTYVIO (vedolizumab) AND - Prescriber acknowledges that administration of IV induction therapy prior to approval of OMVOH (mirikizumab-mrkz) pen for subcutaneous injection using the above criteria should be avoided and will not result in an automatic approval of requests for these formulations. SKYRIZI (risankizumab) syringe for subcutaneous use and on-body injector formulations may receive approval if meeting the following:

	PYZCHIVA (ustekinumab-ttwe) syringe RINVOQ (upadacitinib) tablet RINVOQ LQ (upadacitinib) solution SELARSDI (ustekinumab-AEKN) syringe SIMLANDI (adalimumab-ryvk) auto-injector SIMPONI (golimumab) pen, syringe SKYRIZI (risankizumab-rzaa) OnBody, pen, syringe STELARA (ustekinumab) syringe, vial STEQEYMA (ustekinumab-stba) syringe Ustekinumab (generic Stelara, TTWE, AEKN) syringe, vial VELSIPITY (etrasimod) tablet WEZLANA (ustekinumab-auub) syringe, vial XELJANZ (tofacitinib) solution XELJANZ XR (tofacitinib ER) tablet YESINTEK (ustekinumab-kfce) syringe, vial YUFLYMA (adalimumab-aaty) auto-injector YUSIMRY (adalimumab-aqvh) pen ZYMFENTRA (infliximab-dyyb) pen kit, syringe kit <i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i>	- The requested medication is being prescribed for use for treating moderately-to-severely active Crohn’s disease or for treating moderate-to-severely ulcerative colitis AND - Member is ≥ 18 years of age AND - Request meets one of the following based on prescribed indication: - For treatment of moderately-to-severely active Crohn’s disease, member has trial and failure‡ of one preferred adalimumab product and ENTYVIO (vedolizumab) OR - For treatment of moderately-to-severely active ulcerative colitis, member has trial and failure‡ of one preferred adalimumab product and XELJANZ IR and ENTYVIO (vedolizumab) AND - Prescriber acknowledges that administration of IV induction therapy prior to approval of SKYRIZI (risankizumab) prefilled syringe or on-body injector formulation using the above criteria should be avoided and will not result in an automatic approval of requests for these formulations. Dosing Limit: SKYRIZI on-body formulation maintenance dosing is limited to one 360 mg/2.4 mL single-dose prefilled cartridge or one 180 mg/1.2mL prefilled cartridge every 8 weeks. USTEKINUMAB (Stelara brand/generic and biosimilar agents) syringe for subcutaneous use may receive approval if meeting the following: - The request meets one of the following: - The prescribed agent is one of the following favored Ustekinumab products: Imuldosa, Otulfi, Pyzchiva, Selarsdi, Steqeyma, Ustekinumab (generic Stelara), Ustekinumab-AEKN, Yesintek OR - If the prescribed agent is brand Stelara or a product that is not favored Ustekinumab product, then the member has trialed and failed‡ at least one favored Ustekinumab product AND - The requested medication is being prescribed for use for treating moderately-to-severely active Crohn’s disease or for treating moderately-to-severely active ulcerative colitis AND - Request meets one of the following based on prescribed indication: - For treatment of moderately-to-severely active Crohn’s disease, member has trial and failure‡ of one preferred adalimumab product and ENTYVIO (vedolizumab) OR - For treatment of moderately-to-severely active ulcerative colitis, member has trial and failure‡ of one preferred adalimumab product and XELJANZ IR and ENTYVIO (vedolizumab) AND - The member is ≥ 18 years of age AND - Prescriber acknowledges that loading dose administration prior to approval of ustekinumab for maintenance therapy using the above criteria should be avoided
--	--	--

		and will not result in an automatic approval of ustekinumab for maintenance therapy AND • Prior authorization approval may be given for an initial 16-week supply and authorization approval for continuation may be provided based on clinical response. TREMFYA (guselkumab) pen for subcutaneous injection may receive approval if the following criteria are met: • For treatment of moderately-to-severely active ulcerative colitis, member has trial and failure‡ of one preferred adalimumab product and XELJANZ IR AND • Member is ≥ 18 years of age AND • Prescriber acknowledges that administration of IV induction therapy prior to approval of TREMFYA (guselkumab) pen for subcutaneous injection using the above criteria should be avoided and will not result in an automatic approval of requests for these formulations. XELJANZ (tofacitinib) XR approval will require verification of the clinically relevant reason for use of the XELJANZ XR formulation versus the XELJANZ IR formulation, in addition to meeting non-preferred criteria listed below. All other non-preferred agents may receive approval for FDA-labeled indications if meeting the following: • The requested medication is being prescribed for treating moderately-to-severely active Crohn’s disease or moderately-to-severely active Ulcerative Colitis in alignment with indicated use outlined in FDA-approved product labeling AND • The requested medication meets FDA-labeled indicated age for prescribed use AND • For treatment of moderately-to-severely active Crohn’s disease, member has trial and failure‡ of one preferred adalimumab product OR for treatment of moderately-to-severely active ulcerative colitis, member has trial and failure‡ of one preferred adalimumab product and XELJANZ IR. <u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent may receive approval for continuation of therapy with the prescribed agent. ‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction. Note that trial and failure of Xeljanz IR will not be required when prescribed for ulcerative colitis for members ≥ 50 years of age that have an additional CV risk factor.
--	--	---

		<i>The Department would like to remind providers that many products are associated with patient-centered programs that are available to assist with drug administration, education, and emotional support related to our members' various disease states.</i>
Asthma		
Preferred PA Required (*Must meet eligibility criteria) *DUIXENT (dupilumab) pen, syringe *FASENRA (benralizumab) pen *TEZSPIRE (tezepelumab-ekko) pen *XOLAIR (omalizumab) syringe, autoinjector	Non-Preferred PA Required NUCALA (mepolizumab) auto-injector, syringe <i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i>	*Preferred products (Dupixent, Fasentra, Tezspire, Xolair) may receive approval if meeting the following: DUPIXENT (dupilumab): - Member is 6 years of age or older AND - Member has an FDA-labeled indicated use for treating one of the following: - Moderate to severe asthma (on medium to high dose inhaled corticosteroid and a long-acting beta agonist) with eosinophilic phenotype based on a blood eosinophil level of $\geq 150/\text{mcL}$ OR - Oral corticosteroid dependent asthma AND - Member's asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies AND - Medication is being prescribed as add-on therapy to existing asthma regimen. <u>Quantity Limit:</u> 2 syringes every 28 days after initial 14 days of therapy (first dose is twice the regular scheduled dose) FASENRA (benralizumab): - Member is ≥ 6 years of age AND - Member has an FDA-labeled indicated use for treating severe asthma with an eosinophilic phenotype based on a blood eosinophil level of $\geq 150/\text{mcL}$ AND - Member's asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies AND - The requested medication is being prescribed as add-on therapy to existing asthma regimen. <u>Quantity Limit:</u> One 30 mg unit dose pack every 28 days for the first 3 doses and then every 8 weeks thereafter TEZSPIRE (tezepelumab-ekko): - Member is ≥ 12 years of age AND - Member has a diagnosis of severe asthma AND - Member's asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies AND - The requested medication is being prescribed as add-on therapy to existing asthma regimen. <u>Quantity Limit:</u> Four 210 mg unit dose packs every 28 days

		XOLAIR (omalizumab) may receive approval if meeting the following based on prescribed indication: • Member is ≥ 6 years of age AND • Member has an FDA-labeled indicated use for treating asthma AND • Member has a positive skin test or in vitro reactivity to a perennial inhaled allergen or has a pre-treatment IgE serum concentration ≥ 30 IU/mL AND • Member's asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies AND • The requested medication is being prescribed as add-on therapy to existing asthma regimen. <u>Non-Preferred Agents:</u> Non-preferred FDA-indicated biologic agents for asthma may receive approval if meeting the following: • The requested medication is being prescribed for treating asthma in alignment with indicated use outlined in FDA-approved product labeling (including asthma type and severity) AND • If prescribed for use for asthma with eosinophilic phenotype, member has a blood eosinophil count ≥ 150 cells/mcL AND • The requested medication meets FDA-labeled indicated age for prescribed use AND • Member's asthma has been refractory to recommended evidence-based, guideline-supported pharmacologic therapies AND • The requested medication is being prescribed as add-on therapy to existing asthma regimen AND • Member has trialed and failed‡ two preferred agents. <u>Quantity Limits:</u> Non-preferred medications will be subject to quantity limitations in alignment with FDA-approved dosing per product package labeling. Nucala (mepolizumab) is limited to 100mg every 4 weeks (members ≥ 12 years of age) or 40mg every 4 weeks (members 6-11 years of age). ‡Failure is defined as a lack of efficacy, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions. <u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent may receive approval for continuation of therapy with the prescribed agent.
Atopic Dermatitis		

Preferred (*Must meet eligibility criteria)	Non-Preferred PA Required	*Preferred products (Adbry and Dupixent) may receive approval if meeting the following:
*ADBRY (tralokinumab-ldrm) syringe, autoinjector *DUPIXENT (dupilumab) pen, syringe	CIBINQO (abrocitinib) tablet RINVOQ (upadacitinib) tablet <i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i>	ADBRY (tralokinumab-ldrm): - The requested drug is being prescribed for moderate-to-severe atopic dermatitis AND - Member has trialed and failed‡ the following agents: - One medium potency to very-high potency topical corticosteroid (such as mometasone furoate, betamethasone dipropionate) AND - One topical calcineurin inhibitor (such as pimecrolimus or tacrolimus) <u>Maximum Dose:</u> 600 mg/2 weeks <u>Quantity Limit:</u> Four 150 mg/mL prefilled syringes/2 weeks DUPIXENT (dupilumab): - Member has a diagnosis of moderate to severe atopic dermatitis AND - Member has trialed and failed‡ the following agents: - One medium potency to very-high potency topical corticosteroid [such as mometasone furoate, betamethasone dipropionate, or fluocinonide (see PDL for list of preferred products)] AND - One topical calcineurin inhibitor (such as pimecrolimus or tacrolimus) <u>Quantity Limit:</u> 2 syringes every 28 days after initial 14 days of therapy (first dose is twice the regular scheduled dose) <u>Non-Preferred Agents:</u> Non-preferred agents indicated for the treatment of atopic dermatitis may receive approval if meeting the following: - Member has a diagnosis of moderate to severe chronic atopic dermatitis AND - Member has trialed and failed‡ therapy with two preferred agents for the prescribed indication AND - Member has trialed and failed‡ the following agents: - One medium potency to very-high potency topical corticosteroid (such as mometasone furoate, betamethasone dipropionate, or fluocinonide) - One topical calcineurin inhibitor (such as pimecrolimus and tacrolimus) AND - The medication is being prescribed by or in consultation with a dermatologist, allergist, immunologist, or rheumatologist. <u>Approval:</u> One year

		‡Failure is defined as a lack of efficacy, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions. <u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent may receive approval for continuation of therapy with the prescribed agent.
Other indications		
Preferred (If diagnosis met, No PA required) (Must meet eligibility criteria*) *DUPIXENT (dupilumab) pen, syringe ENBREL (etanercept) *FASENRA (benralizumab) pen HUMIRA (adalimumab) *KEVZARA (sarilumab) OTEZLA (apremilast) tablet XELJANZ IR (tofacitinib) tablet *XOLAIR (omalizumab) syringe, autoinjector	Non-Preferred PA Required ACTEMRA (tocilizumab) syringe, Actpen ARCALYST (rilonacept) injection CIMZIA (certolizumab pegol) syringe COSENTYX (secukinumab) syringe, pen-injector CYLTEZO (adalimumab-adbm) pen, syringe ILARIS (canakinumab) vial KINERET (anakinra) syringe NUCALA (mepolizumab) auto-injector, syringe OLUMIANT (baricitinib) tablet YUFLYMA (adalimumab-aaty) auto-injector <i>Note: Product formulations in the physician administered drug (PAD) category are located on Appendix P</i>	*DUPIXENT (dupilumab) may receive approval if meeting the following based on prescribed indication: <u>Chronic Idiopathic Urticaria</u> - Member is 12 years of age or older AND - Member is diagnosed with chronic idiopathic urticaria AND - Member is symptomatic despite H1 antihistamine treatment AND - Member has tried and failed‡ at least three of the following - High-dose second generation H1 antihistamine - H2 antihistamine - First-generation antihistamine - Leukotriene receptor antagonist - Hydroxyzine or doxepin <u>Chronic Obstructive Pulmonary Disease</u> - Member is ≥ 18 years of age AND - Medication is being prescribed by or in consultation with a pulmonologist or allergist AND - Requested medication is being prescribed as an add-on maintenance treatment for inadequately controlled chronic obstructive pulmonary disease (COPD) AND - Member's COPD is an eosinophilic phenotype based on a blood eosinophil level of ≥ 300 cells/mcL AND - Member is receiving, and will continue, standard maintenance triple therapy for COPD (inhaled corticosteroid, long-acting muscarinic agent, long-acting beta agonist) as recommended by the current Global Initiative for Chronic Obstructive Lung Disease (GOLD) guidelines AND - Member has experienced at least 2 moderate OR 1 severe COPD exacerbation during the past 12 months <u>Chronic Rhinosinusitis with Nasal Polyposis</u> - Member is ≥ 12 years of age AND - Medication is being prescribed as an add-on maintenance treatment for inadequately controlled chronic rhinosinusitis with nasal polyposis (CRSwNP) AND

		- Member has trialed and failed‡ therapy with at least two intranasal corticosteroid regimens <u>Eosinophilic Esophagitis (EoE):</u> - Member is ≥ 1 year of age AND - Member weighs at least 15 kg AND - Member has a diagnosis of eosinophilic esophagitis (EoE) with ≥ 15 intraepithelial eosinophils per high-power field (eos/hpf), with or without a history of esophageal dilations AND - Member is following appropriate dietary therapy interventions AND - Medication is being prescribed by or in consultation with a gastroenterologist, allergist or immunologist AND - Member has trialed and failed‡ one of the following treatment options for EoE: - Proton pump inhibitor trial of at least eight weeks in duration if reflux is a contributing factor OR - Minimum four-week trial of local therapy with a corticosteroid medication <u>Prurigo Nodularis:</u> - Member is ≥ 18 years of age AND - Medication is being prescribed as treatment for prurigo nodularis AND - Member has trialed and failed‡ therapy with at least two corticosteroid regimens (topical or intralesional injection). *FASENRA (benralizumab) may be approved for the treatment of adult patients with eosinophilic granulomatosis with polyangiitis (EGPA). *KEVZARA (sarilumab) treatment of adult patients with polymyalgia rheumatica who have had an inadequate response to corticosteroids or who cannot tolerate corticosteroid taper. TYENNE (tocilizumab-aazg) may receive approval for use for FDA-label indications following trial and failure‡ of a preferred adalimumab product or ENBREL *XOLAIR (omalizumab) may receive approval if meeting the following based on prescribed indication: <u>Chronic Rhinosinusitis with Nasal Polyps:</u> - Member is 18 years of age or older AND - Medication is being prescribed as add-on maintenance treatment of chronic rhinosinusitis with nasal polyps (CRSwNP) in adult patients 18 years of age and older with inadequate response to nasal corticosteroids AND
--	--	--

		- Member has tried and failed‡ therapy with at least two intranasal corticosteroid regimens <u>Chronic Idiopathic Urticaria (CIU):</u> - Member is 12 years of age or older AND - Member is diagnosed with chronic idiopathic urticaria AND - Member is symptomatic despite H1 antihistamine treatment AND - Member has tried and failed‡ at least three of the following: - High-dose second generation H1 antihistamine - H2 antihistamine - First-generation antihistamine - Leukotriene receptor antagonist - Hydroxyzine or doxepin (must include) AND - Prescriber attests that the need for continued therapy will be periodically reassessed (as the appropriate duration of Xolair therapy for CIU has currently not been evaluated). <u>IgE-Mediated Food Allergy:</u> - Medication is being prescribed for reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods in adult and pediatric patients aged 1 year and older with IgE-mediated food allergy. All other preferred agents (preferred adalimumab products, ENBREL, OTEZLA) may receive approval for use for FDA-labeled indications. <u>Non-Preferred Agents:</u> ARCALYST (rilonacept) may receive approval if meeting the following: - Medication is being prescribed for one of the following autoinflammatory periodic fever syndromes (approval for all other indications is subject to meeting non-preferred criteria listed below): - Cryopyrin-associated Autoinflammatory Syndrome (CAPS), including: - Familial Cold Autoinflammatory Syndrome (FCAS) - Muckle-Wells Syndrome (MWS) - Maintenance of remission of Deficiency of Interleukin-1 Receptor Antagonist (DIRA) in adults and pediatric patients weighing at least 10 kg - Treatment of recurrent pericarditis and reduction in risk of recurrence in adults and children ≥ 12 years of age AND
--	--	---

		• Member has trialed and failed‡ colchicine AND • Initial approval will be given for 12 weeks and authorization approval for continuation will be provided based on clinical response. ILARIS (canakinumab) may receive approval if meeting the following: • Medication is being prescribed for one of the following (approval for all other indications is subject to meeting non-preferred criteria listed below): ○ Familial Mediterranean Fever (FMF) ○ Hyperimmunoglobulinemia D syndrome (HIDS) ○ Mevalonate Kinase Deficiency (MKD) ○ Neonatal onset multisystem inflammatory disease (NOMID) ○ TNF Receptor Associated Periodic Syndrome (TRAPS) ○ Cryopyrin-associated Autoinflammatory Syndrome (including Familial Cold Autoinflammatory Syndrome and Muckle-Wells Syndrome) ○ Symptomatic treatment of adult patients with gout flares in whom NSAIDs and colchicine are contraindicated, are not tolerated, or do not provide an adequate response, and in whom repeated courses of corticosteroids are not appropriate (limited to four 150mg doses per one year approval) AND • Member has trialed and failed‡ colchicine. • Quantity Limits: ○ Cryopyrin-associated periodic syndrome: 600mg (4mL) every 8 weeks ○ All other indications: 300mg (2mL) every 4 weeks KINERET (anakinra) may receive approval if meeting the following: • Medication is being prescribed for one of the following indications (approval for all other indications is subject to meeting non-preferred criteria below): ○ Neonatal onset multisystem inflammatory disease (NOMID). ○ Familial Mediterranean Fever (FMF) AND • Member has trialed and failed‡ colchicine. NUCALA (mepolizumab) may receive approval if meeting the following based on prescribed indication (for any FDA-labeled indications in this subclass category that are not listed, approval is subject to meeting non-preferred criteria listed below): <u>Chronic Rhinosinusitis with Nasal Polyps:</u> • Member is 18 years of age or older AND • Medication is being prescribed as an add-on maintenance treatment in adult patients with inadequately controlled chronic rhinosinusitis with nasal polyposis (CRSwNP) AND
--	--	---

		• Member has a baseline bilateral endoscopic nasal polyps score (NPS; scale 0-8) AND nasal congestion/obstruction score (NC; scale 0-3) averaged over 28-day period AND • Member has trialed and failed[‡] therapy with three intranasal corticosteroids (see PDL Class) AND • Medication is being prescribed by or in consultation with a rheumatologist, allergist, ear/nose/throat specialist or pulmonologist AND • Initial authorization will be for 24 weeks, for additional 12-month approval member must meet the following criteria: ○ NC and NPS scores are provided and show a 20% reduction in symptoms from baseline AND ○ Member continues to use primary therapies such as intranasal corticosteroids. <u>Eosinophilic Granulomatosis with polyangiitis (EGPA):</u> • Member is 18 years of age or older AND • Member has been diagnosed with relapsing or refractory EGPA at least 6 months prior to request as demonstrated by ALL the following: ○ Member has a diagnosis of asthma AND ○ Member has a blood eosinophil count of greater than or equal to 1000 cells/mcL or a blood eosinophil level of 10% AND • Member has the presence of two of the following EGPA characteristics: ○ Histopathological evidence of eosinophilic vasculitis, perivascular eosinophilic infiltration, or eosinophil-rich granulomatous inflammation ○ Neuropathy ○ Pulmonary infiltrates ○ Sinonasal abnormality ○ Cardiomyopathy ○ Glomerulonephritis ○ Alveolar hemorrhage ○ Palpable purpura ○ Antineutrophil cytoplasmic antibody (ANCA) positive AND • Member has trialed and failed[‡] Fasenra (benralizumab) AND • Dose of NUCALA (mepolizumab) 300 mg once every 4 weeks is being prescribed. <u>Hypereosinophilic Syndrome (HES):</u> • Member is 12 years of age or older AND • Member has a diagnosis for HES for at least 6 months that is nonhematologic secondary HES AND
--	--	---

		- Member has a blood eosinophil count of greater than or equal to 1000 cells/mcL AND - Member has a history of two or more HES flares (defined as worsening clinical symptoms or blood eosinophil counts requiring an increase in therapy) AND - Member has been on stable dose of HES therapy for at least 4 weeks, at time of request, including at least one of the following: - Oral corticosteroids - Immunosuppressive therapy - Cytotoxic therapy AND - Dose of 300 mg once every 4 weeks is being prescribed. All other non-preferred agent indications may receive approval for FDA-labeled use following trial and failure‡ of all preferred agents that are FDA-indicated or have strong evidence supporting use for the prescribed indication from clinically recognized guideline compendia (only one preferred adalimumab product trial required). ‡Failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction. <u>Continuation of therapy:</u> Members currently taking a preferred agent may receive approval to continue therapy with that agent. Members with current prior authorization approval on file for a non-preferred agent will be subject to meeting reauthorization criteria above when listed for the prescribed indication, or if reauthorization criteria are not listed for the prescribed indication, may receive approval for continuation of therapy. <u>Note:</u> Prior authorization requests for OLUMIANT (baricitinib) prescribed solely for treating alopecia areata will not be approved. <i>The Department would like to remind providers that many products are associated with patient-centered programs that are available to assist with drug administration, education, and emotional support related to our members' various disease states.</i>
X. Miscellaneous		
Therapeutic Drug Class: EPINEPHRINE PRODUCTS – Effective 1/1/2025		
No PA Required <i>Brand/generic changes effective 02/22/2024*</i> *Epinephrine 0.15mg/0.15ml, 0.3mg/0.3ml auto-injector (Mylan only)	PA Required AUVI-Q (epinephrine) auto-injector Epinephrine 0.15mg/0.15ml, 0.3mg/0.3ml auto-injector (All other manufacturers; generic Adrenaclick, Epipen)	Non-preferred products may be approved if the member has failed treatment with one of the preferred products. Failure is defined as allergy to ingredients in product or intolerable side effects. Quantity limit: 4 auto-injectors per year unless used / damaged / lost

EPIPEN 0.3 mg/0.3 ml (epinephrine) auto-injector EPIPEN JR 0.15 mg/0.15 ml, (epinephrine) auto-injector	SYMJEPI 0.15mg/0.3ml, 0.3mg/0.3ml (epinephrine) syringe	
Therapeutic Drug Class: NEWER HEREDITARY ANGIOEDEMA PRODUCTS – <i>Effective 1/1/2025</i>		
PA Required for all agents in this class		<u>Medications Indicated for Routine Prophylaxis:</u>
Preferred <u>Prophylaxis:</u> CINRYZE (C1 esterase inhibitor) kit HAEGARDA (C1 esterase inhibitor) vial <u>Treatment:</u> BERINERT (C1 esterase inhibitor) kit, vial FIRAZYR (icatibant acetate) syringe^{BNR}	Non-Preferred <u>Prophylaxis:</u> Andembry (garadacimab-gxii) autoinjector ORLADEYO (berotralstat) oral capsule TAKHZYRO (lanadelumab-flyo) syringe, vial <u>Treatment:</u> Ekterly (sebetralstat) tablet Icatibant syringe (generic FIRAZYR) RUCONEST (C1 esterase inhibitor, recomb) vial	Members are restricted to coverage of one medication for <u>routine prophylaxis</u> at one time. Prior authorization approval will be for one year. HAEGARDA (C1 esterase inhibitor - human) may be approved for members meeting the following criteria: ○ Member has a diagnosis of HAE Type I or Type II confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, C1-INH level) OR has a diagnosis of HAE Type III based on clinical presentation AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member meets at least one of the following: ▪ Haegarda is being used for short-term prophylaxis to undergo a surgical procedure or major dental work OR ▪ Haegarda is being used for long-term prophylaxis and member meets one of the following: ○ History of ≥1 attack per month resulting in documented ED admission or hospitalization OR ○ History of laryngeal attacks OR ○ History of ≥2 attacks per month involving the face, throat, or abdomen AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications AND ○ Prescriber acknowledges that the member will receive information and/or counseling regarding the information from the FDA-labeled package insert outlining transmission of infectious agents with a medication made from human blood. Maximum Dose: 60 IU/kg Minimum Age: 6 years CINRYZE (C1 esterase inhibitor - human) may be approved for members meeting the following criteria:

		○ Member has history of trial and failure of Haegarda. Failure is defined as lack of efficacy allergy, intolerable side effects, or a significant drug-drug interaction AND ○ Member has a diagnosis of HAE Type I or Type II confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, C1-INH level) OR has a diagnosis of HAE Type III based on clinical presentation AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member meets at least one of the following: ▪ Cinryze is being used for <u>short-term prophylaxis</u> to undergo a surgical procedure or major dental work OR ▪ Cinryze is being used for <u>long-term prophylaxis</u> and member meets one of the following: ○ History of ≥ 1 attack per month resulting in documented ED admission or hospitalization OR ○ History of laryngeal attacks OR ○ History of ≥ 2 attacks per month involving the face, throat, or abdomen AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications AND ○ Prescriber acknowledges that the member will receive information and/or counseling regarding the information from the FDA-labeled package insert outlining transmission of infectious agents with a medication made from human blood. Minimum age: 6 years Maximum dose: 100 Units/kg ORLADEYO (berotralstat) may be approved for members meeting the following criteria: ○ Member has history of trial and failure of HAEGARDA. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND ○ Member has a diagnosis of HAE Type I or Type II confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, C1-INH level) OR has a diagnosis of HAE Type III based on clinical presentation AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ ORLADEYO is prescribed by or in consultation with an allergist or immunologist AND ○ Appropriate drug interaction interventions will be made for members using concomitant medications that may require dose adjustments (such as
--	--	--

		cyclosporine, fentanyl, pimozide, digoxin) AND ○ Member meets at least one of the following: ▪ ORLADEYO is being used for short-term prophylaxis to undergo a surgical procedure or major dental work ▪ ORLADEYO is being used for long-term prophylaxis and member meets one of the following: • History of ≥ 1 attack per month resulting in documented ED admission or hospitalization OR • History of laryngeal attacks OR • History of ≥ 2 attacks per month involving the face, throat, or abdomen AND • Member is not taking medications that may exacerbate HAE, including ACE inhibitors and estrogen-containing medications Minimum age: 12 years Maximum dose: 150 mg once daily TAKHZYRO (lanadelumab-flyo) may be approved for members meeting the following criteria: ○ Member has history of trial and failure of Haegarda. Failure is defined as: lack of efficacy, allergy, intolerable side effects, or a significant drug-drug interaction AND ○ Member has a diagnosis of HAE Type I or Type II confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, C1-INH level) OR has a diagnosis of HAE Type III based on clinical presentation AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications Minimum age: 2 years Maximum dose: The recommended starting dose is 300mg every 2 weeks. A dosing interval of 300 mg every 4 weeks is also effective and may be considered if the patient is well-controlled (attack free) for more than 6 months <u>Medications Indicated for Treatment of Acute Attacks:</u> Members are restricted to coverage of one medication for <u>treatment of acute attacks</u> at one time. Prior authorization approval will be for one year. FIRAZYR (icatibant acetate) may be approved for members meeting the following criteria: ○ Member has a diagnosis of HAE Type I or Type II confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level,
--	--	---

		C1-INH level) OR has a diagnosis of HAE Type III based on clinical presentation AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications Minimum age: 18 years Maximum dose: 30mg BERINERT (C1 esterase inhibitor - human) may be approved for members meeting the following criteria: ○ Member has a diagnosis of HAE Type I or Type II confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, C1-INH level) OR has a diagnosis of HAE Type III based on clinical presentation AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications AND ○ Prescriber acknowledges that the member will receive information and/or counseling regarding the information from the FDA-labeled package insert outlining transmission of infectious agents with a medication made from human blood. Minimum age: 6 years Max dose: 20 IU/kg RUCONEST (C1 esterase inhibitor - recombinant) may be approved for members meeting the following criteria: ○ Member has a history of trial and failure of Firazyr OR Berinert. Failure is defined as lack of efficacy, allergy, intolerable side effects, or a significant drug-drug interaction AND ○ Member has a diagnosis of HAE Type I or Type II confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, C1-INH level) OR has a diagnosis of HAE Type III based on clinical presentation AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND
--	--	--

		Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications Minimum age: 13 years Maximum dose: 4,200 Units/dose All other non-preferred agents may be approved if the member has trialed and failed at least two preferred agents with the same indicated role in therapy as the prescribed medication (prophylaxis or treatment). Failure is defined as lack of efficacy, allergy, intolerable side effects, or a significant drug-drug interaction.
Therapeutic Drug Class: PHOSPHATE BINDERS – Effective 10/1/2025		
No PA Required Calcium acetate capsule PHOSLYRA (calcium acetate) solution Sevelamer carbonate tablet, powder pack	PA Required AURYXIA (ferric citrate) tablet Calcium acetate tablet CALPHRON (calcium acetate) tablet Ferric citrate tablet FOSRENOL (lanthanum carbonate) chewable tablet, powder pack Lanthanum carbonate chewable tablet RENVELA (sevelamer carbonate) powder pack, tablet Sevelamer HCl tablet VELPHORO (sucroferric oxide) chewable tablet XPHOZAH (tenapanor) tablet	Prior authorization for non-preferred products in this class may be approved if member meets all the following criteria: - Member has diagnosis of end stage renal disease AND - Member has elevated serum phosphorus [> 4.5 mg/dL or > 1.46 mmol/L] AND - Provider attests to member avoidance of high phosphate containing foods from diet AND - Member has trialed and failed‡ one preferred agent (lanthanum products require trial and failure‡ of a preferred sevelamer product). Auryxia (ferric citrate) may be approved if the member meets all the following criteria: - Member is diagnosed with end-stage renal disease, receiving dialysis, and has elevated serum phosphate (> 4.5 mg/dL or > 1.46 mmol/L). AND - Provider attests to counseling member regarding avoiding high phosphate containing foods from diet AND - Member has trialed and failed‡ three preferred agents with different mechanisms of action prescribed for hyperphosphatemia in end stage renal disease OR - Member is diagnosed with chronic kidney disease with iron deficiency anemia and is not receiving dialysis AND - Member has tried and failed‡ at least two different iron supplement product formulations (OTC or RX) Velphoro (sucroferric oxyhydroxide tablet, chewable) may be approved if the member meets all of the following criteria: - Member is diagnosed with chronic kidney disease and receiving dialysis and has elevated serum phosphate (> 4.5 mg/dL or > 1.46 mmol/L). AND - Provider attests to counseling member regarding avoiding high phosphate containing foods from diet AND - Member has trialed and failed‡ two preferred agents, one of which must be a preferred sevelamer product Maximum Dose: Velphoro 3000mg daily

		Members currently stabilized on a non-preferred lanthanum product may receive approval to continue therapy with that product. ‡Failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction. <i>Note: Medications administered in a dialysis unit or clinic are billed through the Health First Colorado medical benefit or Medicare with members with dual eligibility.</i>
--	--	--

Therapeutic Drug Class: **PRENATAL VITAMINS / MINERALS** – *Effective 10/1/2025*

Preferred *Must meet eligibility criteria	Non-Preferred PA Required	
COMPLETE NATAL DHA pack M-NATAL PLUS tablet NESTABS tablets PRENATAL VITAMIN PLUS LOW IRON tablet (Patrin Pharma only) SE-NATAL 19 chewable tablet^{BNR} TARON-C DHA capsule THRIVITE RX tablet TRINATAL RX 1 tablet VITAFOL gummies WESNATAL DHA COMPLETE tablet WESTAB PLUS tablet	All other rebateable prescription products are non-preferred	*Preferred and non-preferred prenatal vitamin products are a benefit for members from 11-60 years of age who are pregnant, lactating, or trying to become pregnant. Prior authorization for non-preferred agents may be approved if member fails 7-day trial with four preferred agents. Failure is defined as allergy, intolerable side effects, or significant drug-drug interaction.

XI. Ophthalmic

Therapeutic Drug Class: **OPHTHALMIC, ALLERGY** – *Effective 4/1/2025*

No PA Required	PA Required	
ALREX ^{BNR} (loteprednol) 0.2%	ALAWAY (ketotifen) 0.025% (OTC)	Non-preferred products may be approved following trial and failure of therapy with two preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions).
Azelastine 0.05%	ALOCRIIL (nedocromil) 2%	
Cromolyn 4%	ALOMIDE (lodoxamide) 0.1%	
Ketotifen 0.025% (OTC)	Bepotastine 1.5%	
LASTACRAFT (alcaftadine) 0.25% (OTC)	BEPREVE (bepotastine) 1.5%	
Olopatadine 0.1%, 0.2% (OTC) (generic Pataday Once/Twice Daily)	Epinastine 0.05%	
	Loteprednol 0.2%	
	Olopatadine 0.1%, 0.2% (RX)	
	PATADAY ONCE DAILY (olopatadine) 0.2% (OTC)	
	PATADAY TWICE DAILY (olopatadine) 0.1% (OTC)	
	PATADAY XS ONCE DAILY (olopatadine) 0.7% (OTC)	
	ZADITOR (ketotifen) 0.025% (OTC)	
	ZERVIAE (cetirizine) 0.24%	

Therapeutic Drug Class: **OPHTHALMIC, IMMUNOMODULATORS** – *Effective 4/1/2025*

No PA Required	PA Required	
RESTASIS ^{BNR} (cyclosporine 0.05%) vials	CEQUA (cyclosporine) 0.09% solution	Non-preferred products may be approved for members meeting all of the following criteria: • Member is 18 years and older AND • Member has a diagnosis of chronic dry eye AND • Member has failed a 3-month trial of one preferred product. Failure is defined as a lack of efficacy, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions AND
	Cyclosporine 0.05% vials	
	MIEBO (Perfluorohexyloctane/PF)	
	RESTASIS MULTIDOSE (cyclosporine) 0.05%	

	TYRVAYA (varenicline) nasal spray VERKAZIA (cyclosporin emulsion) VEVYE (cyclosporine) 0.1% XIIDRA (lifitegrast) 5% solution	- Prescriber is an ophthalmologist, optometrist or rheumatologist <u>Maximum Dose/Quantity:</u> 60 single use containers for 30 days 5.5 mL/20 days for Restasis Multi-Dose and Vevye 3mL/30 days for Miebo Verkazia (cyclosporine ophthalmic emulsion) may be approved if the following criteria are met: - Member is ≥ 4 years of age AND - Verkazia is being used for the treatment of vernal keratoconjunctivitis (VKC) AND - Member has trialed and failed therapy with three agents from the following pharmacologic categories: preferred dual-acting mast cell stabilizer/antihistamine from the Ophthalmics-Allergy PDL class, oral antihistamine, preferred topical ophthalmic corticosteroid from the Ophthalmics-Anti-inflammatories PDL class. Failure is defined as lack of efficacy with 2-week trial, allergy, contraindication to therapy, intolerable side effects, or significant drug-drug interaction <u>Quantity limit:</u> 120 single-dose 0.3 mL vials/15 days
Therapeutic Drug Class: OPHTHALMIC, ANTI-INFLAMMATORIES – <i>Effective 4/1/2025</i>		
NSAIDs		Durezol (difluprednate) may be approved if meeting the following criteria: - Member has a diagnosis of severe intermediate uveitis, severe panuveitis, or severe uveitis with the complication of uveitic macular edema AND has trialed and failed prednisolone acetate 1% (failure is defined as lack of efficacy, allergy, contraindication to therapy, intolerable side effects, or significant drug-drug interaction) OR - Members with a diagnosis other than those listed above require trial and failure of three preferred agents (failure is defined as lack of efficacy, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction). Eysuvis (loteprednol etabonate) may be approved if meeting all of the following: - Member is ≥ 18 years of age AND - Eysuvis (loteprednol etabonate) is being used for short-term treatment (up to two weeks) of the signs and symptoms of dry eye disease AND - Member has failed treatment with one preferred product in the Ophthalmic Immunomodulator therapeutic class. Failure is defined as lack of efficacy with a
No PA Required	PA Required	
Diclofenac 0.1%	ACULAR (ketorolac) 0.5%, LS 0.4%	
Flurbiprofen 0.03%	ACUVAIL (ketorolac/PF) 0.45%	
Ketorolac 0.5%, Ketorolac LS 0.4%	Bromfenac 0.07%, 0.075%, 0.09%	
NEVANAC (nepafenac) 0.1%	BROMSITE (bromfenac) 0.075%	
	ILEVRO (nepafenac) 0.03%	
	PROLENSA (bromfenac) 0.07%	
Corticosteroids		
No PA Required	PA Required	
FLAREX (fluorometholone) 0.1%	Dexamethasone 0.1%	
	Difluprednate 0.05%	

Fluorometholone 0.1% drops FML FORTE (fluorometholone) 0.25% drops LOTEMAX^{BNR} (loteprednol) 0.5% drops, gel LOTEMAX (loteprednol) 0.5% ointment MAXIDEX (dexamethasone) 0.1% PRED MILD (prednisolone) 0.12% Prednisolone acetate 1%	DUREZOL (difluprednate) 0.05% EYSUVIS (loteprednol) 0.25% FML LIQUIFILM (fluorometholone) 0.1% drop FML S.O.P (fluorometholone) 0.1% ointment INVELTYS (loteprednol) 1% LOTEMAX SM (loteprednol) 0.38% gel Loteprednol 0.5% drops, 0.5% gel PRED FORTE (prednisolone) 1% Prednisolone sodium phosphate 1%	3-month trial, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction) AND - Member does not have any of the following conditions: - Viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella OR - Mycobacterial infection of the eye and fungal diseases of ocular structures <u>Quantity limit</u>: one bottle/15 days Lotemax SM (loteprednol etabonate) or Inveltys (loteprednol etabonate) may be approved if meeting all of the following: - Member is ≥ 18 years of age AND - Lotemax SM or Inveltys (loteprednol etabonate) is being used for the treatment of post-operative inflammation and pain following ocular surgery AND - Member has trialed and failed therapy with two preferred loteprednol formulations (failure is defined as lack of efficacy with 2-week trial, allergy, contraindication to therapy, intolerable side effects, or significant drug-drug interaction) AND - Member has trialed and failed therapy with two preferred agents that do not contain loteprednol (failure is defined as lack of efficacy with 2-week trial, contraindication to therapy, allergy, intolerable side effects, or significant drug-drug interaction) AND - Member does not have any of the following conditions: - Viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella OR - Mycobacterial infection of the eye and fungal diseases of ocular structures All other non-preferred products may be approved with trial and failure of three preferred agents (failure is defined as lack of efficacy with 2-week trial, allergy, contraindication, intolerable side effects, or significant drug-drug interaction).
Therapeutic Drug Class: OPHTHALMIC, GLAUCOMA – <i>Effective 4/1/2025</i>		
Beta-blockers		
No PA Required	PA Required	
Carteolol 1% Levobunolol 0.5% Timolol (generic Timoptic) 0.25%, 0.5%	Betaxolol 0.5% BETIMOL (timolol) 0.25%, 0.5% BETOPIC-S (betaxolol) 0.25%	Non-preferred products may be approved following trial and failure of therapy with three preferred products, including one trial with a preferred product having the same general mechanism (such as prostaglandin analogue, alpha2-adrenergic agonist, beta-blocking agent, or carbonic anhydrase inhibitor). Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions.

	ISTALOL (timolol) 0.5% Timolol (generic Istalol) 0.5% drops Timolol GFS 0.25%, 0.5% Timolol/PF (generic Timoptic Ocudose) 0.25%, 0.5% TIMOPTIC, TIMOPTIC OCUDOSE (timolol) 0.25%, 0.5% TIMOPTIC-XE (timolol GFS) 0.25%, 0.5%	Non-preferred combination products may be approved following trial and failure of therapy with one preferred combination product AND trial and failure of individual products with the same active ingredients as the combination product being requested (if available) to establish tolerance. Failure is defined as lack of efficacy with 4-week trial, allergy, intolerable side effects or significant drug-drug interactions. Preservative free products may be approved with provider documentation of adverse effect to preservative-containing product.
Carbonic anhydrase inhibitors		
No PA Required	PA Required	
Brinzolamide 1%	AZOPT (brinzolamide) 1%	
Dorzolamide 2%		
Prostaglandin analogue		
No PA Required	PA Required	
Latanoprost 0.005%	Bimatoprost 0.03%	
LUMIGAN ^{BNR} (bimatoprost) 0.01%	IYUZEH (latanoprost/PF) 0.005%	
TRAVATAN Z ^{BNR} (travoprost) 0.004%	Tafluprost 0.0015%	
	Tafluprost PF 0.0015%	
	Travoprost 0.004%	
	VYZULTA (latanoprostene) 0.024%	
	XALATAN (latanoprost) 0.005%	
	XELPROS (latanoprost) 0.005%	
	ZIOPTAN (tafluprost PF) 0.0015%	
Alpha-2 adrenergic agonists		

No PA Required	PA Required	
ALPHAGAN P ^{BNR} 0.1%, 0.15% (brimonidine) Brimonidine 0.2%	Apraclonidine 0.5% Brimonidine 0.1%, 0.15% IOPIDINE (apraclonidine) 0.5%, 1%	
Other ophthalmic, glaucoma and combinations		
No PA Required	PA Required	
COMBIGAN ^{BNR} 0.2%-0.5% (brimonidine/timolol) Dorzolamide/Timolol 2%-0.5% RHOPRESSA (netarsudil) 0.02% ROCKLATAN (netarsudil/latanoprost) 0.02%-0.005%	Brimonidine/Timolol 0.2%-0.5% COSOPT/COSOPT PF (dorzolamide/timolol) 2%-0.5% Dorzolamide/Timolol PF 2%-0.5% PHOSPHOLINE IODIDE (echothiophate) 0.125% Pilocarpine 1%, 1.25%, 2%, 4% SIMBRINZA (brinzolamide/brimonidine) 1%-0.2% Vuity (pilocarpine) 1.25%	

XII. Renal/Genitourinary

Therapeutic Drug Class: **BENIGN PROSTATIC HYPERPLASIA (BPH) AGENTS** – *Effective 10/1/2025*

No PA Required	PA Required	*CIALIS (tadalafil) may be approved if meeting the following criteria:
Alfuzosin ER tablet	AVODART (dutasteride) softgel	- Member has a documented diagnosis of BPH AND - Member has trialed and failed each of the following: - Finasteride. Failure is defined as lack of efficacy with a 3-month trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction AND - Either a nonselective alpha blocker or tamsulosin. Failure is defined as lack of efficacy, allergy, intolerable side effects, contraindication, or significant drug-drug interaction AND - Documentation of BPH diagnosis will require BOTH of the following: - AUA Prostate Symptom Score ≥ 8 AND - Results of a digital rectal exam AND
Doxazosin tablet	CARDURA (doxazosin) tablet	
Dutasteride capsule	CARDURA XL (doxazosin ER) tablet	
Finasteride tablet	*CIALIS (tadalafil) 2.5 mg, 5 mg tablet	
Tamsulosin capsule	Dutasteride/tamsulosin capsule	

Terazosin capsule	Finasteride/tadalafil capsule FLOMAX (tamsulosin) capsule PROSCAR (finasteride) tablet RAPAFLO (silodosin) capsule Silodosin capsule *Tadalafil 2.5 mg, 5 mg tablet Tezruly (terazosin) solution	- Cialis (tadalafil) is not being prescribed for use for continuing alpha blocker therapy, as use of tadalafil in this population is not recommended due to the potential for hypotension. <u>Maximum Dose:</u> Doses exceeding Cialis (tadalafil) 5mg per day will not be approved. Prior authorization for all other non-preferred products may be approved if meeting the following criteria: - Member has tried and failed‡ three preferred agents AND - For combinations agents, member has tried and failed‡ each of the individual agents within the combination agent and one other preferred agent. ‡Failure is defined as lack of efficacy with 8-week trial, allergy, intolerable side effects, contraindication, or significant drug-drug interaction.
Therapeutic Drug Class: ANTI-HYPERURICEMICS – <i>Effective 10/1/2025</i>		
No PA Required Allopurinol 100 mg, 300 mg tablets Colchicine tablet Febuxostat tablet Probenecid tablet Probenecid/Colchicine tablet	PA Required Allopurinol 200 mg tablets Colchicine capsule COLCRYS (colchicine) tablet GLOPERBA (colchicine) oral solution MITIGARE (colchicine) capsule ULORIC (febuxostat) tablet	Non-preferred xanthine oxidase inhibitor products (allopurinol or febuxostat formulations) may be approved following trial and failure of preferred allopurinol. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. If member has tested positive for the HLA-B*58:01 allele, it is not recommended that they trial allopurinol. A positive result on this genetic test will count as a failure of allopurinol. Prior authorization for all other non-preferred agents (non-xanthine oxidase inhibitors) may be approved after trial and failure of two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. GLOPERBA (colchicine) oral solution may be approved for members who require individual doses <0.6 mg OR for members who are unable to use a solid oral dosage form. Colchicine tablet quantity limits: - Chronic hyperuricemia/gout prophylaxis: 60 tablets per 30 days - Familial Mediterranean Fever: 120 tablets per 30 days
Therapeutic Drug Class: OVERACTIVE BLADDER AGENTS – <i>Effective 10/1/2025</i>		
No PA Required Fesoterodine ER tablet MYRBETRIQ (mirabegron) tablet ^{BNR} Oxybutynin IR, ER tablets, syrup	PA Required Darifenacin ER tablet DETROL (tolterodine) tablet DETROL LA (tolterodine) ER capsule	Non-preferred products may be approved for members who have failed treatment with two preferred products. Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.

Solifenacin tablet	Flavoxate tablet	
Tolterodine tablet, ER capsule	GEMTESA (vibegron) tablet	
Trospium ER tablet	Mirabegron tablet	
	MYRBETRIQ (mirabegron) suspension	
	Oxybutynin 2.5 mg tablet	
	OXYTROL (oxybutynin patch)	
	TOVIAZ (Fesoterodine ER) tablet	
	Trospium ER capsule	
	VESICARE (solifenacin) tablet	
	VESICARE LS (solifenacin) suspension	

XIII. RESPIRATORY

Therapeutic Drug Class: **RESPIRATORY AGENTS** – *Effective 4/14/2025*

Inhaled Anticholinergics

Preferred No PA Required (Unless indicated*)	Non-Preferred PA Required	
<u>Solutions</u> Ipratropium solution	<u>Solutions</u> YUPELRI (revefenacin) solution	*SPIRIVA RESPIMAT (tiotropium) 1.25 mcg may be approved for members ≥ 6 years of age with a diagnosis of asthma (qualifying diagnosis verified by AutoPA). SPIRIVA RESPIMAT is intended to be used by members whose asthma is not controlled with regular use of a combination medium-dose inhaled corticosteroid and long-acting beta agonist (LABA).
<u>Short-Acting Inhalation Devices</u> ATROVENT HFA (ipratropium)	<u>Short-Acting Inhalation Devices</u>	*SPIRIVA RESPIMAT (tiotropium) 2.5 mcg may be approved for members with a diagnosis of COPD who have trialed and failed SPIRIVA HANDIHALER. Failure is defined as intolerable side effects or inability to use dry powder inhaler (DPI) formulation.
<u>Long-Acting Inhalation Devices</u> SPIRIVA Handihaler ^{BNR} (tiotropium) *SPIRIVA RESPIMAT (tiotropium)	<u>Long-Acting Inhalation Devices</u> INCRUSE ELLIPTA (umeclidinium) Tiotropium DPI TUDORZA PRESSAIR (aclidinium)	LONHALA MAGNAIR (glycopyrrolate) may be approved for members ≥ 18 years of age with a diagnosis of COPD including chronic bronchitis and emphysema who have trialed and failed‡ treatment with two preferred anticholinergic agents. Non-preferred single agent anticholinergic agents may be approved for members with a diagnosis of COPD including chronic bronchitis and/or emphysema who have trialed and failed‡ treatment with two preferred agents, one of which must be SPIRIVA HANDIHALER.

		‡Failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction.
Inhaled Anticholinergic Combinations		
No PA Required <u>Solutions</u> Ipratropium/Albuterol solution <u>Short-Acting Inhalation Devices</u> COMBIVENT RESPIMAT (albuterol/ipratropium) <u>Long-Acting Inhalation Devices</u> ANORO ELLIPTA (umeclidinium/vilanterol) ^{BNR}	PA Required <u>Solutions</u> <u>Short-Acting Inhalation Devices</u> <u>Long-Acting Inhalation Devices</u> BEVESPI AEROSPHERE (glycopyrrolate /formoterol fumarate) BREZTRI AEROSPHERE (budesonide/glycopyrrolate/ formoterol) DUAKLIR PRESSAIR (aclidinium/formoterol) STIOLTO RESPIMAT (tiotropium/olodaterol) Umeclidinium/Vilanterol	BREZTRI AEROSPHERE (budesonide/glycopyrrolate/formoterol) may be approved for members ≥ 18 years of age with a diagnosis of COPD who have trialed and failed‡ treatment with two preferred anticholinergic-containing agents. DUAKLIR PRESSAIR (aclidinium/formoterol) may be approved for members ≥ 18 years of age with a diagnosis of COPD who have trialed and failed‡ treatment with two preferred anticholinergic-containing agents. All other non-preferred inhaled anticholinergic combination agents may be approved for members with a diagnosis of COPD including chronic bronchitis and/or emphysema who have trialed and failed‡ treatment with two preferred inhaled anticholinergic combination agents OR three preferred inhaled anticholinergic-containing agents (single ingredient or combination). Members who are currently stabilized on Bevespi Aerosphere may receive approval to continue therapy with that product. ‡Failure is defined as lack of efficacy with 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction.
Inhaled Beta2 Agonists (short acting)		
No PA Required <u>Solutions</u> Albuterol solution, for nebulizer <u>Inhalers</u> VENTOLIN ^{BNR} HFA (albuterol)	PA Required <u>Solutions</u> Levalbuterol solution <u>Inhalers</u> AIRSUPRA (budesonide/albuterol) Albuterol HFA Levalbuterol HFA PROAIR RESPICLICK (albuterol) XOPENEX (levalbuterol) Inhaler	Non-preferred short acting beta-2 agonists may be approved for members who have failed treatment with one preferred agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. MDI formulation quantity limits: 2 inhalers / 30 days AIRSUPRA (budesonide/albuterol) <u>Airsupra minimum age:</u> 18 years old
Inhaled Beta2 Agonists (long acting)		

Preferred <u>Solutions</u> <u>Inhalers</u> SEREVENT DISKUS (salmeterol) inhaler	Non-Preferred PA Required <u>Solutions</u> Arformoterol solution BROVANA (arformoterol) solution Formoterol solution PERFOROMIST (formoterol) solution <u>Inhalers</u> STRIVERDI RESPIMAT (olodaterol)	Non-preferred agents may be approved for members with moderate to severe COPD, AND members must have failed a trial of Serevent. Failure is defined as lack of efficacy with a 6-week trial, allergy, intolerable side effects, or significant drug-drug interaction. For treatment of members with diagnosis of asthma needing add-on therapy, please refer to preferred agents in combination Long-Acting Beta Agonist/Inhaled Corticosteroid therapeutic class.
Inhaled Corticosteroids		
No PA Required <u>Solutions</u> Budesonide nebulers <u>Inhalers</u> ARNUITY ELLIPTA^{BNR} (fluticasone furoate) ASMANEX HFA (mometasone furoate) inhaler ASMANEX Twisthaler (mometasone) PULMICORT FLEXHALER (budesonide) QVAR REDHALER (beclomethasone)	PA Required <u>Solutions</u> PULMICORT (budesonide) respules <u>Inhalers</u> ALVESCO (ciclesonide) inhaler Fluticasone Ellipta Fluticasone propionate diskus *Fluticasone propionate HFA	Non-preferred inhaled corticosteroids may be approved in members with asthma who have failed an adequate trial of two preferred agents. An adequate trial is defined as at least 6 weeks. (Failure is defined as: lack of efficacy with a 6-week trial, allergy, contraindication to, intolerable side effects, or significant drug-drug interactions, or dexterity/coordination limitations (per provider notes) that significantly impact appropriate use of a specific dosage form.) *FLUTICASONE PROPIONATE HFA is available to members without prior authorization for: - Members with a diagnosis of eosinophilic esophagitis (EoE) OR - Members ≤ 12 years of age. <u>Maximum Dose:</u> Pulmicort (budesonide) nebulizer suspension: 2mg/day <u>Quantity Limits:</u> Pulmicort flexhaler: 2 inhalers / 30 days
Inhaled Corticosteroid Combinations		
No PA Required (*Must meet eligibility criteria) ADVAIR DISKUS^{BNR} (fluticasone/salmeterol) ADVAIR HFA^{BNR} (fluticasone/salmeterol)	PA Required BREO ELLIPTA (vilanterol/fluticasone furoate) Budesonide/formoterol (generic Symbicort) Fluticasone/salmeterol (generic Airduo/Advair Diskus)	*TRELEGY ELLIPTA (fluticasone furoate/umeclidinium/vilanterol) may be approved if the member has trialed/failed one preferred agent. Failure is defined as lack of efficacy with a 6-week trial, allergy, intolerable side effects, significant drug-drug interactions, or dexterity/coordination limitations (per provider notes) that significantly impact appropriate use of a specific dosage form. Non-preferred inhaled corticosteroid combinations may be approved for members meeting both of the following criteria:

AIRDUO RESPICLICK ^{BNR} (fluticasone/salmeterol) DULERA (mometasone/formoterol) SYMBICORT ^{BNR} (budesonide/formoterol) inhaler *TRELEGY ELLIPTA (fluticasone furoate/ umeclidinium/vilanterol)	Fluticasone/salmeterol HFA (generic Advair HFA) Fluticasone/vilanterol (generic Breo Ellipta) WIXELA INHUB (fluticasone/salmeterol)	- Member has a qualifying diagnosis of asthma or severe COPD; AND - Member has failed two preferred agents (Failure is defined as lack of efficacy with a 6-week trial, allergy, intolerable side effects, significant drug-drug interactions, or dexterity/coordination limitations (per provider notes) that significantly impact appropriate use of a specific dosage form.
Phosphodiesterase Inhibitors (PDEIs)		
No PA Required Roflumilast tablet	PA Required DALIRESP (roflumilast) tablet OHTUVAYRE (ensifentrine) suspension	Requests for use of the non-preferred brand product formulation may be approved if meeting criteria outlined in the Appendix P “Generic Mandate” section.